Rational Drug Design for Oncotherapy Using Molecular Dynamics Simulations

Jack Tuszynski
Allard Chair
Cross Cancer Institute
Edmonton, Canada

The Tuszynski Group of Molecular Biophysics at the University of Alberta -> CCI

- **ØJack Tuszynski, professor**
- **ØTorin Huzil, PDF**
- **ØAvner Priel, Research associate**
- ØEric Carpenter, senior research assistant
- **Ø**Tyler Luchko, Ph.D. student
- **ØRebeccah Marsh, Ph.D. student**
- **OCatherine Beauchemin, Ph.D. student**
- **ØEvan Kelly, PhD. Student (co-supervised)**
- **ØAllan MacDairmid, Ph.D. student (co-supervised)**
- ØRobert Bryce, Ph.D. student (co-supervised)
- **ØKristy Beinert, summer student**
- Matthew Bergstrom, summer student
- **ØJoe Hajduk, summer student**
- **Ø**Achin Gupta, summer students

Acknowledgements

Collaborators:

- D. Sept, Washington University, St. Louis, USA
- R. Luduena, UTHSC San Antonio, USA
- A. Weis, Lipitek Inc., San Antonio, USA
- E. Unger, IMB Jena, Germany
- N. Woolf, UCLA, USA
- H. Cantiello, Harvard Med School, USA
- J. Dixon, Univ. of Warwick, UK
- M. Peyrard, ENS Lyon, France
- Y. Engelborghs, KU Leuven, Belgium
- H. Bolterauer, Univ. of Giessen, Germany
- M. Sataric, Univ Novi Sad, Serbia
- A. Kovalenko, NINT, Canada

<u>Support</u>: Allard Foundation, MITACS, NSERC, US Dept of Defense, Technology Innovations, CPCRI, YeTaDel Foundation, PIMS, Oncovista, Leverhulme Trust UK



OA drug has to bind to its target with a suitable affinity

OThe drug must stimulate or inhibit the activity of the target

OHowever, a good fit and high affinity may not always be the best scenario.

Modification of key interactions may also be sufficient for increased drug activity

Lead compounds from natural sources

- Finding new and effective drugs is a difficult process
- Many are discovered by serendipity
- The systematic screening of potential drugs involves thousands of compounds
- Computational searches offer speedup and cost cuts

How can computers help design better cancer drugs?


- OEnables us to visualize proteins in 3+1 dimensions
- OManipulation of these systems in silico including diffusion calculations
- ODocking fits drug molecules into binding sites and determines the energy of multiple conformations
- OStatistical Mechanics permits molecular motion and solvent effects to be incorporated
- OSearching through chemical and structural databases to uncover novel correlations

Quantitative Structure Activity Relationship (QSAR)

OStructure Activity Relationship that is applied with a degree of predictive ability

OStructural and chemical elements, such as size, hydrophobicity and the electrostatics of a drug are quantitatively scored and used to predict the activity of similar compounds

Spatial Fitting



OVisual inspection of binding sites in a target

OModification of the drug to provide better binding to target


Olnvolves manual or automated docking of a drug into binding site

Microtubules

OMicrotubules are composed of tubulin

OAssembly and disassembly rates are critical for cell function

ODisruption of these rates with drugs like colchicine and taxol results in cell death




MICROTUBULES PLAY CRITICAL ROLES IN:


Mitosis Intracellular transport Intracellular organization Motility









Ciliated Cell



Dividing Cell




Nerve Cell




Q tQ tQ tQ tr


■ North

Microtubules in Mitosis






GTP Lateral Cap Model for MT Dynamic Instability



Individual MT life story: dynamic instability



- •Rescues
- Growth phase
- Shrinking phase



Simulation using a recursive map Shown in the next slide

Modeling of Individual MT Assembly Processes:

Recursive Map

States:

- Attachment
- Detachment
- Pause
- Catastrophe

$\ell(t+1) = (1 - r_3(1 - r_1)(1 - r_2))(\ell(t) + r_1a - r_2a - ar_1r_3 + ar_2r_3)$

Master Equations

$$\frac{d}{dt}P(n,t) = c_T t_{r1} P(n-1,t) + t_{r2} P(n+1,t) - (c_T t_{r1} + t_{r2} + t_{r3}) P(n,t)$$

for n > 0, and

$$\frac{d}{dt}P(0,t) = t_{r2}P(1,t) + t_{r3}\sum_{n=1}^{\infty}P(n,t) - c_T t_{r1}P(0,t)$$

for n=0.

continuum limit

integro-differential equation


$$\frac{\partial P}{\partial t} = \frac{1}{2}(c_T t_{r1} + t_{r2})\frac{\partial^2 P}{\partial x^2} - (c_T t_{r1} - t_{r2})\frac{\partial P}{\partial x} - t_{r3}P + t_{r3}\int_x^\infty w(x' - x)P(x')dx'$$

Microtubules and Cancer

O Cancer is related to a failure of multiple cellular checkpoints that results in uncontrolled division

O we are selectively targeting mitosis during cell division


Oa critical component of mitosis are the microtubules



MICROTUBULES AND TUBULIN

- Microns-long cylindrical organelles 25 nm in diameter
- Associated with MAPs and motor proteins
- Formed from 110 kD tubulin α/β heterodimer
- Tubulin dimers bind end to end to form a protofilament; 13 protofilaments arranged side to side to form microtubule
- Both α and β tubulins exist as numerous isotypes each encoded by different genes


The atomic resolution structure of the tubulin dimer following Downing's electron crystallography (1997)




C-terminus not resolved

PDB entries for tubulin: 1TUB, 1JFF, 1FFX, 1SAO

Molecular Dynamics




Tubulin Structure: key motifs



Charge distribution of tubulin dimer including C-termini.


Blue=negative Red=positive White=neutral




Approx. 25 e per monomer or 50 per dimer, 40% on C-termini

Electric Field around tubulin

- Because each atom has a partial charge we can calculate the electric field of the protein. $\frac{1}{4\pi\epsilon_0} \sum_{i=1}^n \frac{q_i}{r_i^2} \hat{r}_i$
- Bases (e⁺) and acids (e⁻) are colored blue and red respectively.
- Iso-surface of the electric field is coloured yellow.



Electric Potential




- slice through dimer
- red/blue=negative/positive
- green=isopotential lines


Poisson-Boltzmann equation:

$$\nabla \cdot (\epsilon(\vec{x}) \nabla \phi(\vec{x})) - \kappa(\vec{x})^2 \sinh(\phi(\vec{x})) = -4\pi \rho(\vec{x})$$

Monopole in vacuum becomes a dipole in solvent




Potential in vacuum




potential in solvent

Conformational effects of GTP hydrolysis in a tubulin dimer



Map of Tubulin Dimer's Sphere of Attraction for a neighbouring dimer



Map of the volume in which two tubulin dimers of identical orientation attract. The magenta "tufts" indicate locations of the centre of mass of a second dimer when it would be attracted by the central dimer and the direction of the force. Darker "tufts" are near the viewer, lighter ones further away.


Brownian Dynamics as a tool for MT polymerization simulations:


Model protein-protein association using BD methods


- Electrostatic interactions
- Solvent interactions

$$\Delta R = \frac{D\Delta t}{k_B T} F + S$$

Displacement of Protein









Brownian Dynamics Simulation

- Start at 120 A separation (b)
- End upon escape > 500 A (q) or binding
- From fraction of successful trajectories (β) we can calculate a rate constant:






Conformational dynamics of protofilament formation




Stathmin-stabilized MT filament




PDB file: 1FFX

Conclusion: GTP hydrolysis curves protofilaments


Longitudinal Bonds (stronger) in a Protofilament



Lateral Bonds: formation of a sheet




Lattice formation: A or B type lattice




Energy of a lateral tubulin-tubulin interaction: k=4 N/m D. Sept Consistent with the elastic properties of MT's (Young's modulus of 109 N/m²)

Blue= α tubulin Red= β tubulin

Mircotubule lattices




A Lattice




B Lattice

Taxol diffuses through nanometersize surface pores in a microtubule




Outside surface



Inside surface

Charge distribution on a microtubule.




Mammalian β-Tubulin Isotypes

Isotype	Distribution
βΙ	Most cells
βII	Neurons, glia, inner ear, skeletal
	muscle, smooth muscle
βIII	Neurons, Sertoli cells, colon
βIVa	Neurons and glia
βIVb	Most tissues
βV	3T3 and CHO cells
βVI	Platelets, marrow, spleen
βVII	Brain

β-Tubulin Isotypes in Mammals

	% Sequence	% Conservation
	Identity to βI	<u>in Evolution</u>
βΙ	100.0	99.5
βII	94.6	98.9
βIII	90.2	99.6
βIVa	96.0	N/A
βIVb	97.3	96.6
βV	91.4	98.3
βVI	76.7	80.9
βVII	85.5	ND


Dynamics of Isotypically Pure Microtubules



Panda et al. (1994) *Proc. Nat. Acad. Sci. USA* 91,

11358-11362


Slight shape variations



Mary Ann Jordan & Leslie Wilson:

- Mitotic-spindle microtubules are among the most successful targets for anticancer therapy. Microtubule-targeted drugs, including paclitaxel and Vinca alkaloids, suppress microtubule dynamics without changing microtubule mass; leading to mitotic block and apoptosis. Some microtubule-targeted drugs can act as vascular-targeting agents, rapidly depolymerizing microtubules of newly formed vasculature to shut down the blood supply to tumours.
- Microtubule-active drugs bind to one of three main classes of sites on tubulin, the paclitaxel site, the Vinca domain and the colchicine domain. Drugs that bind to the colchicine domain undergo intensive investigation as vascular-targeting agents for cancer therapy.
- Development of resistance to microtubule-targeted drugs has several possible causes, some might involve changes in microtubule dynamics resulting from altered expression of tubulin isotypes, tubulin mutations, and altered expression or binding of microtubule-regulatory proteins.
- Microtubule-targeted drugs can synergize with one another.
- Understanding their modes of action might lead to improved dosing regimens and combinations with other microtubule-targeted drugs, as well as combinations with 'molecularly targeted' drugs.
- "Conceivably, if one knew the tubulin isotype and microtubule regulatory protein composition of a specific tumor cell, one could design or choose drugs to selectively target that tumor"


Taxol, colchicine and vinca bound to a microtubule




LIGAND BINDING SITES ON TUBULIN

- 1) Exchangeable GTP site (β) regulates assembly and dynamic instability.
- 2) Non-exchangeable GTP site (α). Function unknown.
- 3) Colchicine site (α/β interface). Ligands inhibit MT assembly (colchicine, podophyllotoxin, TN16).
- 4) Vinca area (β). Ligands inhibit MT assembly.
- 5) Taxol site (β) . Ligands promote MT assembly, inhibit dynamics.
- 6) Other sites: FHIT, Mg++,


Binding pockets for colchicine and taxol compounds




colchicine




docetaxel



paclitaxel






Structure Activity Relationship

OA means by which drugs can be related to their molecular structure

OAssessed by considering a series of modifications to a drug

OThe effect upon biological activity is then determined for each modification

Colchicine binding pocket view from alpha tubulin towards beta




Tubulin residues involved in interactions with Colchicine

class II class III class III class VI class II
class III
class VI


RFPGQLNADLRKLAVNMVPFPRLHFFMPGFAPLTSRGSQQYRALTVPELTQQVFDAKNMM 300 RFPGQLNADLRKLAVNMVPFPRLHFFMPGFAPLTSRGSQQYRALTVPELTQQMFDSKNMM 300 RFPGQLNADLRKLAVNMVPFPRLHFFMPGFAPLTARGSQQYRALTVPELTQQMFDAKNMM 300 RFPGQLNADLRKLAVNMVPFPRLHFFMPGFAPLTAQGSQQYRALSVAELTQQMFDARNTM 300

class II class III class VI AACDPRHGRYLTVAAVFRGRMSMKEVDEQMLNVQNKNSSYFVEWIPNNVKTAVCDIPPRG 360
AACDPRHGRYLTVAAIFRGRMS/KEVDEQMLNVQNKNSSYFVEWIPNNVKTA/CDIPPRG 360
AACDPRHGRYLTVATVFRGRI DEQMLAIQSKNSSYFVEWIPNNVKY PRG 360
AACDLRRGRYLTVACIFRGY VDQQLLSVQTRNSSCFVEWIPNNVKVAVCDIPPRG 360

The colchicine binding site βII and βIII Tubulin




Thr (polar)-> Val (no)
Cys (polar)->Ser (polar)





Tyr (polar)->Arg (+charged)

Colchicine bound to β III and β III Tubulin





Colchicine derivatives that may differentiate isotypes







Colchicine derivatives bound to Tubulin isotypes








Structure-activity relationship of taxol



Drug binding example: Taxol bound to Tubulin







Taxol in its binding site

top view and at a 45 deg angle (bottom). The white bump in the centre of the pocket is the ring of His 227 viewed edge-on;


the taxoid is wrapped around it.
The oxygen in serine (green in the bottom pic) is about 4 A from taxol which is replaced in TBB4_HUMAN by alanine. That means that OH is striped off.






Taxol Binding Site
Acidic residues (red); basic residues (blue)

Taxol binding site in a MT




start 300 K, than heat to 500 K for 50 ps and then cool down to 300 K for 240 ps

Taxol bound to a MT




start 300 K, then heat to 500 K for 50 ps then keep at 500 K for 50 ps and finally cool down to 300 K for 240 ps.

Force application to remove taxol in b 1 tubulin




Used to quantify the binding strength for different isotypes of tubulin

New compound to target beta VI: taxol with a methyl group at C-10



Further possibilites: SNP's



Alberta/Texas/New York collaboration: OUR GENERAL STRATEGY:

- 1. Model the isotype-drug binding sites
- 2. Design isotype-specific drugs
- 3. Simulate their action by computer models
- 4. Synthesize the drugs using well-known organic synthetic procedures and try to design drugs less likely to cross the blood-brain barrier.
- 5. Test the drugs for isotype specificity in vitro using fluorescence assays
- 6. Test the drugs on normal and transformed cells (e.g., breast and prostate). Effects on cell viability, mitotic arrest, microtubule distribution and tumor colonyforming ability will be measured.
- 7. Test the drugs in animal models. Mice with tumors will be used. Effectiveness of drugs against tumors will be measured. Side effects will be identified and toxicity determined.
- 8. Iterate the process.