

TOPOLOGICAL HYPERPLANE ARRANGEMENTS

David Forge, LRI, Université Paris-Sud

and

Thomas Zaslavsky, Binghamton University (SUNY)

Conference in Honour of Peter Orlik

Fields Institute, Toronto

19 August 2008

Topological hyperplane (topoplane):

$Y \subset X$ such that $(X, Y) \cong (\mathbb{R}^n, \mathbb{R}^{n-1})$.

\mathcal{A} : finite set of topoplanes.

Intersection semilattice:

$$\mathcal{L} := \{\bigcap \mathcal{S} : \mathcal{S} \subseteq \mathcal{A} \text{ and } \bigcap \mathcal{S} \neq \emptyset\},$$

partially ordered (as is customary) by reverse inclusion.

Flat: An intersection (an element of \mathcal{L}).

Main definition:

\mathcal{A} is an *arrangement of topoplanes* if:

$\forall H \in \mathcal{A}$ and $\forall Y \in \mathcal{L}$, either

$$Y \subseteq H$$

or

$$H \cap Y = \emptyset$$

or

$H \cap Y$ is a topoplane in Y .

Main Examples:

- Arrangement of real hyperplanes in \mathbb{R}^n (homogeneous or affine). (Winder, Zaslavsky, Las Vergnas)
- Arrangement of affine pseudohyperplanes representing an oriented matroid. (Las Vergnas)
- Intersection of a real hyperplane arrangement with a convex set. (Alexanderson & Wetzel, Zaslavsky)

Induced arrangement in a flat Y :

$$\mathcal{A}^Y := \{Y \cap H : H \in \mathcal{A} \text{ and } Y \not\subseteq H \text{ and } Y \cap H \neq \emptyset\}.$$

Region: Connected component of complement $X \setminus \bigcup \mathcal{A}$.

Face: Region of any \mathcal{A}^Y .

Theorem (Zaslavsky, 1977):

$$(1) \quad \# \text{ regions of } \mathcal{A} = \sum_{Y \in \mathcal{L}} |\mu(X, Y)|,$$

where μ is the Möbius function of \mathcal{L} ,
assuming the side condition that every region is a topological cell.

Primary Question:

Is this really new? Can we finagle it out of something ‘simpler’? Las Vergnas’s theorem for pseudohyperplane arrangements (oriented matroids)?

I.e.: $\exists \mathcal{A}'$ such that $\bigcup \mathcal{A}' = \bigcup \mathcal{A}$ and \mathcal{A}' is a pseudohyperplane arrangement?

Answer:


No !

Intersecting topoplanes may have the topology of two crossing hyperplanes,


$$(2) \quad (X, H_1, H_2, H_1 \cap H_2) \cong (\mathbb{R}^n, x_1 = 0, x_2 = 0, x_1 = x_2 = 0),$$

or of two noncrossing ‘flat’ topoplanes,

$$(3) \quad (X, H_1, H_2, H_1 \cap H_2) \cong (\mathbb{R}^n, G_+, G_-, x_1 = x_2 = 0).$$



(2)




(3)

Definition: \mathcal{A} is *transsective* if every intersecting pair of topoplanes crosses.

Fact: An arrangement of (affine) pseudohyperplanes is transsective.

Types of Topoplane Arrangement



Reglueing

This means there is another arrangement, \mathcal{A}' , that has the same faces as \mathcal{A} :

$$\bigcup \mathcal{A}' = \bigcup \mathcal{A}.$$

In the Plane:

Theorem 9. *For any arrangement of topolines, there is a transsective reglueing (i.e., every topline intersection is a crossing).*

(Proof by construction.)

Higher Dimensions:

Theorem 10. *For a simple topoplane arrangement in which every region is a cell, there is a transsective reglueing.*

(Proof by construction.)

Theorem 10'. *For a nonsimple such arrangement, there need not be a transsective reglueing.*

(Proof by example.)

PROOFS BY PICTURES

Elementary properties

- (1) If \mathcal{A} is an arrangement of topoplanes and Y is a flat, then the induced collection \mathcal{A}^Y is an arrangement of topoplanes.
- (2) For an arrangement of topoplanes, each interval in \mathcal{L} is a geometric lattice with rank given by codimension.
- (3) Suppose every region is a cell. Topoplanes H_1 and H_2 cross if and only if they intersect each other and each of the regions they form has boundary that meets both $H_1 \setminus H_2$ and $H_2 \setminus H_1$.
- (4) In a topoline arrangement every face is a cell.

Lemma 4. *Suppose every region is a cell. H_1 and H_2 cross iff they intersect and each region they form has boundary that meets both $H_1 \setminus H_2$ and $H_2 \setminus H_1$.*

Proof: Easy.


Lemma 6. *Suppose every region is a cell. If H_1 and H_2 , cross, then $Y \cap H_1$ and $Y \cap H_2$ cross in \mathcal{A}^Y for each $Y \in \mathcal{L}$ such that $Y \cap H_1, Y \cap H_2$ are distinct topoplanes in Y .*

Proof: Not as easy as you might think.

Reglueing in the Plane

Theorem 9. *For any arrangement of topolines, there is an arrangement that has the same faces, and in which every intersection is a crossing.*

Proof Sketch. We apply the method of descent to the number of noncrossing pairs of intersecting topolines. Suppose noncrossing topolines H^1, H^2 have intersection point Z .



$$\mathcal{A}' := \{K^1, K^2, K^3, K^4\}.$$

\mathcal{A}' has the same faces. Must check: \mathcal{A}' is an arrangement of topolines (takes some work), with fewer noncrossing pairs (nearly obvious).

Since there are fewer noncrossing topline pairs in the new arrangement, by continuing the process we get a transsective arrangement. \square

Reglueing Fails in Three Dimensions

Proof of Theorem 10' by a counterexample of five topoplanes in \mathbb{R}^3 :

$$H_1 = \{x : x_1 = 0\},$$

$$H_2 = \{x : x_2 = 0\},$$

$$H_3 = \{x : x_2 = |x_1|\},$$

$$H_4 = \{x : x_3 = 0\},$$

$$H_5 = \{x : x_2 + x_3 = 0\}.$$

Every pair crosses except H_2 and H_3 . The common point of all topoplanes is O , the origin.

The 1-dimensional flats are:

$$Z := H_1 \cap H_2 \cap H_3 = \{x : x_1 = x_2 = 0\},$$

$$H_1 \cap H_4 = \{x : x_1 = x_3 = 0\},$$

$$H_1 \cap H_5 = \{x : x_1 = 0, x_2 + x_3 = 0\},$$

$$Y := H_2 \cap H_4 \cap H_5 = \{x : x_2 = x_3 = 0\},$$

$$H_3 \cap H_4 = \{x : x_2 = |x_1|, x_3 = 0\},$$

$$H_3 \cap H_5 = \{x : x_2 = |x_1| = -x_3\}.$$

The only two 1-dimensional flats that lie in three topoplanes are Z and Y .

Fact: It is impossible to have a transsective arrangement whose regions are the same as those of this one.

Simple Arrangements Reglue

\mathcal{A} is *simple* if every flat is the intersection of the fewest possible topoplanes. Otherwise it is *multiple*.

Theorem 10. *For a simple topoplane arrangement in which every region is a cell, there is an arrangement that has the same faces, and in which every topoplane intersection is a crossing.*

Proof Sketch. Similar to the planar proof: the method of descent on the number of noncrossing intersecting pairs of topoplanes. The construction is the same. The complications are greater, but not too bad.

To show that \mathcal{A}' is an arrangement of topoplanes we consider the intersection of a topoplane H and a flat Y of the reglued arrangement \mathcal{A}' . This is the hard part of the proof. There are four cases, depending mostly on whether either H or Y is a topoplane or flat in the original arrangement \mathcal{A} . \square

Topoplanes vs. pseudohyperplanes

Projective pseudohyperplane arrangement:

A finite set \mathcal{P} of subspaces in \mathbb{RP}^n such that

- each $(\mathbb{RP}^n, H) \cong (\mathbb{RP}^n, \mathbb{RP}^{n-1})$,
- the intersection Y of any members of \mathcal{P} is a \mathbb{RP}^d , and
- for any other H , either $Y \subseteq H$ or $H \cap Y$ is a pseudohyperplane in Y .

Known: Every region is an open cell and its closure is a closed cell.

Affine pseudohyperplane arrangement:

$$\mathcal{P}_0 := \{H \setminus H_0\} \text{ in } \mathbb{R}^n = \mathbb{RP}^n \setminus H_0.$$

\mathcal{A} is *projectivizable* if it is homeomorphic to a \mathcal{P}_0 .

Two topoplanes are *parallel* if they are disjoint.

Lemma 11. *If a topoplane arrangement is projectivizable then it is transsective, parallelism is an equivalence relation on topoplanes, and every region is a cell.*

Proof: Easy.

Look at a transsective topoplane arrangement in which parallelism is an equivalence relation?

HOW TO AVOID BEING PROJECTIVE

1. Disconnection:

\mathcal{A} is *connected* if $\bigcup \mathcal{A}$ is connected.

Disconnected \mathcal{A} may be a pseudohyperplane arrangement. But:

Counterexample:

Put \mathcal{A}_1 and \mathcal{A}_2 in the right and left halfspaces of \mathbb{R}^n . Then $\mathcal{A}_1 \cup \mathcal{A}_2$ is disconnected.


Proposition 12. *If \mathcal{A}_1 has a pair of intersecting topoplanes, $\mathcal{A}_1 \cup \mathcal{A}_2$ is not projectivizable.*

Proof: Easy.

In the Plane:


Theorem 13. *A topoline arrangement is projectivizable iff it is transsective and parallelism is an equivalence relation.*

The diagram (next) shows the construction.



2. Connection:

Counterexample:



$$L := \{x : x_1 x_2 = 0 \text{ and } x_1, x_2 \geq 0\}.$$

Parallelism is not transitive. Connected, transsective, but not projectivizable.

Question:

In higher dimensions, is intransitivity of parallelism the only obstruction?

3. Restriction to a Domain

Restriction of \mathcal{A} to a domain:

$$\mathcal{A}^D := \{\text{components of } H \cap D : H \in \mathcal{A} \text{ and } H \cap D \neq \emptyset\},$$

where $D \subseteq \mathbb{R}^n$ is a cellular domain and \mathcal{A}^D is a topoline arrangement in D .
(Alexanderson and Wetzel, Zaslavsky)

Properties:

- \mathcal{A}^D is transsective if \mathcal{A} is transsective.
- Parallelism could be transitive in \mathcal{A} but not in \mathcal{A}^D .

Theorem (Las Vergnas, unpublished). *Any transsective topoline arrangement is the restriction to a cellular domain of a projectivizable arrangement.*

Question:

In higher dimensions, is failure of transsectivity the only obstruction to being the restriction of a projectivizable arrangement?

Las Vergnas has an apparent counterexample in dimension 3, being studied by J. Ramírez Alfonsín.

OPEN QUESTIONS

- (1) Is the condition that every region be a cell ever superfluous?
- (2) Are there simple properties that imply all intersecting topoplanes cross? For instance, if there are enough topoplanes?
- (3) Complexify!

REFERENCES

- [1] G.L. Alexanderson and John E. Wetzel, Dissections of a plane oval.
Amer. Math. Monthly **84** (1977), 442–449.
MR 58 #23976. Zbl. 375.50009.
- [2] Michel Las Vergnas, Matroïdes orientables.
C. R. Acad. Sci. Paris Sér. A–B **280** (1975), Ai, A61–A64.
MR 51 #7910. Zbl. 304.05013.
- [3] Thomas Zaslavsky, A combinatorial analysis of topological dissections.
Adv. Math. **25** (1977), 267–285.
MR 56 #5310. Zbl. 406.05004.
- [4] David Forge and Thomas Zaslavsky, On the division of space by topological hyperplanes.
European J. Combin., to appear.