

ASTALA'S CONJECTURE ON DISTORTION OF HAUSDORFF MEASURES UNDER QUASICONFORMAL MAPS IN THE PLANE AND RELATED REMOVABILITY PROBLEMS

Ignacio Uriarte-Tuero

(joint works with Michael T. Lacey, Eric T. Sawyer, and Xavier
Tolsa)

May 31, 2010

SIZE OF SETS: HAUSDORFF MEASURES AND DIMENSION

DEFINITION

A “**gauge function**” is a continuous increasing $h : [0, \infty) \rightarrow [0, \infty)$ such that $h(0) = 0$.

Let $E \subset \mathbb{C}$ be compact and let $0 < \delta \leq \infty$. Define

$$\mathcal{H}_\delta^h(E) = \inf \left\{ \sum_{i=1}^{\infty} h(r_i) : E \subset \bigcup_{i=1}^{\infty} B(x_i, r_i), 2r_i \leq \delta \right\}$$

If $\delta = \infty$: \mathcal{H}_∞^h is called **Hausdorff content**.

$$\mathcal{H}^h(E) = \lim_{\delta \rightarrow 0} \mathcal{H}_\delta^h(E) \quad \text{“Hausdorff Measure”}$$

If $h(t) = t^s$, then $\mathcal{H}^h(E) = \mathcal{H}^s(E)$

$$\text{Dim}(E) = \inf\{s > 0 : \mathcal{H}^s(E) = 0\} = \sup\{t > 0 : \mathcal{H}^t(E) = \infty\}$$

$\mathcal{H}^s(E) = 0$ if and only if $\mathcal{H}_\infty^s(E) = 0$

An orientation-preserving homeomorphism $\phi : \Omega \rightarrow \Omega'$ between planar domains $\Omega, \Omega' \subset \mathbb{C}$ is called **K -quasiconformal (K -QC)** if it belongs to the Sobolev space $W_{loc}^{1,2}(\Omega)$ and satisfies the *distortion inequality*

$$\max_{\alpha} |\partial_{\alpha} \phi| \leq K \min_{\alpha} |\partial_{\alpha} \phi| \quad \text{a.e. in } \Omega. \quad (1)$$

Actually, $J(z, \phi) \leq |D\phi(z)|^2 \leq K J(z, \phi)$

If ϕ is just required to be continuous instead of homeomorphism: ϕ is **K -quasiregular (K -QR)**


In \mathbb{C} , any ϕ K -QR solves the Beltrami equation $\bar{\partial}\phi = \mu\partial\phi$, where $\|\mu\|_{\infty} \leq \frac{K-1}{K+1} < 1$.

Using Calderón-Zygmund theory: there is a unique homeomorphic normalized solution f to Beltrami (e.g. fixing 0, 1, ∞ .)

Infinitesimally, quasiconformal mappings carry circles to ellipses with eccentricity at most K .

Macroscopically: they take discs to “quasidisks” (one can inscribe and circumscribe circles of radii r and R such that $\frac{R}{r} \leq C(K)$).
I.e. something somewhat round, neither “cigarettes” nor “horseshoes”.

THESE MAPPINGS CAN DISTORT DIMENSION



THEOREM (STOILOW FACTORIZATION)

Let f be the homeomorphic normalized solution to the Beltrami equation:

$$\bar{\partial}\phi = \mu\partial\phi, \text{ where } \|\mu\|_{\infty} < 1. \quad (2)$$

Let $g \in W_{loc}^{1,2}(\Omega)$ be another solution to (2) Then there exists a holomorphic $\psi : f(\Omega) \rightarrow \mathbb{C}$ such that $g(z) = \psi(f(z))$. Conversely, for any holomorphic ψ in $f(\Omega)$, $\psi \circ f \in W_{loc}^{1,2}$ and solves Beltrami in $f^{-1}(\Omega)$.

Examples of K -quasiconformal f :

- f conformal
- f bilipschitz ($c|z - w| \leq |f(z) - f(w)| \leq C|z - w|$)
- radial stretching: $z \rightarrow |z|^{K-1}z$; $z \rightarrow |z|^{\frac{1}{K}-1}z$

THEOREM (ASTALA - AREA DISTORTION - ACTA '94)

If ϕ is K -quasiconformal, then

$$|\phi(E)| \lesssim |E|^{1/K}.$$

THEOREM (ASTALA - HIGHER INTEGRABILITY - ACTA '94)

If ϕ is K -quasiconformal, then

$$\|\phi\|_{W_{\text{loc}}^{1,p}} < \infty, \quad p < \frac{2K}{K-1} (> 2!).$$

THEOREM (ASTALA - H-DIMENSION DISTORTION - ACTA '94)

For any compact set E with Hausdorff dimension $0 < t < 2$ and any K -quasiconformal mapping ϕ we have

$$\frac{1}{K} \left(\frac{1}{t} - \frac{1}{2} \right) \leq \frac{1}{\dim(\phi E)} - \frac{1}{2} \leq K \left(\frac{1}{t} - \frac{1}{2} \right) \quad (3)$$

Finally, these bounds are optimal, in that equality may occur in either estimate.

In other words, $\dim(E) \leq t$ implies $\dim(\phi E) \leq t' = \frac{2Kt}{2+(K-1)t}$.

Equation (3) conjectured by Iwaniec, Martin - Acta '93 (n instead of 2.)

Progress on these questions in higher dimensions is much harder.

MAIN THEOREM (LACEY, SAWYER, UT '08 - ACTA, TO APPEAR)

If ϕ is a planar K -quasiconformal mapping, $0 \leq t \leq 2$ and $t' = \frac{2Kt}{2+(K-1)t}$, then we have the implication below for all compact sets $E \subset \mathbb{C}$.

$$\mathcal{H}^t(E) = 0 \implies \mathcal{H}^{t'}(\phi E) = 0, \quad (4)$$

IF ϕ IS K -QUASICONFORMAL, ϕ^{-1} IS ALSO
 K -QUASICONFORMAL


So, it follows that

$$\mathcal{H}^{t'}(\phi E) > 0 \implies \mathcal{H}^t(E) > 0.$$

$\mathcal{H}^t(E) < \infty$ NEED NOT IMPLY $\mathcal{H}^{t'}(\phi E) = 0$. (UT - IMRN '08)

THE CONCLUSION $\mathcal{H}^{t'}(E) = 0$ CANNOT BE STRENGTHENED TO
H-D W.R.T. A GAUGE. (UT - IMRN '08)

For any gauge function h satisfying $\lim_{s \rightarrow 0} \frac{s^{t'}}{h(s)} = 0$, there exists a
compact set E and K -quasiconformal mapping ϕ with $\mathcal{H}^t(E) = 0$
but $\mathcal{H}^h(\phi E) = \infty$.



RELEVANCE OF THE CONJECTURE: ANALYTIC CAPACITY

- $E \subset \mathbb{C}$ compact
- $\gamma(E) = 0$ if and only if for any $U \supset E$ open, any bounded analytic $f : U \setminus E \rightarrow \mathbb{C}$, extends analytically to U .
- Independent of U
- $\mathcal{H}^1(E) = 0$ implies $\gamma(E) = 0$ (Painlevé): hence $\dim_H(E) < 1$ implies $\gamma(E) = 0$.
- $\dim_H(E) > 1$ implies $\gamma(E) > 0$ (Ahlfors - Duke '47)
- If $0 < \mathcal{H}^1(E) < \infty$ (or sigma-finite): $E = G \cup R \cup N$ (disjoint union), with $\mathcal{H}^1(G) = 0$, $\mathcal{H}^1(R \setminus \bigcup_{i=1}^{\infty} \Gamma_i) = 0$ (Γ_i Lipschitz or C^1 curves), for any rectifiable Γ , $\mathcal{H}^1(N \cap \Gamma) = 0$ (e.g. Garnett set)
Then, $\gamma(E) = 0$ if and only if $R = \emptyset$ (David - Rev. Mat. Ib. '98; Tolsa - Acta '03)
- General case: crucial Tb theorem (Nazarov-Treil-Volberg - Acta '03); complete solution (Tolsa - Acta '03).

RELEVANCE OF THE CONJECTURE:

L^∞ - K -REMOVABILITY

- $E \subset \mathbb{C}$ compact is L^∞ - K -removable if any $f : \mathbb{C} \setminus E \rightarrow \mathbb{C}$ in L^∞ , K -quasiregular, can be extended to a global ($\mathbb{C} \rightarrow \mathbb{C}$) K -quasiregular map.
By Stoilow and Liouville's thms, such extension is constant.
- By Stoilow's theorem, the critical dimension is $\frac{2}{K+1}$.

THEOREM (ASTALA, CLOP, MATEU, OROBITG, UT - DUKE '08)

Let $E \subset \mathbb{C}$ compact and f K -quasiregular

- (A) $\mathcal{H}^{\frac{2}{K+1}}(E) = 0$ (sigma-finite) implies $\mathcal{H}^1(fE) = 0$ (resp. sigma-finite)
- (B) $\mathcal{H}^{\frac{2}{K+1}}(E)$ sigma-finite and $K > 1$ implies E is L^∞ - K -removable.
- (C) There exists E with $\dim_H(E) = \frac{2}{K+1}$ not L^∞ - K -removable.

RELEVANCE OF THE CONJECTURE:

L^∞ - K -REMOVABILITY II

- Proof of easier version of case (B): if $\mathcal{H}^{\frac{2}{K+1}}(E) = 0$, E is L^∞ - K -removable. (Conjectured Iwaniec-Martin, Acta'93; proved Astala '04 approx.)
- Let E compact, $\mathcal{H}^{\frac{2}{K+1}}(E) = 0$, and $g : \mathbb{C} \setminus E \rightarrow \mathbb{C}$ K -quasiregular, bounded.
- By Stoilow: $g = h \circ f$ with h analytic and bounded, and f K -quasiconformal.
- μ defined a.e. : solve Beltrami : f defined everywhere. The problem is whether h can be extended, i.e. whether $\gamma(fE) = 0$ for any K -quasiconformal f .
- By part (A), $\mathcal{H}^1(fE) = 0$ (quasiconformal distortion of sets: Astala's conjecture)
- By Painlevé, $\gamma(fE) = 0$, so h can be extended analytically (removability results for bounded analytic functions, i.e. analytic capacity).

RELEVANCE OF THE CONJECTURE: L^∞ - K -REMOVABILITY III

MAIN THEOREM (TOLSA, UT '09)

Let $E \subset \mathbb{C}$ compact and $\varphi : \mathbb{C} \rightarrow \mathbb{C}$ K -quasiconformal, $K > 1$. If E is contained in a ball B , then

$$\frac{\dot{C}_{\frac{2K}{2K+1}, \frac{2K+1}{K+1}}(E)}{\text{diam}(B)^{\frac{2}{K+1}}} \geq c^{-1} \left(\frac{\gamma(\varphi(E))}{\text{diam}(\varphi(B))} \right)^{\frac{2K}{K+1}}.$$

Remarks

- $\dot{C}_{\alpha,p}$ is a Riesz capacity associated to non linear potential.
- Indices $\alpha = \frac{2K}{2K+1}$, $p = \frac{2K+1}{K+1}$ are sharp.
- E non K -removable $\Rightarrow \dot{C}_{\frac{2K}{2K+1}, \frac{2K+1}{K+1}}(E) > 0 \Rightarrow \mathcal{H}^{\frac{2}{K+1}}(E)$ non σ -finite. (Recover Astala, Clop, Mateu, Orobitg, UT '08 (B))
- Theorem only holds for $K > 1$.

$$\begin{aligned}\dot{C}_{\alpha,p}(E) &= \inf\{\|\psi\|_{\dot{W}_{\alpha,p}}^p : \psi \in C_c^\infty(\mathbb{C}), \psi \geq 1 \text{ on } E\} \\ &= \sup_{\mu \in \mathcal{M}^+(E)} \left(\frac{\mu(E)}{\|I_\alpha * \mu\|_{p'}} \right)^p ; \quad I_\alpha(x) = \frac{c_\alpha}{|x|^{n-\alpha}} \text{ in } \mathbb{R}^n.\end{aligned}$$

$\mathcal{M}^+(E)$ are the positive Radon measures on E .

Wolff:

$$\dot{C}_{\alpha,p}(E) \approx \sup\{\mu(E) : \text{supp}(\mu) \subset E; \dot{W}_{\alpha,p}^\mu(x) \leq 1 \forall x \in \mathbb{C}\}, \quad (5)$$

where

$$\dot{W}_{\alpha,p}^\mu(x) = \int_0^\infty \left(\frac{\mu(B(x,r))}{r^{2-\alpha p}} \right)^{p'-1} \frac{dr}{r}.$$

For $\dot{C}_{\frac{2K}{2K+1}, \frac{2K+1}{K+1}}(E)$,

$$\dot{W}_{\frac{2K}{2K+1}, \frac{2K+1}{K+1}}^\mu(x) = \int_0^\infty \left(\frac{\mu(B(x,r))}{r^{\frac{2}{K+1}}} \right)^{\frac{K+1}{K}} \frac{dr}{r}.$$


L^∞ - K -REMOVABILITY IV: SHARPNESS

There are sets E with $\mathcal{H}^{\frac{2}{K+1}}(E)$ non σ -finite such that $\dot{C}_{\alpha,p}(E) = 0$ for all α, p with $2 - \alpha p = \frac{2}{K+1}$. \longrightarrow New examples of K -removable sets.

THEOREM (TOLSA, UT '09)

For all $\beta, q > 0$ such that $2 - \beta q = \frac{2}{K+1}$ and $q' < \frac{K+1}{K}$, there exists $E \subset \mathbb{C}$ and a K -quasiconformal map φ such that $\gamma(\varphi E) > 0$ (and so $\dot{C}_{\frac{2K}{2K+1}, \frac{2K+1}{K+1}}(E) > 0$), but $\dot{C}_{\beta,q}(E) = 0$.

$$\Rightarrow \quad \dot{C}_{\alpha,p}(E) = 0 \quad \text{if } 2 - \alpha p > \frac{2}{K+1}.$$



THEOREM (TOLSA, UT '09)

Let h be a positive function on $(0, \infty)$ such that

$$\varepsilon(r) = \frac{h(r)}{r^{\frac{2}{K+1}}} \rightarrow 0 \quad \text{as } r \rightarrow 0.$$

Then there is a compact set $E \subset \mathbb{C}$ such that $\mathcal{H}^h(E) = 0$ and a K -quasiconformal map φ such that $\gamma(\varphi(E)) > 0$ (and thus $\dot{C}_{\frac{2K}{2K+1}, \frac{2K+1}{K+1}}(E) > 0$.)

- E can be taken so that $\mathcal{H}^h(E) = 0$ for all h which is nondecreasing and

$$\int_0^\infty \left(\frac{h(r)}{r^{\frac{2}{K+1}}} \right)^a \frac{dr}{r} < \infty$$

for some $a > 0$ (E independent of h).

MORE ON K -REMOVABILITY

- Instead of L^∞ , work with BMO.
- Analytic case: $\gamma_{BMO}(E) = 0$ if and only if $\mathcal{H}^1(E) = 0$ (Král - Wilhelm Pieck Univ. '78; Kaufman - Pacific J. '82)
- K -quasiregular case: if $\mathcal{H}^{\frac{2}{K+1}}(E) = 0$, then E is BMO- K -removable (Astala, Clop, Mateu, Orobitg, UT '08).
There exists E with $0 < \mathcal{H}^{\frac{2}{K+1}}(E) < \infty$, not BMO- K -removable (UT '08). Hence the L^∞ and BMO K -removability problems are different (answering Question 4.2 in Astala, Clop, Mateu, Orobitg, UT '08)
- Can also work with Lipschitz (or Hölder) (α).
- Analytic case: $\gamma_{Lip(\alpha)}(E) = 0$ if and only if $\mathcal{H}^{1+\alpha}(E) = 0$ (Dolženko - Uspekhi Mat. Nauk. '63)
- K -quasiregular case: let $d_\alpha = \frac{2}{K+1}(1 + \alpha K)$. If $\mathcal{H}^{d_\alpha}(E) = 0$, E is $Lip(\alpha)$ - K -removable (Clop - Ann. Acad. Sci. Fenn. '07).
There exists E with $0 < \mathcal{H}^{d_\alpha}(E) < \infty$, not $Lip(\alpha)$ - K -removable (Clop, UT - J. d'Analyse, '09).

IDEAS OF PROOF OF ASTALA'S CONJECTURE

Recall statement of Astala's conjecture: (4)

- Can reduce to the case of K -quasiconformal ϕ , with $0 < K < 1 + \epsilon$ very close to one.
- ϕ is a solution to the Beltrami equation $\bar{\partial}\phi = \mu\partial\phi$, where $\|\mu\|_\infty \leq K - 1 \leq \epsilon$.
- We have the Beurling operator which intertwines ∂ and $\bar{\partial}$:

$$B f(z) = \int \frac{f(w)}{(z - w)^2} dz \wedge d\bar{z}$$
$$\partial f = 1 + B(\bar{\partial} f) \quad (\bar{\partial} f = \mu \partial f; \text{homeo, normalized})$$

- And so, we have a formal expansion

$$\bar{\partial} f = \mu \partial f = \mu + \mu B \mu + \mu B(\mu B \mu) + \dots$$

- There is one additional factorization that we will come to.

APPROXIMATING E : PACKING CONDITION


DEFINITION

Let \mathcal{P} be a finite collection of disjoint dyadic cubes in the plane. Let $0 < t < 2$. We denote the t -Carleson packing norm of \mathcal{P} as follows:

$$\|\mathcal{P}\|_{t\text{-pack}} = \sup_Q \left[\ell(Q)^{-t} \sum_{\substack{P \in \mathcal{P} \\ P \subset Q}} \ell(P)^t \right]^{1/t},$$

where the supremum is taken over all dyadic cubes Q .



Q



Q



not allowed!



$$\sum_{P \in \mathcal{P}} \ell(P) \leq \ell(Q)$$

APPROXIMATING E : PACKING LEMMA

LEMMA (PACKING LEMMA)

Suppose E is a compact subset of $(0, 1)^2 \subset \mathbb{C}$, $0 < t < 2$, and $\varepsilon > 0$. Then there is a positive constant C and a finite collection of closed dyadic cubes $\mathcal{P} = \{P_i\}_{i=1}^N$ such that

- (A) $4P_i \cap 4P_j = \emptyset$ for $i \neq j$.
- (B) $E \subset \bigcup_{i=1}^N 12P_i$.
- (C) $\|\mathcal{P}\|_{t\text{-pack}} \leq 1$.
- (D) $\sum_{i=1}^N \ell(P_i)^t \leq C (\mathcal{H}_\infty^t(E) + \varepsilon)$.

$$d\alpha = \sum_{P \in \mathcal{P}} \frac{\mathbf{1}_P}{\ell(P)^{2-t}} dx. \quad \text{Remember this!}$$

A WEIGHTED ESTIMATE FOR BEURLING TRANSFORM

THEOREM

For any $0 < t < 2$, and any collection \mathcal{P} with $\|\mathcal{P}\|_{t\text{-pack}} < \infty$, we have

$$\|Bf\|_{L^p(\alpha)} \lesssim \|f\|_{L^p(\alpha)}, \quad 1 < p < \infty.$$

HEURISTICS OF THE PROOF:

- 1 The measures α are **not** A_p weights, so the inequality above is certainly not trivial.
- 2 Indeed, one may need to use the general theory of two-weight inequalities, which is a rather thin literature.
- 3 A direct proof is the hardest route. Slightly easier is the weak-type approach. Easier still is the restricted weak-type approach.
- 4 The combinatorial properties of t -packing permit a simple self-contained proof.

COMBINING THE ELEMENTS OF THE PROOF

- 1 Given $\epsilon > 0$, approximate E by the Packing Lemma, approximate by the cubes \mathcal{P} .
- 2 Factorize $\varphi = \varphi_{\text{in}} \circ \varphi_{\text{out}}$, where $\varphi_{\text{in/out}}$ is conformal in/out—side $\varphi_{\text{out}}(\bigcup P_j) / \bigcup P_j$.
- 3 The inside piece has **improved** endpoint integrability for Sobolev spaces, so it is easier to handle (Astala - Nesi.)
- 4 So it remains to control the outside piece, and here the weighted inequality is decisive.

Recall $d\alpha = dx \sum_{P \in \mathcal{P}} \mathbf{1}_P / \ell(P)^{2-t}$, and $\sum_{P \in \mathcal{P}} \ell(P)^t < \epsilon$.

$$\begin{aligned}
 \left(\sum_{P \in \mathcal{P}} \text{diam}(f(P))^t \right)^{\frac{2}{t}} &\lesssim \int_E J(z, f) \alpha(dz \wedge d\bar{z}) \\
 &= \int_E (|f_z|^2 - |f_{\bar{z}}|^2) \alpha(dz \wedge d\bar{z}) \quad (f_z = 1 + B(f_{\bar{z}})) \\
 &\leq 2 \int_E (\mathbf{1} + |\mathbf{B}(f_{\bar{z}})|^2 + |f_{\bar{z}}|^2) \alpha(dz \wedge d\bar{z}) \\
 \int_E \mathbf{1} \alpha(dz \wedge d\bar{z}) &= \sum_{P \in \mathcal{P}} \ell(P)^t < \epsilon \quad \text{by construction.} \\
 \int_E |\mathbf{B}(f_{\bar{z}})|^2 \alpha(dz \wedge d\bar{z}) &\lesssim \int_E |f_{\bar{z}}|^2 \alpha(dz \wedge d\bar{z}) \quad \text{by wtd ineq}
 \end{aligned}$$

Recall: $\bar{\partial}f = \mu\partial f = \mu + \mu B \mu + \mu B(\mu B \mu) + \dots$, so

$$\begin{aligned} \left[\int_E |f_{\bar{z}}|^2 \alpha(dz \wedge d\bar{z}) \right]^{1/2} &\leq \sum_{n=0}^{\infty} \|\mu B \mu \cdots B \mu\|_{L^2(\alpha)} \\ &\leq \sum_{n=0}^{\infty} [\|\mu\|_{\infty} \|B\|_{L^2(\alpha) \rightarrow L^2(\alpha)}]^n \alpha(\mathbb{C})^{1/2} \\ &\lesssim \epsilon^{1/2} \end{aligned}$$

IDEAS OF PROOF OF REMOVABILITY RESULT: ANALYTIC CAPACITY AND NON LINEAR POTENTIALS

THEOREM (TOLSA - INDIANA '02)

$$\gamma(E) \approx \sup \{ \mu(E) : \text{supp}(\mu) \subset E; M\mu(x) + c_\mu(x) \leq 1 \ \forall x \in \mathbb{C} \},$$

$$\text{where } M\mu(x) = \sup_{r>0} \frac{\mu(B(x,r))}{r}, \quad c_\mu(x)^2 = \iint \frac{1}{R(x,y,z)^2} d\mu(y) d\mu(z).$$

Recall:

$$\dot{W}_{2/3,3/2}^\mu(x) = \int_0^\infty \left(\frac{\mu(B(x,r))}{r} \right)^2 \frac{dr}{r} \approx \sum_{k \in \mathbb{Z}} \left(\frac{\mu(B(x,2^k))}{2^k} \right)^2.$$

We have (recall Wolff: (5))

$$M\mu(x)^2 + c_\mu(x)^2 \lesssim \dot{W}_{2/3,3/2}^\mu(x) \quad \Rightarrow \quad \gamma(E) \gtrsim \dot{C}_{2/3,3/2}(E).$$

γ AND $\dot{C}_{2/3,3/2}$


$$\gamma \gtrsim \dot{C}_{2/3,3/2}.$$

But it is **false** that $\gamma \approx \dot{C}_{2/3,3/2}$.

Example: if L is a segment, then

$$\gamma(L) = \frac{\mathcal{H}^1(L)}{4}, \quad \dot{C}_{2/3,3/2}(L) = 0.$$

But for “typical Cantor sets”, $\gamma(F) \approx \dot{C}_{2/3,3/2}(F)$.



FIRST STEP FOR PROOF OF MAIN RESULT

Distortion of $\dot{C}_{2/3,3/2}$:

THEOREM (TOLSA, UT '09)

Let $E \subset \mathbb{C}$ compact and $\varphi : \mathbb{C} \rightarrow \mathbb{C}$ K -quasiconformal, $K > 1$. If E is contained in a ball B , then

$$\frac{\dot{C}_{\frac{2K}{2K+1}, \frac{2K+1}{K+1}}(E)}{\text{diam}(B)^{\frac{2}{K+1}}} \geq c^{-1} \left(\frac{\dot{C}_{2/3,3/2}(\varphi(E))}{\text{diam}(\varphi(B))} \right)^{\frac{2K}{K+1}}.$$

- The theorem also holds for other capacities $\dot{C}_{\alpha,p}$.

SECOND STEP: CORONA TYPE DECOMPOSITION

\mathcal{D} = dyadic lattice.

$\mathcal{T} \subset \mathcal{D}$ is a **tree** if:

- There exists $Q_0 \in \mathcal{T}$ which contains all other $Q \in \mathcal{T}$.
- If $Q, R \in \mathcal{T}$, with $Q \subset R$, then \mathcal{T} contains also the intermediate squares

$$P \text{ such that } Q \subset P \subset R.$$

Q_0 = Top square of \mathcal{T} .

Leaves of \mathcal{T} = stopping squares.

A **corona type decomposition** is a partition of \mathcal{D} (or $\mathcal{D}(R_0)$) into trees.

THE CORONA TYPE DECOMPOSITION

We set $\theta(Q) := \frac{\mu(Q)}{\ell(Q)}$,

$$c^2(\mu) := \iiint \frac{1}{R(x, y, z)^2} d\mu(x) d\mu(y) d\mu(z).$$

LEMMA (CORONA DECOMPOSITION [TOLSA - ANNALS OF MATH. '05])

Let μ with linear growth and $\text{supp}(\mu) \subset Q_0$.

There exists a corona type decomposition of $\mathcal{D}(Q_0)$ such that

$$\sum_{Q \in \text{Top}} \theta(Q)^2 \mu(Q) \leq C (\mu(\mathbb{C}) + c^2(\mu)),$$

where, for each tree \mathcal{T} with top square $Q \in \text{Top}$ there exists a chord arc curve $\Gamma_{\mathcal{T}}$ such that

- If $P \in \mathcal{T}$, then $8P \cap \Gamma_{\mathcal{T}} \neq \emptyset$.*
- If $P \in \mathcal{T}$, then $\theta(P) \approx \theta(Q)$.*

SOME IDEAS FOR THE PROOF OF THE MAIN THEOREM

Take μ supported on E such that $\mu(E) \approx \gamma(E)$, with linear growth and $c^2(\mu) \leq \mu(E)$.

By Tchebychev, there exists $F \subset E$, $\mu(F) \geq \mu(E)/2$, such that

$$\sum_{Q \in \text{Top}: x \in Q} \theta(Q)^2 \lesssim 1 \quad \text{for } x \in F.$$

If $\text{Top} = \mathcal{D}$, then

$$W_{2/3, 3/2}^\mu(x) \approx \sum_{Q \in \text{Top}: x \in Q} \theta(Q)^2 \lesssim 1 \quad \text{for } x \in F..$$

In the general case we combine ideas from distortion of $\dot{C}_{2/3, 3/2}$ with improved distortion of estimates of subset of chord arc curves.

Recall [ACM+]: If $E \subset \Gamma$, Γ chord arc curve, then

$$\dim_H(\varphi(E)) > \frac{2}{K+1}.$$

REFERENCES:

- ACM+ Astala, Clop, Mateu, Orobitg, U-T, “Distortion of Hausdorff measures and improved Painlevé removability for quasiregular mappings”, *Duke Math. J.* 141, no. 3 (2008), 539–571
- CUT Clop, U-T, “Sharp nonremovability examples for Hölder continuous quasiregular maps in the plane”, *J. d’Anal.*, (2009)
- LSUTA Lacey, Sawyer, U-T, “Astala’s conjecture on distortion of Hausdorff measures under quasiconformal maps in the plane”, (2008), *Acta Math.*, ??
- LSUTB Lacey, Sawyer, U-T, “A characterization of two weight norm inequalities for maximal singular integrals”, (2008), [Preprint]
- TUT Tolsa, U-T, “Quasiconformal maps, analytic capacity, and non linear potentials”, (2009), [Preprint]
- UT U-T, “Sharp Examples for Planar Quasiconformal Distortion of Hausdorff Measures and Removability”, *Int. Math. Res. Not.* (2008), no. 14, Art. ID rnn047, 43 pp.