

Some Analytic Aspects of Vafa-Witten Twisted $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory

(thesis work under Tom Mrowka)

Ben Mares

May 14, 2011

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Goal

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- Vafa-Witten invariants are only conjecturally defined.
- For many Kähler manifolds they have been “computed” by algebraic methods, and the answers satisfy the conjectured properties.
- We want to prove that these invariants exist for any oriented Riemannian manifold.
- We construct a partial Uhlenbeck compactification.
- Many issues remain unresolved before invariants become rigorous.

Goal

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- Vafa-Witten invariants are only conjecturally defined.
- For many Kähler manifolds they have been “computed” by algebraic methods, and the answers satisfy the conjectured properties.
- We want to prove that these invariants exist for any oriented Riemannian manifold.
- We construct a partial Uhlenbeck compactification.
- Many issues remain unresolved before invariants become rigorous.

Goal

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- Vafa-Witten invariants are only conjecturally defined.
- For many Kähler manifolds they have been “computed” by algebraic methods, and the answers satisfy the conjectured properties.
- We want to prove that these invariants exist for any oriented Riemannian manifold.
- We construct a partial Uhlenbeck compactification.
- Many issues remain unresolved before invariants become rigorous.

Goal

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- Vafa-Witten invariants are only conjecturally defined.
- For many Kähler manifolds they have been “computed” by algebraic methods, and the answers satisfy the conjectured properties.
- We want to prove that these invariants exist for any oriented Riemannian manifold.
- We construct a partial Uhlenbeck compactification.
- Many issues remain unresolved before invariants become rigorous.

Goal

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- Vafa-Witten invariants are only conjecturally defined.
- For many Kähler manifolds they have been “computed” by algebraic methods, and the answers satisfy the conjectured properties.
- We want to prove that these invariants exist for any oriented Riemannian manifold.
- We construct a partial Uhlenbeck compactification.
- Many issues remain unresolved before invariants become rigorous.

Geometry in four dimensions

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Possibly the most prominent feature of four-dimensional geometry:

$$1 \longrightarrow \{\pm I\} \longrightarrow \mathrm{SO}(4) \longrightarrow \mathrm{SO}(3) \times \mathrm{SO}(3) \longrightarrow 1$$

If $\dim V = 4$, then

$$\Lambda^2 V = \Lambda^+ V \oplus \Lambda^- V$$

$$6 = 3 + 3$$

$$1 \longrightarrow \{\pm I\} \longrightarrow \mathrm{SO}(V) \longrightarrow \mathrm{SO}(\Lambda^+ V) \times \mathrm{SO}(\Lambda^- V) \longrightarrow 1$$

$\Lambda^+ V$ is an **oriented** three-dimensional vector space associated to any 4-D oriented Euclidean V .

Geometry in four dimensions

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Possibly the most prominent feature of four-dimensional geometry:

$$1 \longrightarrow \{\pm I\} \longrightarrow \mathrm{SO}(4) \longrightarrow \mathrm{SO}(3) \times \mathrm{SO}(3) \longrightarrow 1$$

If $\dim V = 4$, then

$$\begin{aligned}\Lambda^2 V &= \Lambda^+ V \oplus \Lambda^- V \\ 6 &= 3 + 3\end{aligned}$$

$$1 \longrightarrow \{\pm I\} \longrightarrow \mathrm{SO}(V) \longrightarrow \mathrm{SO}(\Lambda^+ V) \times \mathrm{SO}(\Lambda^- V) \longrightarrow 1$$

$\Lambda^+ V$ is an **oriented** three-dimensional vector space associated to any 4-D oriented Euclidean V .

Geometry in four dimensions

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Possibly the most prominent feature of four-dimensional geometry:

$$1 \longrightarrow \{\pm I\} \longrightarrow \mathrm{SO}(4) \longrightarrow \mathrm{SO}(3) \times \mathrm{SO}(3) \longrightarrow 1$$

If $\dim V = 4$, then

$$\begin{aligned}\Lambda^2 V &= \Lambda^+ V \oplus \Lambda^- V \\ 6 &= 3 + 3\end{aligned}$$

$$1 \longrightarrow \{\pm I\} \longrightarrow \mathrm{SO}(V) \longrightarrow \mathrm{SO}(\Lambda^+ V) \times \mathrm{SO}(\Lambda^- V) \longrightarrow 1$$

$\Lambda^+ V$ is an **oriented** three-dimensional vector space associated to any 4-D oriented Euclidean V .

The cross product

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Let X be an oriented Riemannian four-manifold. For any $x \in X$, take $V = T_x^*X$.

- Choose an oriented orthonormal basis $\{e^0, e^1, e^2, e^3\}$.
- An oriented orthonormal basis for $\Lambda^+ T_x^*X$ is

$$\sigma^1 = e^0 \wedge e^1 + e^2 \wedge e^3,$$

$$\sigma^2 = e^0 \wedge e^2 + e^3 \wedge e^1,$$

$$\sigma^3 = e^0 \wedge e^3 + e^1 \wedge e^2.$$

- Define the cross product on $\Lambda^+ T_x^*X$ via $\{\sigma^i\}$ components, so $\sigma^1 \times \sigma^2 = \sigma^3$.

The cross product

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Let X be an oriented Riemannian four-manifold. For any $x \in X$, take $V = T_x^*X$.

- Choose an oriented orthonormal basis $\{e^0, e^1, e^2, e^3\}$.
- An oriented orthonormal basis for $\Lambda^+ T_x^*X$ is

$$\sigma^1 = e^0 \wedge e^1 + e^2 \wedge e^3,$$

$$\sigma^2 = e^0 \wedge e^2 + e^3 \wedge e^1,$$

$$\sigma^3 = e^0 \wedge e^3 + e^1 \wedge e^2.$$

- Define the cross product on $\Lambda^+ T_x^*X$ via $\{\sigma^i\}$ components, so $\sigma^1 \times \sigma^2 = \sigma^3$.

The cross product

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Let X be an oriented Riemannian four-manifold. For any $x \in X$, take $V = T_x^*X$.

- Choose an oriented orthonormal basis $\{e^0, e^1, e^2, e^3\}$.
- An oriented orthonormal basis for $\Lambda^+ T_x^*X$ is

$$\sigma^1 = e^0 \wedge e^1 + e^2 \wedge e^3,$$

$$\sigma^2 = e^0 \wedge e^2 + e^3 \wedge e^1,$$

$$\sigma^3 = e^0 \wedge e^3 + e^1 \wedge e^2.$$

- Define the cross product on $\Lambda^+ T_x^*X$ via $\{\sigma^i\}$ components, so $\sigma^1 \times \sigma^2 = \sigma^3$.

The de Rham complex

Latest version at <http://tensorial.com/math/>

Inside the de Rham complex

$$0 \rightarrow \Omega^0 \xrightarrow{d} \Omega^1 \xrightarrow{d} \Omega^2 \xrightarrow{d} \Omega^3 \xrightarrow{d} \Omega^4 \rightarrow 0$$

is the subcomplex

$$\begin{array}{ccccccc} 0 & \rightarrow & \Omega^0 & \xrightarrow{d} & \Omega^1 & \xrightarrow{d^+} & \Omega^{2,+} \rightarrow 0. \\ & & b^0 & & b^1 & & b^+ \end{array}$$

Given a principal bundle P with connection A ,

$$0 \rightarrow \Omega^0(\mathfrak{g}_P) \xrightarrow{d_A} \Omega^1(\mathfrak{g}_P) \xrightarrow{d_A^+} \Omega^{2,+}(\mathfrak{g}_P) \rightarrow 0.$$

The double-composition is

$$d_A^+ \circ d_A = [F_A^+, \bullet].$$

This defines a complex when $F_A^+ = 0$.

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The de Rham complex

Latest version at <http://tensorial.com/math/>

Inside the de Rham complex

$$0 \rightarrow \Omega^0 \xrightarrow{d} \Omega^1 \xrightarrow{d} \Omega^2 \xrightarrow{d} \Omega^3 \xrightarrow{d} \Omega^4 \rightarrow 0$$

is the subcomplex

$$\begin{array}{ccccccc} 0 & \rightarrow & \Omega^0 & \xrightarrow{d} & \Omega^1 & \xrightarrow{d^+} & \Omega^{2,+} \rightarrow 0. \\ & & b^0 & & b^1 & & b^+ \end{array}$$

Given a principal bundle P with connection A ,

$$0 \rightarrow \Omega^0(\mathfrak{g}_P) \xrightarrow{d_A} \Omega^1(\mathfrak{g}_P) \xrightarrow{d_A^+} \Omega^{2,+}(\mathfrak{g}_P) \rightarrow 0.$$

The double-composition is

$$d_A^+ \circ d_A = [F_A^+, \bullet].$$

This defines a complex when $F_A^+ = 0$.

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The de Rham complex

Latest version at <http://tensorial.com/math/>

Inside the de Rham complex

$$0 \rightarrow \Omega^0 \xrightarrow{d} \Omega^1 \xrightarrow{d} \Omega^2 \xrightarrow{d} \Omega^3 \xrightarrow{d} \Omega^4 \rightarrow 0$$

is the subcomplex

$$\begin{array}{ccccccc} 0 & \rightarrow & \Omega^0 & \xrightarrow{d} & \Omega^1 & \xrightarrow{d^+} & \Omega^{2,+} \rightarrow 0. \\ & & b^0 & & b^1 & & b^+ \end{array}$$

Given a principal bundle P with connection A ,

$$0 \rightarrow \Omega^0(\mathfrak{g}_P) \xrightarrow{d_A} \Omega^1(\mathfrak{g}_P) \xrightarrow{d_A^+} \Omega^{2,+}(\mathfrak{g}_P) \rightarrow 0.$$

The double-composition is

$$d_A^+ \circ d_A = [F_A^+, \bullet].$$

This defines a complex when $F_A^+ = 0$.

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Anti-self-dual equation

Latest version at <http://tensorial.com/math/>

People often study the equation $F_A^+ = 0$.

It is called the anti-self-dual equation since

$$F_A^+ = 0 \iff F_A = F_A^-.$$

Solutions arise as absolute minimizers of $\|F_A\|_{L^2}$.

If $g \in \mathcal{G}_P$ is a gauge transformation, then

$$F_{g(A)}^+ = g \cdot F_A^+ \cdot g^{-1},$$

so \mathcal{G}_P preserves solutions to $F_A^+ = 0$. The moduli space

$$\mathcal{M}_{\text{ASD}} = \{[A] \in \mathcal{A}_P / \mathcal{G}_P \mid F_A^+ = 0\}$$

is finite-dimensional. Roughly speaking, it defines a homology class in $\mathcal{A}_P / \mathcal{G}_P$. This leads to Donaldson invariants.

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Anti-self-dual equation

Latest version at <http://tensorial.com/math/>

People often study the equation $F_A^+ = 0$.

It is called the anti-self-dual equation since

$$F_A^+ = 0 \iff F_A = F_A^-.$$

Solutions arise as absolute minimizers of $\|F_A\|_{L^2}$.

If $g \in \mathcal{G}_P$ is a gauge transformation, then

$$F_{g(A)}^+ = g \cdot F_A^+ \cdot g^{-1},$$

so \mathcal{G}_P preserves solutions to $F_A^+ = 0$. The moduli space

$$\mathcal{M}_{\text{ASD}} = \{[A] \in \mathcal{A}_P / \mathcal{G}_P \mid F_A^+ = 0\}$$

is finite-dimensional. Roughly speaking, it defines a homology class in $\mathcal{A}_P / \mathcal{G}_P$. This leads to Donaldson invariants.

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Anti-self-dual equation

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

People often study the equation $F_A^+ = 0$.

It is called the anti-self-dual equation since

$$F_A^+ = 0 \iff F_A = F_A^-.$$

Solutions arise as absolute minimizers of $\|F_A\|_{L^2}$.

If $g \in \mathcal{G}_P$ is a gauge transformation, then

$$F_{g(A)}^+ = g \cdot F_A^+ \cdot g^{-1},$$

so \mathcal{G}_P preserves solutions to $F_A^+ = 0$. The moduli space

$$\mathcal{M}_{\text{ASD}} = \{[A] \in \mathcal{A}_P / \mathcal{G}_P \mid F_A^+ = 0\}$$

is finite-dimensional. Roughly speaking, it defines a homology class in $\mathcal{A}_P / \mathcal{G}_P$. This leads to Donaldson invariants.

Motivating question for studying Vafa-Witten

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- What is the Euler characteristic of the ASD moduli space \mathcal{M}_{ASD} ?
 - Is this question meaningful?
 - \mathcal{M}_{ASD} can be singular
 - \mathcal{M}_{ASD} depends on the choice of a metric
 - \mathcal{M}_{ASD} has multiple possible compactifications

Motivating question for studying Vafa-Witten

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- What is the Euler characteristic of the ASD moduli space \mathcal{M}_{ASD} ?
 - Is this question meaningful?
 - \mathcal{M}_{ASD} can be singular
 - \mathcal{M}_{ASD} depends on the choice of a metric
 - \mathcal{M}_{ASD} has multiple possible compactifications

Comparison with Donaldson invariants

Latest version at <http://tensorial.com/math/>

Write \mathcal{M}_{ASD} as a zero set:

$$\mathcal{M}_{\text{ASD}} = \{[A] \in \mathcal{A}/\mathcal{G} \mid F_A^+ = 0\}.$$

If $\dim(\mathcal{M}_{\text{ASD}}) = 0$, then the Donaldson invariant is a signed count $\#\mathcal{M}_{\text{ASD}}$.

Analogy with polynomials:

- Let $M = \{x \mid x^2 - c = 0\}$. How many points are in m ?
 - Signed count $\#M$ gives $+1 - 1 = 0$.
 - Generically well-defined on \mathbb{R} .
 - Unsigned count $\chi(M)$ gives $1 + 1 = 2$.
 - Generically well-defined on \mathbb{C} .

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Comparison with Donaldson invariants

Latest version at <http://tensorial.com/math/>

Write \mathcal{M}_{ASD} as a zero set:

$$\mathcal{M}_{\text{ASD}} = \{[A] \in \mathcal{A}/\mathcal{G} \mid F_A^+ = 0\}.$$

If $\dim(\mathcal{M}_{\text{ASD}}) = 0$, then the Donaldson invariant is a signed count $\#\mathcal{M}_{\text{ASD}}$.

Analogy with polynomials:

- Let $M = \{x \mid x^2 - c = 0\}$. How many points are in m ?
 - Signed count $\#M$ gives $+1 - 1 = 0$.
 - Generically well-defined on \mathbb{R} .
 - Unsigned count $\chi(M)$ gives $1 + 1 = 2$.
 - Generically well-defined on \mathbb{C} .

Comparison with Donaldson invariants

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Write \mathcal{M}_{ASD} as a zero set:

$$\mathcal{M}_{\text{ASD}} = \{[A] \in \mathcal{A}/\mathcal{G} \mid F_A^+ = 0\}.$$

If $\dim(\mathcal{M}_{\text{ASD}}) = 0$, then the Donaldson invariant is a signed count $\#\mathcal{M}_{\text{ASD}}$.

Analogy with polynomials:

- Let $M = \{x \mid x^2 - c = 0\}$. How many points are in m ?
 - Signed count $\#M$ gives $+1 - 1 = 0$.
 - Generically well-defined on \mathbb{R} .
 - Unsigned count $\chi(M)$ gives $1 + 1 = 2$.
 - Generically well-defined on \mathbb{C} .

Comparison with Donaldson invariants

Latest version at <http://tensorial.com/math/>

Write \mathcal{M}_{ASD} as a zero set:

$$\mathcal{M}_{\text{ASD}} = \{[A] \in \mathcal{A}/\mathcal{G} \mid F_A^+ = 0\}.$$

If $\dim(\mathcal{M}_{\text{ASD}}) = 0$, then the Donaldson invariant is a signed count $\#\mathcal{M}_{\text{ASD}}$.

Analogy with polynomials:

- Let $M = \{x \mid x^2 - c = 0\}$. How many points are in m ?
 - Signed count $\#M$ gives $+1 - 1 = 0$.
 - Generically well-defined on \mathbb{R} .
 - Unsigned count $\chi(M)$ gives $1 + 1 = 2$.
 - Generically well-defined on \mathbb{C} .

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

“Complexification” of configuration space

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

When $\dim(\mathcal{M}_{\text{ASD}}) = 0$, we expect the signed count of the Donaldson invariant $\#\mathcal{M}_{\text{ASD}}$ to typically differ from the unsigned count “ $\chi(\mathcal{M}_{\text{ASD}})$ ”.

In analogy with complexification, we will “double” the degrees of freedom in our configuration space by adding extra fields. This leads to an augmented moduli space \mathcal{M}_{VW} with

$$\mathcal{M}_{\text{ASD}} \subset \mathcal{M}_{\text{VW}}.$$

“Complexification” of configuration space

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

When $\dim(\mathcal{M}_{\text{ASD}}) = 0$, we expect the signed count of the Donaldson invariant $\#\mathcal{M}_{\text{ASD}}$ to typically differ from the unsigned count “ $\chi(\mathcal{M}_{\text{ASD}})$ ”.


In analogy with complexification, we will “double” the degrees of freedom in our configuration space by adding extra fields. This leads to an augmented moduli space \mathcal{M}_{VW} with

$$\mathcal{M}_{\text{ASD}} \subset \mathcal{M}_{\text{VW}}.$$

Counting zeroes of a section

Latest version at <http://tensorial.com/math/>

Consider a vector bundle $V \rightarrow X$ with a section $s : X \rightarrow V$.




Consider s as a vector field over the zero section in the total space.

Somehow extend the vector field to the total space...

Counting zeroes of a section

Latest version at <http://tensorial.com/math/>

Consider a vector bundle $V \rightarrow X$ with a section $s : X \rightarrow V$.




Consider s as a vector field over the zero section in the total space.

Somehow extend the vector field to the total space...

Counting zeroes of a section

Latest version at <http://tensorial.com/math/>

Consider a vector bundle $V \rightarrow X$ with a section $s : X \rightarrow V$.



Consider s as a vector field over the zero section in the total space.

Somehow extend the vector field to the total space...

Orienting the zeroes

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Consider the horizontal/vertical components of the derivative along the zero section:

$$\begin{pmatrix} 0 & \bullet \\ ds & \bullet \end{pmatrix}$$

(The horizontal component vanishes identically along the zero section.) We can choose the \bullet .

Choose the \bullet as

$$\begin{pmatrix} 0 & ds^* \\ ds & 0 \end{pmatrix}$$

to achieve consistently-signed determinant along the zero-section.

Orienting the zeroes

Latest version at <http://tensorial.com/math/>

Consider the horizontal/vertical components of the derivative along the zero section:

$$\begin{pmatrix} 0 & \bullet \\ ds & \bullet \end{pmatrix}$$

(The horizontal component vanishes identically along the zero section.) We can choose the \bullet .

Choose the \bullet as

$$\begin{pmatrix} 0 & ds^* \\ ds & 0 \end{pmatrix}$$

to achieve consistently-signed determinant along the zero-section.

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Vanishing theorem

Latest version at <http://tensorial.com/math/>

Ideally, our extended vector field will have no additional zeroes. This is the content of a “vanishing theorem.”



If a vanishing theorem holds, the zeroes of our vector field agree with the zeroes of our section, and the signed zero count of our vector field equals the unsigned zero count of our section.

Euler characteristic of \mathcal{M}_{ASD}

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

In this finite-dimensional analogy, \mathcal{M}_{ASD} is the zero-set of the section, and \mathcal{M}_{VW} is the zero-set of the vector field.

When a vanishing theorem holds, $\mathcal{M}_{\text{ASD}} = \mathcal{M}_{\text{VW}}$. In this case, we expect

$$“\#\mathcal{M}_{\text{VW}}” = “\chi(\mathcal{M}_{\text{ASD}}).”$$

What's wrong with $\chi(\mathcal{M}_{\text{ASD}})$?

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations


Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Poincaré-Hopf index theorem only computes the Euler characteristic of a compact manifold. Since \mathcal{M}_{ASD} is non-compact, we need a compactification $\overline{\mathcal{M}}_{\text{ASD}}$:



The invariant should be independent of the metric g , but different choices of g typically lead to *cobordant* compactified ASD moduli spaces $\overline{\mathcal{M}}_{\text{ASD}}(g)$.

Euler characteristic is *not* invariant under cobordism!
(\mathbb{S}^2)

What's wrong with $\chi(\mathcal{M}_{\text{ASD}})$?

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations


Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Poincaré-Hopf index theorem only computes the Euler characteristic of a compact manifold. Since \mathcal{M}_{ASD} is non-compact, we need a compactification $\overline{\mathcal{M}}_{\text{ASD}}$:



The invariant should be independent of the metric g , but different choices of g typically lead to *cobordant* compactified ASD moduli spaces $\overline{\mathcal{M}}_{\text{ASD}}(g)$.

Euler characteristic is *not* invariant under cobordism!
(S^2)

What's wrong with $\chi(\mathcal{M}_{\text{ASD}})$?

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations


Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Poincaré-Hopf index theorem only computes the Euler characteristic of a compact manifold. Since \mathcal{M}_{ASD} is non-compact, we need a compactification $\overline{\mathcal{M}}_{\text{ASD}}$:



The invariant should be independent of the metric g , but different choices of g typically lead to *cobordant* compactified ASD moduli spaces $\overline{\mathcal{M}}_{\text{ASD}}(g)$.

Euler characteristic is *not* invariant under cobordism!
(\mathbb{S}^2)

What's wrong with $\chi(\mathcal{M}_{\text{ASD}})$?

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations


Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Poincaré-Hopf index theorem only computes the Euler characteristic of a compact manifold. Since \mathcal{M}_{ASD} is non-compact, we need a compactification $\overline{\mathcal{M}}_{\text{ASD}}$:



The invariant should be independent of the metric g , but different choices of g typically lead to *cobordant* compactified ASD moduli spaces $\overline{\mathcal{M}}_{\text{ASD}}(g)$.

Euler characteristic is *not* invariant under cobordism!
(\mathbb{S}^2)

What's wrong with $\chi(\mathcal{M}_{\text{ASD}})$?

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations


Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Poincaré-Hopf index theorem only computes the Euler characteristic of a compact manifold. Since \mathcal{M}_{ASD} is non-compact, we need a compactification $\overline{\mathcal{M}}_{\text{ASD}}$:




The invariant should be independent of the metric g , but different choices of g typically lead to *cobordant* compactified ASD moduli spaces $\overline{\mathcal{M}}_{\text{ASD}}(g)$.

Euler characteristic is *not* invariant under cobordism!
(\mathbb{S}^2)

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Behavior of families

Latest version at <http://tensorial.com/math/>

Consider $0 = x^2 - c$ as c goes from positive to negative:



Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Alternative behavior

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

We could have extended the vector field differently.

Alternative behavior

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation


The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing



Alternative behavior

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation


The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing



Alternative behavior

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation


The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing



Alternative behavior

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation


The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing



Alternative behavior

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation


The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing



Alternative behavior

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation


The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing



Alternative behavior

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation


The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing



What is the purpose of extra fields?

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations


Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Consider a one-parameter family of metrics $\{g_t\}$ for $t \in \mathbb{R}$.



As the Euler characteristic of $\overline{\mathcal{M}}_{\text{ASD}}$ changes, points in \mathcal{M}_{VW} should be created or destroyed to compensate.

Additional pathologies

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The previous picture is a fantasy. The crux of this program is to understand how to deal with pathologies.




- Sequences of solutions in $\overline{\mathcal{M}}_{VW}$ could have unbounded L^2 norms.
- Rays appear in $\overline{\mathcal{M}}_{VW}$ at reducible points of $\overline{\mathcal{M}}_{ASD}$.
- Despite having expected dimension zero, there are often manifolds of non-ASD solutions

Additional pathologies

Latest version at <http://tensorial.com/math/>

The previous picture is a fantasy. The crux of this program is to understand how to deal with pathologies.




- Sequences of solutions in $\overline{\mathcal{M}}_{VW}$ could have unbounded L^2 norms.
- Rays appear in $\overline{\mathcal{M}}_{VW}$ at reducible points of $\overline{\mathcal{M}}_{ASD}$.
- Despite having expected dimension zero, there are often manifolds of non-ASD solutions

Additional pathologies

Latest version at <http://tensorial.com/math/>

The previous picture is a fantasy. The crux of this program is to understand how to deal with pathologies.



- Sequences of solutions in $\overline{\mathcal{M}}_{VW}$ could have unbounded L^2 norms.
- Rays appear in $\overline{\mathcal{M}}_{VW}$ at reducible points of $\overline{\mathcal{M}}_{ASD}$.
- Despite having expected dimension zero, there are often manifolds of non-ASD solutions

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Additional pathologies

Latest version at <http://tensorial.com/math/>

The previous picture is a fantasy. The crux of this program is to understand how to deal with pathologies.



- Sequences of solutions in $\overline{\mathcal{M}}_{VW}$ could have unbounded L^2 norms.
- Rays appear in $\overline{\mathcal{M}}_{VW}$ at reducible points of $\overline{\mathcal{M}}_{ASD}$.
- Despite having expected dimension zero, there are often manifolds of non-ASD solutions

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Atiyah-Jeffrey supersymmetry

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

There is an Atiyah-Jeffrey style supersymmetric path integral expression for the Euler characteristic of \mathcal{M}_{ASD} .

$$“\chi(\mathcal{M}_{\text{ASD}})” = “\int e^{-L}.”$$

Vafa and Witten recognized Yamron's twist of $\mathcal{N} = 4$ supersymmetric Yang-Mills as such.

They were studying $\mathcal{N} = 4$ supersymmetry in the context of S-duality.

S-duality and geometric Langlands

Latest version at <http://tensorial.com/math/>

In this context, S-duality roughly means that the generating function

$$\sum_k \chi(\mathcal{M}_{\text{ASD}}(k)) q^k$$

should be a modular form.

In several specific examples, they “computed” these generating functions and verified their modularity.

This Vafa-Witten theory is one of three twists of $\mathcal{N} = 4$ supersymmetric Yang-Mills theory. In 2006, Kapustin and Witten explored the relation of another such twist is to geometric Langlands. More recently, the Vafa-Witten twist has appeared in the work of Haydys and Witten on five-dimensional gauge theory.

S-duality and geometric Langlands

Latest version at <http://tensorial.com/math/>

In this context, S-duality roughly means that the generating function

$$\sum_k \chi(\mathcal{M}_{\text{ASD}}(k)) q^k$$

should be a modular form.

In several specific examples, they “computed” these generating functions and verified their modularity.

This Vafa-Witten theory is one of three twists of $\mathcal{N} = 4$ supersymmetric Yang-Mills theory. In 2006, Kapustin and Witten explored the relation of another such twist is to geometric Langlands. More recently, the Vafa-Witten twist has appeared in the work of Haydys and Witten on five-dimensional gauge theory.

S-duality and geometric Langlands

Latest version at <http://tensorial.com/math/>

In this context, S-duality roughly means that the generating function

$$\sum_k \chi(\mathcal{M}_{\text{ASD}}(k)) q^k$$

should be a modular form.

In several specific examples, they “computed” these generating functions and verified their modularity.

This Vafa-Witten theory is one of three twists of $\mathcal{N} = 4$ supersymmetric Yang-Mills theory. In 2006, Kapustin and Witten explored the relation of another such twist is to geometric Langlands. More recently, the Vafa-Witten twist has appeared in the work of Haydys and Witten on five-dimensional gauge theory.

Explicit example of S-duality

Latest version at <http://tensorial.com/math/>

Consider the four-manifold $X = K3$. The generating functions for $G = \mathrm{SU}(2)$ and $\hat{G} = \mathrm{SO}(3)$ are

$$Z_{\mathrm{SU}(2)}(q) = \frac{1}{2}q^{-2}\left(\frac{1}{4} + 0q + 30q^2 + 3200q^3 + \dots\right. \\ \left. \dots + \frac{10189790756178504975}{4}q^{16} + \dots\right)$$

$$Z_{\mathrm{SO}(3)}(q) = q^{-2}\left(\frac{1}{4} + 0q^{1/2} + 0q + 2096128q^{3/2} + \dots\right. \\ \left. + 50356230q^2 + 679145472q^{5/2} + \dots\right. \\ \left. \dots + \frac{21379974409572270922824975}{4}q^{16} + \dots\right)$$

Define $q^{1/2} = e^{i\pi\tau}$. In this case, S-duality is the “modular relation”

$$Z_{\mathrm{SU}(2)}(-1/\tau) = (2\tau)^{-12} Z_{\mathrm{SO}(3)}(\tau).$$

Explicit example of S-duality

Latest version at <http://tensorial.com/math/>

Consider the four-manifold $X = K3$. The generating functions for $G = \mathrm{SU}(2)$ and $\hat{G} = \mathrm{SO}(3)$ are

$$Z_{\mathrm{SU}(2)}(q) = \frac{1}{2}q^{-2}\left(\frac{1}{4} + 0q + 30q^2 + 3200q^3 + \dots\right. \\ \left. \dots + \frac{10189790756178504975}{4}q^{16} + \dots\right)$$

$$Z_{\mathrm{SO}(3)}(q) = q^{-2}\left(\frac{1}{4} + 0q^{1/2} + 0q + 2096128q^{3/2} + \dots\right. \\ \left. + 50356230q^2 + 679145472q^{5/2} + \dots\right. \\ \left. \dots + \frac{21379974409572270922824975}{4}q^{16} + \dots\right)$$

Define $q^{1/2} = e^{i\pi\tau}$. In this case, S-duality is the “modular relation”

$$Z_{\mathrm{SU}(2)}(-1/\tau) = (2\tau)^{-12}Z_{\mathrm{SO}(3)}(\tau).$$

The equations

Latest version at <http://tensorial.com/math/>

The equations

$$F_A^+ - \frac{1}{4} [B \times B] - \frac{1}{2} [C, B] = 0$$

$$d_A C + d_A^* B = 0$$

Let $P \rightarrow X^4$ be a principal bundle over an oriented Riemannian four-manifold. A *configuration* (C, A, B) consists of

- A section of the adjoint bundle $C \in \Omega^0(M; \mathfrak{g}_P)$
- A connection $A \in \mathcal{A}_P$
- An adjoint-valued self-dual two-form $B \in \Omega^{2,+}(M; \mathfrak{g}_P)$

The quadratic term

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The equations

$$F_A^+ - \frac{1}{4}[B \times B] - \frac{1}{2}[C, B] = 0$$

$$d_A C + d_A^* B = 0$$

This quadratic term on $\mathfrak{g} \otimes \Lambda^{2,+}$ is the tensor product of the Lie bracket and the cross product.

Since $[\cdot, \cdot]$ is antisymmetric on \mathfrak{g} and \times is antisymmetric on $\Lambda^{2,+}$, their product $[B_1 \times B_2]$ is *symmetric*.

Only semi-definite

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Note that $[B \times B]$ has a nontrivial kernel. Later we will see that this has dire consequences.

For example, if B has rank one

$$B = \chi \otimes \sigma^1,$$

then

$$[B \times B] = [\chi, \chi] \otimes (\sigma^1 \times \sigma^1) = 0 \otimes 0.$$

The quartic form $|[B \times B]|^2$ is only semi-definite.

Only semi-definite

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Note that $[B \times B]$ has a nontrivial kernel. Later we will see that this has dire consequences.

For example, if B has rank one

$$B = \chi \otimes \sigma^1,$$

then

$$[B \times B] = [\chi, \chi] \otimes (\sigma^1 \times \sigma^1) = 0 \otimes 0.$$

The quartic form $||[B \times B]||^2$ is only semi-definite.

The standard compactness strategy

Latest version at <http://tensorial.com/math/>

- Use energy identities to establish a priori L_1^2 bounds
- These L_1^2 bounds imply weak compactness (Hodge theory for abelian case, Uhlenbeck/Sedlacek theory for non-abelian case)
- Elliptic regularity implies strong (Uhlenbeck) compactness

Summary

Using established analytic machinery, *a priori* L_1^2 bounds imply compactness

Examples: ASD, Seiberg-Witten, $PU(2)$ monopoles

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The standard compactness strategy

Latest version at <http://tensorial.com/math/>

- Use energy identities to establish a priori L_1^2 bounds
- These L_1^2 bounds imply weak compactness (Hodge theory for abelian case, Uhlenbeck/Sedlacek theory for non-abelian case)
- Elliptic regularity implies strong (Uhlenbeck) compactness

Summary

Using established analytic machinery, *a priori* L_1^2 bounds imply compactness

Examples: ASD, Seiberg-Witten, $PU(2)$ monopoles

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The standard compactness strategy

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- Use energy identities to establish a priori L_1^2 bounds
- These L_1^2 bounds imply weak compactness (Hodge theory for abelian case, Uhlenbeck/Sedlacek theory for non-abelian case)
- Elliptic regularity implies strong (Uhlenbeck) compactness

Summary

Using established analytic machinery, *a priori* L_1^2 bounds imply compactness

Examples: ASD, Seiberg-Witten, $PU(2)$ monopoles

The standard compactness strategy

Latest version at <http://tensorial.com/math/>

- Use energy identities to establish a priori L_1^2 bounds
- These L_1^2 bounds imply weak compactness (Hodge theory for abelian case, Uhlenbeck/Sedlacek theory for non-abelian case)
- Elliptic regularity implies strong (Uhlenbeck) compactness

Summary

Using established analytic machinery, *a priori* L_1^2 bounds imply compactness

Examples: ASD, Seiberg-Witten, PU(2) monopoles

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Seiberg-Witten equations

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\begin{aligned}F_A^+ - \frac{1}{4}(\phi \otimes \phi^*)_0 &= 0, \\ \not{D}_A \phi &= 0.\end{aligned}$$

P is a $U(1)$ bundle with connection A and associated line bundle L .

ϕ is a section of the twisted spinor bundle $\mathcal{S} \otimes L$.

Clifford multiplication identifies F_A^+ with a traceless endomorphism of $\mathcal{S} \otimes L$.

Define energy as the sum of squares:

$$\mathcal{E}_{\text{SW}}(A, \phi) := \int_X |F_A^+ - \frac{1}{4}(\phi \otimes \phi^*)_0|^2 + \int_X |\not{D}_A \phi|^2.$$

The Seiberg-Witten equations

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\begin{aligned}F_A^+ - \frac{1}{4}(\phi \otimes \phi^*)_0 &= 0, \\ \not{D}_A \phi &= 0.\end{aligned}$$

P is a $U(1)$ bundle with connection A and associated line bundle L .

ϕ is a section of the twisted spinor bundle $\mathbb{S} \otimes L$.

Clifford multiplication identifies F_A^+ with a traceless endomorphism of $\mathbb{S} \otimes L$.

Define energy as the sum of squares:

$$\mathcal{E}_{\text{SW}}(A, \phi) := \int_X |F_A^+ - \frac{1}{4}(\phi \otimes \phi^*)_0|^2 + \int_X |\not{D}_A \phi|^2.$$

Energy identities for Seiberg-Witten

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\mathcal{E}_{\text{SW}}(A, \phi) := \int_X |F_A^+ - \tfrac{1}{4}(\phi \otimes \phi^*)_0|^2 + \int_X |\not\partial_A \phi|^2.$$

The Weitzenböck formula for spinors gives

$$\int_X |\not\partial_A \phi|^2 = \int_X (|\nabla_A \phi|^2 + \tfrac{1}{4}s|\phi|^2 - \tfrac{1}{2}\langle \phi, F_A^+ \cdot \phi \rangle).$$

Expanding out \mathcal{E}_{SW} gives

$$\begin{aligned} \mathcal{E}_{\text{SW}} = & \int_X (|F_A^+|^2 + \tfrac{1}{16}|(\phi \otimes \phi^*)_0|^2 + \cancel{\tfrac{1}{2}\langle F_A^+, (\phi \otimes \phi^*)_0 \rangle}) \\ & + \int_X (|\nabla_A \phi|^2 + \tfrac{1}{4}s|\phi|^2 - \cancel{\tfrac{1}{2}\langle \phi, F_A^+ \cdot \phi \rangle}) \end{aligned}$$

$$\mathcal{E}_{\text{SW}} = \int_X (|F_A^+|^2 + |\nabla_A \phi|^2 + \tfrac{1}{32}|\phi|^4 + \tfrac{1}{4}s|\phi|^2)$$

Energy identities for Seiberg-Witten

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\mathcal{E}_{\text{SW}}(A, \phi) := \int_X |F_A^+ - \tfrac{1}{4}(\phi \otimes \phi^*)_0|^2 + \int_X |\not\partial_A \phi|^2.$$

The Weitzenböck formula for spinors gives

$$\int_X |\not\partial_A \phi|^2 = \int_X (|\nabla_A \phi|^2 + \tfrac{1}{4}s|\phi|^2 - \tfrac{1}{2}\langle \phi, F_A^+ \cdot \phi \rangle).$$

Expanding out \mathcal{E}_{SW} gives

$$\begin{aligned} \mathcal{E}_{\text{SW}} = & \int_X (|F_A^+|^2 + \tfrac{1}{16}|(\phi \otimes \phi^*)_0|^2 + \cancel{\tfrac{1}{2}\langle F_A^+, (\phi \otimes \phi^*)_0 \rangle}) \\ & + \int_X (|\nabla_A \phi|^2 + \tfrac{1}{4}s|\phi|^2 - \cancel{\tfrac{1}{2}\langle \phi, F_A^+ \cdot \phi \rangle}) \end{aligned}$$

$$\mathcal{E}_{\text{SW}} = \int_X (|F_A^+|^2 + |\nabla_A \phi|^2 + \tfrac{1}{32}|\phi|^4 + \tfrac{1}{4}s|\phi|^2)$$

Energy identities for Seiberg-Witten

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\mathcal{E}_{\text{SW}}(A, \phi) := \int_X |F_A^+ - \tfrac{1}{4}(\phi \otimes \phi^*)_0|^2 + \int_X |\not\partial_A \phi|^2.$$

The Weitzenböck formula for spinors gives

$$\int_X |\not\partial_A \phi|^2 = \int_X (|\nabla_A \phi|^2 + \tfrac{1}{4}s|\phi|^2 - \tfrac{1}{2}\langle \phi, F_A^+ \cdot \phi \rangle).$$

Expanding out \mathcal{E}_{SW} gives

$$\begin{aligned} \mathcal{E}_{\text{SW}} = & \int_X (|F_A^+|^2 + \tfrac{1}{16}|(\phi \otimes \phi^*)_0|^2 + \cancel{\tfrac{1}{2}\langle F_A^+, (\phi \otimes \phi^*)_0 \rangle}) \\ & + \int_X (|\nabla_A \phi|^2 + \tfrac{1}{4}s|\phi|^2 - \cancel{\tfrac{1}{2}\langle \phi, F_A^+ \cdot \phi \rangle}) \end{aligned}$$

$$\mathcal{E}_{\text{SW}} = \int_X (|F_A^+|^2 + |\nabla_A \phi|^2 + \tfrac{1}{32}|\phi|^4 + \tfrac{1}{4}s|\phi|^2)$$

Completing the square

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\mathcal{E}_{\text{SW}} = \int_X \left(|F_A^+|^2 + |\nabla_A \phi|^2 + \frac{1}{32} |\phi|^4 + \frac{1}{4} s |\phi|^2 \right).$$

Problem When scalar curvature is negative, the term $\int_X \frac{1}{4} s |\phi|^2$ could be large and negative.

Solution Complete the square:

$$\begin{aligned} \mathcal{E}_{\text{SW}} &= \int_X \left(|F_A^+|^2 + |\nabla_A \phi|^2 + \frac{1}{32} (|\phi|^2 + 4s)^2 - \frac{1}{2} s^2 \right) \\ &= \int_X \left(\frac{1}{2} |F_A|^2 + |\nabla_A \phi|^2 + \frac{1}{32} (|\phi|^2 + 4s)^2 \right) - \kappa - \int_X \frac{1}{2} s^2. \end{aligned}$$

Completing the square

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\mathcal{E}_{\text{SW}} = \int_X \left(|F_A^+|^2 + |\nabla_A \phi|^2 + \frac{1}{32} |\phi|^4 + \frac{1}{4} s |\phi|^2 \right).$$

Problem When scalar curvature is negative, the term $\int_X \frac{1}{4} s |\phi|^2$ could be large and negative.

Solution Complete the square:

$$\begin{aligned} \mathcal{E}_{\text{SW}} &= \int_X \left(|F_A^+|^2 + |\nabla_A \phi|^2 + \frac{1}{32} (|\phi|^2 + 4s)^2 - \frac{1}{2} s^2 \right) \\ &= \int_X \left(\frac{1}{2} |F_A|^2 + |\nabla_A \phi|^2 + \frac{1}{32} (|\phi|^2 + 4s)^2 \right) - \kappa - \int_X \frac{1}{2} s^2. \end{aligned}$$

Compactness for Seiberg-Witten

Latest version at <http://tensorial.com/math/>

A priori estimate

$$\mathcal{E}_{\text{SW}} = 0 \iff$$

$$\int_X \left(\frac{1}{2} |F_A|^2 + |\nabla_A \phi|^2 + \frac{1}{32} (|\phi|^2 + 4s)^2 \right) = \kappa + \int_X \frac{1}{2} s^2.$$

The left hand side is a sum of positive terms. The quartic term is essentially $\|\phi\|_{L^4}^4$.

The right hand side depends only on the (fixed) topology of the bundle, and geometry of the manifold.

Since the gauge group is abelian, elementary Hodge theory yields genuine L_1^2 bounds on A and ϕ , and elliptic bootstrapping implies compactness.

Compactness for Seiberg-Witten

Latest version at <http://tensorial.com/math/>

A priori estimate

$$\mathcal{E}_{\text{SW}} = 0 \iff$$

$$\int_X \left(\frac{1}{2} |F_A|^2 + |\nabla_A \phi|^2 + \frac{1}{32} (|\phi|^2 + 4s)^2 \right) = \kappa + \int_X \frac{1}{2} s^2.$$

The left hand side is a sum of positive terms. The quartic term is essentially $\|\phi\|_{L^4}^4$.

The right hand side depends only on the (fixed) topology of the bundle, and geometry of the manifold.

Since the gauge group is abelian, elementary Hodge theory yields genuine L_1^2 bounds on A and ϕ , and elliptic bootstrapping implies compactness.

Compactness for Seiberg-Witten

Latest version at <http://tensorial.com/math/>

A priori estimate

$$\mathcal{E}_{\text{SW}} = 0 \iff$$

$$\int_X \left(\frac{1}{2} |F_A|^2 + |\nabla_A \phi|^2 + \frac{1}{32} (|\phi|^2 + 4s)^2 \right) = \kappa + \int_X \frac{1}{2} s^2.$$

The left hand side is a sum of positive terms. The quartic term is essentially $\|\phi\|_{L^4}^4$.

The right hand side depends only on the (fixed) topology of the bundle, and geometry of the manifold.

Since the gauge group is abelian, elementary Hodge theory yields genuine L_1^2 bounds on A and ϕ , and elliptic bootstrapping implies compactness.

Compactness for Seiberg-Witten

Latest version at <http://tensorial.com/math/>

A priori estimate

$$\mathcal{E}_{\text{SW}} = 0 \iff$$

$$\int_X \left(\frac{1}{2} |F_A|^2 + |\nabla_A \phi|^2 + \frac{1}{32} (|\phi|^2 + 4s)^2 \right) = \kappa + \int_X \frac{1}{2} s^2.$$

The left hand side is a sum of positive terms. The quartic term is essentially $\|\phi\|_{L^4}^4$.

The right hand side depends only on the (fixed) topology of the bundle, and geometry of the manifold.

Since the gauge group is abelian, elementary Hodge theory yields genuine L_1^2 bounds on A and ϕ , and elliptic bootstrapping implies compactness.

Energy identities for Vafa-Witten

Latest version at <http://tensorial.com/math/>

We emulate the standard approach:

$$\begin{aligned}\mathcal{E}_{\text{VW}}(C, A, B) &:= \frac{1}{2} \|d_A C + d_A^* B\|^2 + \|F_A^+ - \frac{1}{4} [B \times B] - \frac{1}{2} [C, B]\|^2 \\ &= \frac{1}{2} \|d_A C\|^2 + \frac{1}{2} \|d_A^* B\|^2 + \|F_A^+ - \frac{1}{4} [B \times B]\|^2 + \frac{1}{4} \|[C, B]\|^2 \\ &\quad + \int_X (\langle d_A C, d_A^* B \rangle - \langle F_A^+, [C, B] \rangle) + \int_X \frac{1}{4} \langle [B \times B], [C, B] \rangle.\end{aligned}$$

The bottom line cancels since

$$\langle F_A^+, [C, B] \rangle = \langle [F_A^+, C], B \rangle = \langle d_A d_A C, B \rangle,$$

and the Jacobi identity implies

$$[[B \times B] \cdot B] = 0.$$

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Energy identities for Vafa-Witten

Latest version at <http://tensorial.com/math/>

We emulate the standard approach:

$$\begin{aligned}\mathcal{E}_{\text{VW}}(C, A, B) &:= \frac{1}{2} \|d_A C + d_A^* B\|^2 + \|F_A^+ - \frac{1}{4} [B \times B] - \frac{1}{2} [C, B]\|^2 \\ &= \frac{1}{2} \|d_A C\|^2 + \frac{1}{2} \|d_A^* B\|^2 + \|F_A^+ - \frac{1}{4} [B \times B]\|^2 + \frac{1}{4} \|[C, B]\|^2 \\ &\quad + \int_X (\langle d_A C, d_A^* B \rangle - \langle F_A^+, [C, B] \rangle) + \int_X \frac{1}{4} \langle [B \times B], [C, B] \rangle.\end{aligned}$$

The bottom line cancels since

$$\langle F_A^+, [C, B] \rangle = \langle [F_A^+, C], B \rangle = \langle d_A d_A C, B \rangle,$$

and the Jacobi identity implies

$$[[B \times B] \cdot B] = 0.$$

Energy identities for Vafa-Witten

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

We emulate the standard approach:

$$\begin{aligned}\mathcal{E}_{\text{VW}}(C, A, B) &:= \frac{1}{2} \|d_A C + d_A^* B\|^2 + \|F_A^+ - \frac{1}{4} [B \times B] - \frac{1}{2} [C, B]\|^2 \\ &= \frac{1}{2} \|d_A C\|^2 + \frac{1}{2} \|d_A^* B\|^2 + \|F_A^+ - \frac{1}{4} [B \times B]\|^2 + \frac{1}{4} \|[C, B]\|^2 \\ &\quad + \int_X (\langle d_A C, d_A^* B \rangle - \langle F_A^+, [C, B] \rangle) + \int_X \frac{1}{4} \langle [B \times B], [C, B] \rangle.\end{aligned}$$

The bottom line cancels since

$$\langle F_A^+, [C, B] \rangle = \langle [F_A^+, C], B \rangle = \langle d_A d_A C, B \rangle,$$

and the Jacobi identity implies

$$[[B \times B] \cdot B] = 0.$$

Simplification

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Thus, assuming that the base manifold X is closed, we have the identity

$$\mathcal{E}_{\text{VW}} = \frac{1}{2} \|d_A C\|^2 + \frac{1}{2} \|d_A^* B\|^2 + \|F_A^+ - \frac{1}{4} [B \times B]\|^2 + \frac{1}{4} \|[C, B]\|^2.$$

This is a different sum of squares, equivalent equations are

$$\begin{aligned} F_A^+ - \frac{1}{4} [B \times B] &= 0, & [C, B] &= 0, \\ d_A^* B &= 0, & d_A C &= 0. \end{aligned}$$

These equations are linear in C . The interesting nonlinear part with B decouples. WLOG, set $C = 0$.

Simplification

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Thus, assuming that the base manifold X is closed, we have the identity

$$\mathcal{E}_{\text{VW}} = \frac{1}{2} \|d_A C\|^2 + \frac{1}{2} \|d_A^* B\|^2 + \|F_A^+ - \frac{1}{4} [B \times B]\|^2 + \frac{1}{4} \|[C, B]\|^2.$$

This is a different sum of squares, equivalent equations are

$$\begin{aligned} F_A^+ - \frac{1}{4} [B \times B] &= 0, & [C, B] &= 0, \\ d_A^* B &= 0, & d_A C &= 0. \end{aligned}$$

These equations are linear in C . The interesting nonlinear part with B decouples. WLOG, set $C = 0$.

Analogy with Seiberg-Witten

Latest version at <http://tensorial.com/math/>

The equations

$$\begin{aligned}F_A^+ &= \frac{1}{4} [B \times B] \\ d_A^* B &= 0\end{aligned}$$

These equations say that B has a harmonic square root, if we interpret “ B^2 ” = $[B \times B]$.

$$\begin{aligned}B &= “2\sqrt{F_A^+}” \\ d_A^* B &= 0 \quad (\Rightarrow d_A B = 0)\end{aligned}$$

Contrast this with the Seiberg-Witten equations

$$\begin{aligned}F_A^+ - (\phi \otimes \phi^*)_0 &= 0 \\ \not{D}_A \phi &= 0\end{aligned}$$

Analogy with Seiberg-Witten

Latest version at <http://tensorial.com/math/>

The equations

$$\begin{aligned}F_A^+ &= \frac{1}{4} [B \times B] \\ d_A^* B &= 0\end{aligned}$$

These equations say that B has a harmonic square root, if we interpret “ B^2 ” = $[B \times B]$.

$$\begin{aligned}B &= “2\sqrt{F_A^+}” \\ d_A^* B &= 0 \quad (\Rightarrow d_A B = 0)\end{aligned}$$

Contrast this with the Seiberg-Witten equations

$$\begin{aligned}F_A^+ - (\phi \otimes \phi^*)_0 &= 0 \\ \not{D}_A \phi &= 0\end{aligned}$$

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Weitzenböck formula

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\frac{1}{2} \|d_A^* B\|^2 = \frac{1}{4} \|\nabla_A B\|^2 + \int_X \left(\frac{1}{2} \langle B, [F_A^+ \times B] \rangle + \left(\frac{1}{12} s - \frac{1}{2} W^+ \right) \cdot \langle B \otimes B \rangle \right).$$

$$\mathcal{E}_{\text{VW}}(0, A, B) = \|F_A^+ - \frac{1}{4} [B \times B]\|^2 + \frac{1}{2} \|d_A^* B\|^2.$$

Once again, the cross-term miraculously cancels:

$$\mathcal{E}_{\text{VW}} = \|F_A^+\|^2 + \frac{1}{16} \|[B \times B]\|^2 + \frac{1}{4} \|\nabla_A B\|^2 + \int_X \frac{1}{2} \left(\langle B, [F_A^+ \times B] \rangle - \langle F_A^+, [B \times B] \rangle \right) + \int_X \left(\frac{1}{12} s - \frac{1}{2} W^+ \right) \cdot \langle B \otimes B \rangle.$$

The Weitzenböck formula

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\frac{1}{2} \|d_A^* B\|^2 = \frac{1}{4} \|\nabla_A B\|^2 + \int_X \left(\frac{1}{2} \langle B, [F_A^+ \times B] \rangle + \left(\frac{1}{12} s - \frac{1}{2} W^+ \right) \cdot \langle B \otimes B \rangle \right).$$

$$\mathcal{E}_{\text{VW}}(0, A, B) = \|F_A^+ - \frac{1}{4} [B \times B]\|^2 + \frac{1}{2} \|d_A^* B\|^2.$$

Once again, the cross-term miraculously cancels:

$$\mathcal{E}_{\text{VW}} = \|F_A^+\|^2 + \frac{1}{16} \|[B \times B]\|^2 + \frac{1}{4} \|\nabla_A B\|^2 + \int_X \frac{1}{2} \left(\langle B, [F_A^+ \times B] \rangle - \langle F_A^+, [B \times B] \rangle \right) + \int_X \left(\frac{1}{12} s - \frac{1}{2} W^+ \right) \cdot \langle B \otimes B \rangle.$$

Vanishing theorem

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Vafa-Witten equations (with $C = 0$) are equivalent to

$$0 = \frac{1}{2} \|F_A^+\|^2 + \frac{1}{4} \|\nabla_A B\|^2 + \frac{1}{16} \|[B \times B]\|^2 + \\ + \int_X \left(\frac{1}{12} s - \frac{1}{2} W^+ \right) \cdot \langle B \otimes B \rangle.$$

If furthermore the curvature part is positive semi-definite, then M must be Kähler, hyper-Kähler, or $b^+ = 0$, and the equations decouple further to

$$F_A^+ = 0 \quad \nabla_A B = 0 \quad [B \times B] = 0.$$

Vanishing theorem

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

The Vafa-Witten equations (with $C = 0$) are equivalent to

$$0 = \frac{1}{2} \|F_A^+\|^2 + \frac{1}{4} \|\nabla_A B\|^2 + \frac{1}{16} \|[B \times B]\|^2 + \\ + \int_X \left(\frac{1}{12} s - \frac{1}{2} W^+ \right) \cdot \langle B \otimes B \rangle.$$

If furthermore the curvature part is positive semi-definite, then M must be Kähler, hyper-Kähler, or $b^+ = 0$, and the equations decouple further to

$$F_A^+ = 0 \quad \nabla_A B = 0 \quad [B \times B] = 0.$$

Kähler manifolds

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Let (M, g, ω) be a closed Kähler manifold. The equations

$$F_A^+ - \frac{1}{4} [B \times B] = 0$$

$$d_A^* B = 0$$

reduce to

$$\beta \in \Omega^{2,0}(X; \mathfrak{g}_P \otimes \mathbb{C}), \quad B = \beta - \beta^*, \quad \bar{\partial}_A \beta = 0,$$

$$\omega \wedge iF_A + \frac{1}{2} [\beta \wedge \beta^*] = 0.$$

Note the extra symmetry $\beta \mapsto e^{i\theta} \beta$ for θ constant.

Semistability

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

We say a subbundle $E' \subset E$ is β -invariant if

$$\beta(E') \subset E' \otimes K.$$

Suppose E is a Hermitian vector bundle, A is a holomorphic connection on E , the traceless part of F_A is F_A^0 , and $\beta \in \Omega^{2,0}(\text{End}(E))$ satisfies

$$\omega \wedge iF_A^0 + \frac{1}{2} [\beta \wedge \beta^*] = 0,$$

then E is β -semistable.

Failure of a priori bound

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Recall that our quartic term $|[B \cdot B]|^2$ is only positive *semi*-definite. If it were positive-definite, then it would dominate the curvature part, and the identity

$$0 = \frac{1}{2} \|F_A\|^2 + \frac{1}{4} \|\nabla_A B\|^2 + \frac{1}{16} \|[B \times B]\|^2 + \\ + \int_X \left(\frac{1}{12} s - \frac{1}{2} W^+ \right) \cdot \langle B \otimes B \rangle - \kappa$$

would yield a priori bounds on $\|F_A\|$, $\|\nabla_A B\|$, and $\|B\|_{L^4}$. Instead, we get no such bounds since $|[B \times B]|^2$ could vanish while the curvature terms go to $-\infty$ unchecked.

Scaling the metric

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Although these equations are not conformally invariant, there is a scaling law.

For all $\eta \in \mathbb{R}$, the space of solutions is invariant under

$$(C, A, B, g) \mapsto (e^{-\eta}C, A, e^{\eta}B, e^{2\eta}g).$$

Thus rescaling the metric can be absorbed by this rescaling of B and C .

More Weitzenböck

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

For a more concrete application of the width heuristic, consider the following identity for solutions:

$$\frac{1}{8} \Delta |B|^2 + \frac{1}{4} |\nabla_A B|^2 + \frac{1}{8} |[B \times B]|^2 = \langle B \cdot (-\frac{1}{12} s + \frac{1}{2} W^+) B \rangle$$

In particular,

$$\Delta |B|^2 \leq \lambda |B|^2$$

where λ depends on curvature.

With slightly more work,

$$\Delta |B| \leq \lambda |B|$$

More Weitzenböck

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

For a more concrete application of the width heuristic, consider the following identity for solutions:

$$\frac{1}{8} \Delta |B|^2 + \frac{1}{4} |\nabla_A B|^2 + \frac{1}{8} |[B \times B]|^2 = \langle B \cdot (-\frac{1}{12} s + \frac{1}{2} W^+) B \rangle$$

In particular,

$$\Delta |B|^2 \leq \lambda |B|^2$$

where λ depends on curvature.

With slightly more work,

$$\Delta |B| \leq \lambda |B|$$

Bounding B

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Thanks to a mean-value inequality due to Morrey

$$\Delta |B| \leq \lambda |B| \implies \|B\|_{L^\infty} \leq c \|B\|_{L^2}$$

Thus

$$\|F_A^+\|_{L^\infty} = \|\tfrac{1}{4} [B \times B]\|_{L^\infty} \leq c' \|B\|_{L^2}^2.$$

Assuming a bound on $\|B\|_{L^2}$, we get bounds on $\|F_A^+\|_{L^\infty}$ and $\|B\|_{L^\infty}$.

If $|[B \times B]|^2$ were positive-definite, such bounds would follow automatically from a priori estimates plus maximum principle.

Bounding B

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Thanks to a mean-value inequality due to Morrey

$$\Delta |B| \leq \lambda |B| \implies \|B\|_{L^\infty} \leq c \|B\|_{L^2}$$

Thus

$$\|F_A^+\|_{L^\infty} = \|\tfrac{1}{4} [B \times B]\|_{L^\infty} \leq c' \|B\|_{L^2}^2.$$

Assuming a bound on $\|B\|_{L^2}$, we get bounds on $\|F_A^+\|_{L^\infty}$ and $\|B\|_{L^\infty}$.

If $|[B \times B]|^2$ were positive-definite, such bounds would follow automatically from a priori estimates plus maximum principle.

Bounding B

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Thanks to a mean-value inequality due to Morrey

$$\Delta |B| \leq \lambda |B| \implies \|B\|_{L^\infty} \leq c \|B\|_{L^2}$$

Thus

$$\|F_A^+\|_{L^\infty} = \|\tfrac{1}{4} [B \times B]\|_{L^\infty} \leq c' \|B\|_{L^2}^2.$$

Assuming a bound on $\|B\|_{L^2}$, we get bounds on $\|F_A^+\|_{L^\infty}$ and $\|B\|_{L^\infty}$.

If $|[B \times B]|^2$ were positive-definite, such bounds would follow automatically from a priori estimates plus maximum principle.

Bounding B

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Thanks to a mean-value inequality due to Morrey

$$\Delta |B| \leq \lambda |B| \implies \|B\|_{L^\infty} \leq c \|B\|_{L^2}$$

Thus

$$\|F_A^+\|_{L^\infty} = \|\tfrac{1}{4} [B \times B]\|_{L^\infty} \leq c' \|B\|_{L^2}^2.$$

Assuming a bound on $\|B\|_{L^2}$, we get bounds on $\|F_A^+\|_{L^\infty}$ and $\|B\|_{L^\infty}$.

If $|[B \times B]|^2$ were positive-definite, such bounds would follow automatically from a priori estimates plus maximum principle.

Feehan-Leness program for $\text{PU}(2)$

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Only major property distinguishing $\text{PU}(2)$ monopoles and Vafa-Witten equations is $\| [B \times B] \|^2$ being semi-definite. Their analytic framework extends to give:

- Slice theorem
- Elliptic estimates
- Removal of singularities
- Compactness (almost!)

Compactness requires bounds on $\|F_A^+\|_{L^\infty}$ and $\|B\|_{L^\infty}$.

Feehan-Leness program for $\text{PU}(2)$

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Only major property distinguishing $\text{PU}(2)$ monopoles and Vafa-Witten equations is $\| [B \times B] \|^2$ being semi-definite. Their analytic framework extends to give:

- Slice theorem
- Elliptic estimates
- Removal of singularities
- Compactness (almost!)

Compactness requires bounds on $\|F_A^+\|_{L^\infty}$ and $\|B\|_{L^\infty}$.

Truncated Vafa-Witten moduli space

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\mathcal{M}_{\text{VW},k}^b := \{[0, A, B] \in \mathcal{M}_{\text{VW},k} \mid \|B\|_{L^2} \leq b\}, \quad b \in \mathbb{R}$$

- $\mathcal{M}_{\text{VW},k}^b \subset \mathcal{M}_{\text{VW},k}^{b'}$ for $b \leq b'$.
- $\mathcal{M}_{\text{VW},k}^0 = \mathcal{M}_{\text{ASD},k}$.
- $\mathcal{M}_{\text{VW},k}^b = \emptyset$ for $b < 0$ or $k < -cb^4$

Each $\mathcal{M}_{\text{VW},k}^b$ has an Uhlenbeck compactification $\overline{\mathcal{M}}_{\text{VW},k}^b$.

A partial compactification is given by

$$\overline{\mathcal{M}}_{\text{VW},k} := \bigcup_{b \in \mathbb{R}} \overline{\mathcal{M}}_{\text{VW},k}^b.$$

Truncated Vafa-Witten moduli space

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\mathcal{M}_{\text{VW},k}^b := \{[0, A, B] \in \mathcal{M}_{\text{VW},k} \mid \|B\|_{L^2} \leq b\}, \quad b \in \mathbb{R}$$

- $\mathcal{M}_{\text{VW},k}^b \subset \mathcal{M}_{\text{VW},k}^{b'}$ for $b \leq b'$.
- $\mathcal{M}_{\text{VW},k}^0 = \mathcal{M}_{\text{ASD},k}$.
- $\mathcal{M}_{\text{VW},k}^b = \emptyset$ for $b < 0$ or $k < -cb^4$

Each $\mathcal{M}_{\text{VW},k}^b$ has an Uhlenbeck compactification $\overline{\mathcal{M}}_{\text{VW},k}^b$.

A partial compactification is given by

$$\overline{\mathcal{M}}_{\text{VW},k} := \bigcup_{b \in \mathbb{R}} \overline{\mathcal{M}}_{\text{VW},k}^b.$$

Truncated Vafa-Witten moduli space

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\mathcal{M}_{\text{VW},k}^b := \{[0, A, B] \in \mathcal{M}_{\text{VW},k} \mid \|B\|_{L^2} \leq b\}, \quad b \in \mathbb{R}$$

- $\mathcal{M}_{\text{VW},k}^b \subset \mathcal{M}_{\text{VW},k}^{b'}$ for $b \leq b'$.
- $\mathcal{M}_{\text{VW},k}^0 = \mathcal{M}_{\text{ASD},k}$.
- $\mathcal{M}_{\text{VW},k}^b = \emptyset$ for $b < 0$ or $k < -cb^4$

Each $\mathcal{M}_{\text{VW},k}^b$ has an Uhlenbeck compactification $\overline{\mathcal{M}}_{\text{VW},k}^b$.

A partial compactification is given by

$$\overline{\mathcal{M}}_{\text{VW},k} := \bigcup_{b \in \mathbb{R}} \overline{\mathcal{M}}_{\text{VW},k}^b.$$

Kuranishi complex

Latest version at <http://tensorial.com/math/>

The Kuranishi complex for an instanton $A \in \mathcal{A}_P$ is

$$0 \rightarrow \Omega^0(\mathfrak{g}_P) \xrightarrow{d_A} \Omega^1(\mathfrak{g}_P) \xrightarrow{d_A^+} \Omega^{2,+}(\mathfrak{g}_P) \rightarrow 0$$

with (harmonic) cohomology

$$\mathcal{H}_A^0$$

$$\mathcal{H}_A^1$$

$$\mathcal{H}_A^2$$

- \mathcal{H}_A^0 is the infinitesimal stabilizer (vanishes for irreducibles)
- \mathcal{H}_A^1 is the tangent space of \mathcal{M}_{ASD}
- \mathcal{H}_A^2 is the obstruction to transversality

Kuranishi complex

Latest version at <http://tensorial.com/math/>

The Kuranishi complex for an instanton $A \in \mathcal{A}_P$ is

$$0 \rightarrow \Omega^0(\mathfrak{g}_P) \xrightarrow{d_A} \Omega^1(\mathfrak{g}_P) \xrightarrow{d_A^+} \Omega^{2,+}(\mathfrak{g}_P) \rightarrow 0$$

with (harmonic) cohomology

$$\mathcal{H}_A^0$$

$$\mathcal{H}_A^1$$

$$\mathcal{H}_A^2$$

- \mathcal{H}_A^0 is the infinitesimal stabilizer (vanishes for irreducibles)
- \mathcal{H}_A^1 is the tangent space of \mathcal{M}_{ASD}
- \mathcal{H}_A^2 is the obstruction to transversality

Index theorem

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

$$0 \rightarrow \underbrace{\Omega^0(\mathfrak{g}_P)}_{\mathcal{H}_A^0} \xrightarrow{d_A} \underbrace{\Omega^1(\mathfrak{g}_P)}_{\mathcal{H}_A^1} \xrightarrow{d_A^+} \underbrace{\Omega^{2,+}(\mathfrak{g}_P)}_{\mathcal{H}_A^2} \rightarrow 0$$

We will always assume A is irreducible, so $\mathcal{H}_A^0 = 0$.

By the index theorem,

$$\dim \mathcal{H}_A^1 = d + c \qquad \dim \mathcal{H}_A^2 = c$$

where d is the expected dimension $8k - 3(1 - b_1 + b^+)$ for $G = \mathrm{SU}(2)$.

For generic metrics, $c = 0$, and a neighborhood of $[A] \in \mathcal{M}_{\mathrm{ASD}}$ is modeled by \mathcal{H}_A^1 with dimension d .

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Index theorem

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$0 \rightarrow \Omega^0(\mathfrak{g}_P) \xrightarrow{d_A} \Omega^1(\mathfrak{g}_P) \xrightarrow{d_A^+} \Omega^{2,+}(\mathfrak{g}_P) \rightarrow 0$$
$$\mathcal{H}_A^0 \qquad \mathcal{H}_A^1 \qquad \mathcal{H}_A^2$$

We will always assume A is irreducible, so $\mathcal{H}_A^0 = 0$.

By the index theorem,

$$\dim \mathcal{H}_A^1 = d + c \qquad \dim \mathcal{H}_A^2 = c$$

where d is the expected dimension $8k - 3(1 - b_1 + b^+)$ for $G = \mathrm{SU}(2)$.

For generic metrics, $c = 0$, and a neighborhood of $[A] \in \mathcal{M}_{\mathrm{ASD}}$ is modeled by \mathcal{H}_A^1 with dimension d .

Inverse function theorem

Latest version at <http://tensorial.com/math/>

Consider

$$\dim \mathcal{H}_A^1 = d + 1 \quad \dim \mathcal{H}_A^2 = 1$$

The differential

$$\Omega^1(\mathfrak{g}_P) \xrightarrow{d_A^+} \Omega^{2,+}(\mathfrak{g}_P)$$

is no longer surjective. For $a \in H_A^1$,

$$F_{A+a}^+ = F_A^+ + d_A^+ a + \frac{1}{2} [a \wedge a]^+ = \frac{1}{2} [a \wedge a]^+.$$

Can use inverse function theorem to find \tilde{a} with $F_{A+\tilde{a}}^+ = 0$ when $\frac{1}{2} [a \wedge a]^+ \perp \mathcal{H}_A^2$.

Quadratic model

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\dim \mathcal{H}_A^1 = d + 1 \quad \dim \mathcal{H}_A^2 = 1$$

Fix $\omega \in \mathcal{H}_A^2$, with $\omega \neq 0$. Define a quadratic form $q(a)$ on \mathcal{H}_A^1 by

$$q(a) := \int_X \left\langle \frac{1}{2} [a \wedge a]^+ \cdot \omega \right\rangle = 0.$$

If q is nondegenerate, a neighborhood of $[A] \in \mathcal{M}_{\text{ASD}}$ is modeled on $q(a) = 0$.

Vafa-Witten quadratic model

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

$$\int_X \langle [a \times b] \cdot \alpha_1 \rangle = 0,$$

$$\vdots$$

$$\int_X \langle [a \times b] \cdot \alpha_{d+1} \rangle = 0,$$

$$\int_X \left\langle \frac{1}{2} [a \wedge a]^+ - \frac{1}{4} [b \times b] \cdot \omega \right\rangle = 0.$$

When nondegenerate, first equations say $a = 0$ or $b = 0$.

Perturbing the metric

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten


Properness

Singular
instantons and
gluing

Consider a perturbation of conformal structure $m \in \Omega^0(X; \text{Hom}(\Lambda^{2,-} V^*, \Lambda^{2,+} V^*))$, and the parameterized family tm , $t \in \mathbb{R}$. We get an extra term

$$\int_X \left\langle \left(\frac{1}{2} [a \wedge a]^+ - \frac{1}{4} [b \times b] - tm F_A^- \right) \cdot \omega \right\rangle = 0.$$

When nondegenerate, we get the picture



Gluing theory and future work

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Following the work of Donaldson and Taubes, we can construct approximate solutions by grafting concentrated instantons.

The obstruction to repairing the graft to obtain a genuine solution can be approximated by a quadratic map.

Assuming nondegeneracy, we obtain quadratic models of the Uhlenbeck ends of the moduli space.

We hope to gain insight into the lingering compactness issues by studying these models.

New nonlinear estimates will hopefully lead to better a priori bounds.

Gluing theory and future work

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Following the work of Donaldson and Taubes, we can construct approximate solutions by grafting concentrated instantons.

The obstruction to repairing the graft to obtain a genuine solution can be approximated by a quadratic map.

Assuming nondegeneracy, we obtain quadratic models of the Uhlenbeck ends of the moduli space.

We hope to gain insight into the lingering compactness issues by studying these models.

New nonlinear estimates will hopefully lead to better a priori bounds.

Gluing theory and future work

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Following the work of Donaldson and Taubes, we can construct approximate solutions by grafting concentrated instantons.

The obstruction to repairing the graft to obtain a genuine solution can be approximated by a quadratic map.

Assuming nondegeneracy, we obtain quadratic models of the Uhlenbeck ends of the moduli space.

We hope to gain insight into the lingering compactness issues by studying these models.

New nonlinear estimates will hopefully lead to better a priori bounds.

Gluing theory and future work

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Following the work of Donaldson and Taubes, we can construct approximate solutions by grafting concentrated instantons.

The obstruction to repairing the graft to obtain a genuine solution can be approximated by a quadratic map.

Assuming nondegeneracy, we obtain quadratic models of the Uhlenbeck ends of the moduli space.

We hope to gain insight into the lingering compactness issues by studying these models.

New nonlinear estimates will hopefully lead to better a priori bounds.

Gluing theory and future work

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

Following the work of Donaldson and Taubes, we can construct approximate solutions by grafting concentrated instantons.

The obstruction to repairing the graft to obtain a genuine solution can be approximated by a quadratic map.

Assuming nondegeneracy, we obtain quadratic models of the Uhlenbeck ends of the moduli space.

We hope to gain insight into the lingering compactness issues by studying these models.

New nonlinear estimates will hopefully lead to better a priori bounds.

IT'S OVER!

Latest version at <http://tensorial.com/math/>

Some Analytic
Aspects of
Vafa-Witten
Twisted $\mathcal{N} = 4$
Supersymmetric
Yang-Mills Theory

Ben Mares

An introduction
to anti-self-duality

History and
motivation

The equations

Energy identities
in gauge theory

Energy identities
for Vafa-Witten

Properness

Singular
instantons and
gluing

- Thanks for listening!