

# Seiberg-Witten invariants and surface singularities

Liviu I. Nicolaescu

This is a joint project with Andras Nemethi. For details one can consult [math.AG/0111298](#), [math.AG/0201120](#).

## Contents

<b>1</b>	<b>Topological Invariants</b>	<b>1</b>
<b>2</b>	<b>Analytic invariants</b>	<b>4</b>
<b>3</b>	<b>The Main Problem and a Bit of History</b>	<b>5</b>
<b>4</b>	<b>The Main Conjecture and Evidence in its Favor</b>	<b>6</b>

$(X, p)$  (germ) of isolated surface singularity (*i.s.s.* for brevity). Assume  $X$  is Stein.

## 1 Topological Invariants

**The link.** Embed  $(X, p) \hookrightarrow (\mathbb{C}^N, 0)$ , and set

$$M = X \cap S_\varepsilon^{2N-1}(0).$$

$M$  is an oriented 3-manifold independent on the embedding and  $\varepsilon \ll 1$ .




Figure 1: *The link of an isolated singularity*

**Good resolutions.** A resolution of  $(X, p)$  is a pair  $(\tilde{X}, \pi)$  where

- $\tilde{X}$  is a smooth complex surface;
- $\tilde{X} \xrightarrow{\pi} X$  is holomorphic;
- $\tilde{X} \setminus \pi^{-1}(p) \rightarrow X \setminus p$  is biholomorphic;

The resolution is called *good* if the exceptional divisor  $E := \pi^{-1}(p)$  is a normal crossing divisor i.e its irreducible components  $(E_i)_{1 \leq i \leq n}$  are smooth curves intersecting transversally.

**FACT.** *Good resolutions exist but are not unique. There exists a unique minimal resolution  $\tilde{X}$ , i.e. a resolution containing no  $-1$ -spheres. There exists a unique minimal good resolution. (It may have  $-1$  spheres, but when blown down the exceptional divisor will no longer be a normal crossing divisor). Any other resolution is obtained from the minimal one by blowing-up/down  $-1$  spheres.*

Suppose  $\tilde{X}$  is a resolution of  $X$ . We set

$$\Lambda = \Lambda(\tilde{X}) := \text{span}_{\mathbb{Z}}\{E_i\} \subset H_2(\tilde{X}, \mathbb{Z}),$$

$$\Lambda_+(\tilde{X}) := \left\{ \sum_i m_i E_i \in \Lambda; m_i \geq 0 \right\}.$$

**Theorem.** (D. Mumford) *The symmetric matrix  $(E_i \cdot E_j)_{i,j}$  is  $< 0$ .*

**The dual resolution graph.** Suppose  $(\tilde{X}, \pi)$  is a good resolution of the i.s.s.  $(X, p)$  with exceptional divisor  $E = \bigcup_i E_i$ . The (dual) resolution graph is a decorated graph  $\Gamma = \Gamma_{\tilde{X}}$  obtained as follows.

- There is one vertex  $v_i$  for each component  $E_i$ .
- Two vertices  $v_i, v_j, i \neq j$  are connected by  $E_i \cdot E_j$  edges.
- Each vertex  $v_i$  is decorated by two integers, the genus  $g_i$  of  $E_i$ , and the self intersection number  $e_i := E_i^2$ .

We see that  $\tilde{X}$  is a plumbing of disk bundles over the Riemann surfaces  $E_i$ , with plumbing instructions contained in the graph  $\Gamma$ . The boundary of this plumbing is precisely the link of the singularity.




Figure 2: A plumbing and its associated dual graph

**Theorem.** (W. Neumann) *Suppose  $(X_i, p), i = 0, 1$  are two i.s.s. Denote by  $M_i$  their links, and by  $\tilde{X}_i$  their minimal good resolutions. The following statements are equivalent.*

(a) The graphs  $\Gamma_{\tilde{X}_i}$  are isomorphic (as weighted graphs).

(b) The links  $M_i$  are diffeomorphic as oriented 3-manifolds.

**Definition.** We say that a property of an i.s.s. is *topological* if it can be described in terms of the combinatorics of the dual graph of the minimal good resolution. □

**The arithmetic genus.**  $\tilde{X}$  resolution of  $(X, p)$ ,  $E = \cup_i E_i$ , the exceptional divisor. Note that every  $Z = \sum_i n_i E_i \in \Lambda_+$  can be identified with a compact complex curve on  $\tilde{X}$ . The *arithmetic genus* of  $Z$  is defined by

$$p_a(Z) = 1 + \frac{1}{2}(Z \cdot Z + \langle K_{\tilde{X}}, Z \rangle),$$

where  $K_{\tilde{X}} \in H^2(\tilde{X}, \mathbb{Z})$  is the canonical line bundle of  $\tilde{X}$ . When  $Z$  is a smooth curve  $p_a(Z)$  is the usual genus of  $Z$ . Set

$$p_a(\tilde{X}) := \sup\{p_a(Z); Z \in \Lambda_+ \setminus \{0\}\}.$$

This nonnegative integer is independent of the resolution and thus it is a *topological* invariant of  $(X, p)$ . We will denote it by  $p_a(X, p)$ , and we will refer to it as the arithmetic genus of the singularity.

**The canonical cycle.**  $(X, p)$  - i.s.s. and  $(\tilde{X}, \pi)$  is a resolution. The canonical cycle is the cycle  $Z_K = Z_K(\tilde{X}) \in \Lambda \otimes \mathbb{Q}$  defined by

$$Z_K \cdot E_j = -\langle K_{\tilde{X}}, E_j \rangle = 2 - p_a(E_j) + E_j^2, \forall i.$$

Set

$$\gamma(\tilde{X}) = Z_{K_{\tilde{X}}}^2 + b_2(\tilde{X}) \in \mathbb{Q}.$$

This number is independent of the resolution  $\tilde{X}$ , and thus it is a topological invariant of  $(X, p)$ . We will denote it by  $\gamma(X, p)$ . Note that if  $\tilde{X}$  is the minimal good resolution then  $Z_{K_{\tilde{X}}}$  is a topological invariant of  $M$ .

Observe that

$$\gamma(X, p) = \left( K_{\tilde{X}}^2 - (2\chi(\tilde{X}) + 3\text{sign}(\tilde{X})) \right) + 2 - 2b_1(\tilde{X}). \quad (\gamma)$$

**Definition.** Suppose  $(X, p)$  is an i.s.s., and  $(\tilde{X}, \pi, E)$  is a good resolution. The singularity is called **Gorenstein** if  $K_{\tilde{X}}|_{\tilde{X} \setminus E}$  is **holomorphically trivial**. The singularity is called **numerically Gorenstein** if  $K_{\tilde{X}}|_{\tilde{X} \setminus E}$  is **topologically trivial**.

Observe that  $(X, p)$  is numerically Gorenstein iff

$$K_{\tilde{X}} \in H^2(\tilde{X}, \partial\tilde{X}; \mathbb{Z}) \iff Z_K \in \Lambda.$$

**Example.** All local complete intersection singularities are Gorenstein. Recall that the i.s.s.  $(X, p)$  is a local complete intersection singularity if near  $p$  it can be described as the zero set of a holomorphic map  $F: \mathbb{C}^N \rightarrow \mathbb{C}^{N-2}$ . □

## 2 Analytic invariants

**The geometric genus.**  $(X, p)$  i.s.s.,  $X$  Stein,  $\tilde{X}$  resolution.

$$\tilde{X} \text{ Levi pseudoconvex} \implies \dim H^k(\tilde{X}, \mathcal{O}_{\tilde{X}}) < \infty, \quad \forall k \geq 1.$$

The integer  $\dim H^1(\tilde{X}, \mathcal{O}_{\tilde{X}})$  is independent of the resolution, and thus it is an *analytic* invariant of  $(X, p)$ . It is called the *geometric genus* and is denoted by  $p_g(X, p)$ . It is known that

$$p_g(X, p) \geq p_a(X, p).$$

**Example.** Suppose  $L \rightarrow \Sigma$  is a degree  $d < 0$  holomorphic line bundle over the Riemann surface  $\Sigma$  of genus  $g$ . By a theorem of Grauert there exists a natural Stein space  $X$  with an isolated singularity at  $p \in X$ , and a holomorphic map

$$(L, \Sigma) \xrightarrow{\pi} (X, p)$$

which makes  $L$  a good resolution of  $(X, p)$  with exceptional divisor  $\Sigma \hookrightarrow L$ . Then

$$p_a(X, p) = g, \quad Z_K = \left(1 + \frac{2-2g}{d}\right)\Sigma, \quad \gamma(X, p) = d\left(1 + \frac{2-2g}{d}\right)^2 + 1,$$

$$p_g(X, p) = \sum_{n \geq 0} \dim H^1(\Sigma, \mathcal{O}(-nL)) \stackrel{Serre}{=} \sum_{n \geq 0} \dim H^0(\Sigma, \mathcal{O}(K_\Sigma + nL)). \quad (p_g)$$

We deduce that  $p_g(X, p)$  depends on the complex structure on  $\Sigma$ , and on the complex structure on  $L$ , i.e. on the holomorphic embedding  $\Sigma \hookrightarrow L$ . These dependencies on analytic data become irrelevant under appropriate topological constraints.

- $\Sigma$  is rational, i.e.  $g = 0$ .
- $\Sigma$  is elliptic, i.e.  $g = 1$ .
- The degree of  $L$  is sufficiently negative,  $\deg L \leq -g$ .

In all these cases  $p_g(X, p) = p_a(X, p) = g$ .

□

**Smoothings.** A smoothing of an i.s.s.  $(X, p)$  is a proper flat map  $(\mathcal{X}, q) \xrightarrow{F} (\mathbb{C}, 0)$  together with an embedding  $\iota : (X, p) \hookrightarrow (\mathcal{X}, q)$  which induces an isomorphism  $(X, p) \cong (F^{-1}(0), q)$ . For  $t \in \mathbb{C}^*$  sufficiently small the fiber  $X_t := f^{-1}(t)$  is smooth. Its topology is independent of  $t$ .  $X_t$  is called the *Milnor fiber* of the smoothing. The Milnor number  $\mu$  of the smoothing is  $b_2(X_t)$ .

**Example.** Suppose  $(X, p)$  is the complete intersection, described by the zero set of a map  $F : \mathbb{C}^N \rightarrow \mathbb{C}^{N-2}$ . To construct smoothings of  $(X, p)$  it suffices to pick a line through the origin  $L \subset \mathbb{C}^{N-2}$  and set  $\mathcal{X} := F^{-1}(L)$ .

□

**Theorem.** (Durfee-Laufer-Steenbrink-Wahl) *Suppose  $(X, p)$  is a smoothable Gorenstein i.s.s. We denote by  $F$  its Milnor fiber. Then*

$$p_g(X, p) = -\frac{1}{8} \text{sign}(F) - \frac{1}{8} \gamma(X, p).$$

Motivated by the above result, we define the *virtual signature* of an i.s.s. by

$$\sigma_{virt}(X, p) := -8p_g(X, p) - \gamma(X, p).$$

For smoothable Gorenstein singularities, the virtual signature is the signature of the Milnor fiber.

### 3 The Main Problem and a Bit of History

**The Main Problem.** *How much information about the analytic structure of the i.s.s.  $(X, p)$  is encoded in the topology of its link  $M$ . In particular, can we determine  $p_g(X, p)$  from combinatorial data contained in the dual resolution graph of the minimal good resolution?*

**History.** Work of the past four decades indicates that the link often contains nontrivial information about the analytic structure.

❶ (D. Mumford, 1961)  $(X, p)$  is smooth at  $p$  if and only if the link is  $\cong S^3$ .

❷ (M. Artin, 1962-66)  $p_g(X, p) = 0 \iff p_a(X, p) = 0$ . In this case, the link  $M$  is a rational homology sphere ( $\mathbb{Q}HS$  for brevity).

❸ (H. Laufer, 1977) Assume that  $(X, p)$  is elliptic, i.e.  $p_a(X, p) = 1$ , and Gorenstein. Then the condition  $p_g(X, p) = 1$  is topological.

❹ (A. Nemethi, 1999) Assume that  $(X, p)$  is elliptic, Gorenstein, and the link  $M$  is a  $\mathbb{Q}HS$ . Then  $p_g(X, p)$  is equal to a certain topological invariant of  $M$ , the length of the elliptic sequence defined by S.S.-T. Yau.

**Remark.** (a)  $M$  is a  $\mathbb{Q}HS$  iff  $\Gamma$  is a tree and all the components  $E_i$  are rational curves.

(b) The condition that  $M$  is a  $\mathbb{Q}HS$  cannot be removed from ❹. To see this consider the singularities  $(X_1, 0) = \{x^2 + y^3 + z^{18} = 0\}$ ,  $(X_2, 0) = \{z^2 + y(x^4 + y^6) = 0\}$ . They have isomorphic resolution graphs (see below), but  $p_g(X_1, 0) = 3$ ,  $p_g(X_2, 0) = 2$ .  $\square$



❺ (Fintushel-Stern, Neumann-Wahl, 1990) Suppose  $(X, p)$  is a Brieskorn complete intersection singularity and the link  $M$  is an *integral* homology sphere ( $\mathbb{Z}HS$ ). Then

$$\text{Casson}(M) = -\frac{1}{8}\sigma_{virt}(X, p).$$

Here we recall that a Brieskorn complete intersection singularity is a complete intersection singularity of the form

$$\begin{cases} a_{11}z_1^{p_1} + \cdots + a_{1n}z_n^{p_n} = 0 \\ \vdots \\ a_{(n-2)1}z_1^{p_1} + \cdots + a_{(n-2)n}z_n^{p_n} = 0 \end{cases}$$

**Remark.** If in ❺ we assume only that  $M$  is a  $\mathbb{Q}HS$  then the obvious generalization

$$\text{Casson-Walker}(M) = -\frac{1}{8}\sigma_{virt}(X, p).$$

is no longer true. □

#### 4 The Main Conjecture and Evidence in its Favor

**The Main Conjecture.** Suppose  $(X, p)$  is a rational, or Gorenstein singularity such that its link is a  $\mathbb{Q}HS$ . Then  $M$  is equipped with a canonical  $spin^c$  structure  $\sigma_{can}$ , which depends only on the resolution graph  $\Gamma$  of the minimal good resolution, and

$$sw_M(\sigma_{can}) = -\frac{1}{8}\sigma_{virt}(X, p),$$

where  $sw_M(\sigma_{can})$  denotes the Seiberg-Witten invariant of the canonical  $spin^c$  structure. In particular, if  $M$  is a  $\mathbb{Z}HS$  then there is a unique  $spin^c$  structure on  $M$  whose Seiberg-Witten invariant equals the Casson invariant of  $M$  so that

$$\text{Casson}(M) = -\frac{1}{8}\sigma_{virt}(X, p).$$

□

**Evidence.** We need to describe the various terms in the Main Conjecture.

Denote by  $\tilde{X}$  the minimal good resolution of  $(X, p)$ . Then

$$\Lambda = H_2(\tilde{X}, \mathbb{Z}), \quad H^2(\tilde{X}, \mathbb{Z}) \cong \check{\Lambda} := \text{Hom}(\Lambda, \mathbb{Z})$$

Set  $H := H_1(M, \mathbb{Z})$ , and denote the group operation on  $H$  multiplicatively. The intersection form on  $\Lambda$  defines an embedding  $\Lambda \hookrightarrow \check{\Lambda}$ , and we have

$$H \cong \check{\Lambda}/\Lambda.$$

$H$  acts freely and transitively on the set  $Spin^c(M)$  of  $spin^c$  structures on  $M$

$$H \times Spin^c(M) \ni (h, \sigma) \mapsto h \cdot \sigma \in Spin^c(M).$$

To define the canonical  $spin^c$  structure  $\sigma_{can}$  let us recall that a choice of a  $spin^c$  structure on  $M$  is equivalent to a choice of an almost complex structure on the stable tangent bundle  $\mathbb{R} \oplus TM$  of  $M$ . The stable tangent bundle of  $M$  is equipped with a natural complex structure induced by the complex structure on  $\tilde{X}$ .  $\sigma_{can}$  is the  $spin^c$  structure associated to this complex structure.

\* **Proposition.**  $\sigma_{can}$  can be described only in terms of the combinatorics of  $\Gamma_{\tilde{X}}$ .

*Proof.* Denote by  $\mathbf{lk}_M : H \times H \rightarrow \mathbb{Q}/\mathbb{Z}$  the linking form of  $M$ . An enhancement of  $\mathbf{lk}_M$  is a function

$$q : H \rightarrow \mathbb{Q}/\mathbb{Z}$$

such that

$$q(h_1 h_2) - q(h_1) - q(h_2) = \mathbf{lk}_M(h_1, h_2), \quad \forall h_1, h_2 \in H.$$

There is a natural bijection between  $Spin^c(M)$  and the set of enhancements,  $\sigma \mapsto q_\sigma$ . Recalling that  $H \cong \check{\Lambda}/\Lambda$  we define

$$q_{can} : \check{\Lambda}/\Lambda \rightarrow \mathbb{Q}, \quad q_{can}(h) = -\frac{1}{2}(K_{\check{X}} \cdot \check{h} + \check{h} \cdot \check{h}) \bmod \mathbb{Z}$$

for every  $h \in \check{\Lambda}/\Lambda$ , and every  $\check{h} \in \check{\Lambda}$  which projects onto  $h$ . The expression in the right hand side depends only on  $\Gamma$ .  $\sigma_{can}$  is the  $spin^c$  structure corresponding to  $q_{can}$ .  $\square$

**Remark.** (a)  $q_{can}$  first appeared in work of Looijenga-Wahl.

(b) From the above description of  $\sigma_{can}$  and the equality  $(\gamma)$  we deduce that  $\gamma(X, p) - 2$  equals the Gompf invariant of the  $spin^c$  structure  $\sigma_{can}$ .  $\square$

The Seiberg-Witten invariant is a function

$$sw_M : Spin^c(M) \rightarrow \mathbb{Q}, \quad \sigma \mapsto sw_M(\sigma).$$

$sw_M(\sigma) = \#$  of Seiberg-Witten  $\sigma$ -monopoles + the Kreck-Stolz invariant of  $\sigma$  (a certain combination of eta invariants). For each  $\sigma \in Spin^c(M)$  define

$$SW_{M,\sigma} : H \rightarrow \mathbb{Q}, \quad SW_{M,\sigma}(h) = sw_M(h^{-1} \cdot \sigma).$$

One can give a combinatorial description of this invariant. For each  $spin^c$  structure  $\sigma$ , the Reidemeister-Turaev torsion of  $(M, \sigma)$  is a function

$$\mathcal{T}_{M,\sigma} : H \rightarrow \mathbb{Q}.$$

Denote by  $CW_M$  the Casson-Walker invariant of  $M$ , and define the modified Reidemeister-Turaev torsion of  $M$  by

$$\mathcal{T}_{M,\sigma}^0 : H \rightarrow \mathbb{Q}, \quad \mathcal{T}_{M,\sigma}^0(h) := \frac{1}{|H|} CW_M + \mathcal{T}_{M,\sigma}(h), \quad \forall h \in H.$$

**Theorem.** (L.I. Nicolaescu) For every  $\sigma \in Spin^c(M)$  we have

$$SW_{M,\sigma} \equiv \mathcal{T}_{M,\sigma}^0.$$

Denote by  $\hat{H}$  the Pontryagin dual of  $H$ ,  $\hat{H} := \text{Hom}(H, U(1))$ . The Fourier transform of  $\mathcal{T}_{M,\sigma_{can}}$  is the function

$$\hat{\mathcal{T}}_{M,\sigma_{can}} : \hat{H} \rightarrow \mathbb{C}, \quad F(\chi) = \sum_{h \in H} \mathcal{T}_{M,\sigma_{can}}(h) \bar{\chi}(h).$$

The Fourier inversion formula implies

$$sw_M(\sigma_{can}) = SW_{M,\sigma_{can}}(1) = \frac{1}{|H|} CW_M + \frac{1}{|H|} \sum_{\chi \in \hat{H}} \hat{\mathcal{T}}_{M,\sigma_{can}}(\chi). \quad (*)$$

**Theorem.**(Lescop-Rațiu) *The Casson-Walker invariant can be described **explicitly** in terms of the combinatorics of  $\Gamma$ .*

**Main Technical Result.** (Nemethi-Nicolaescu)  $\hat{\mathcal{J}}_{M,\sigma_{can}}$  can be described **explicitly** in terms of the combinatorics of  $\Gamma_{\tilde{X}}$ .

*Idea of Proof.* Using surgery formulæ we produce an *explicit* holomorphic regularization  $R_M$  of  $\hat{\mathcal{J}}_{M,\sigma_{can}}$ . This is an element in the group algebra  $\mathbb{C}(t)[\hat{H}]$ ,

$$R_M = \sum_{\chi \in \hat{H}} R_\chi(t)\chi, \quad R_\chi \in \mathbf{C}(t) = \text{the field of rational functions in one variable} \quad (**)$$

such that for every character  $\chi$

$$\lim_{t \rightarrow 1} R_\chi(t) = \hat{\mathcal{J}}_{M,\sigma_{can}}(\chi).$$

□

In applications the sum in the right-hand side of (\*) is difficult to compute if the combinatorics of the graph is very involved.

**Theorem.** (Nemethi-Nicolaescu) *The Main Conjecture is true for **all** the quasihomogeneous singularities whose links are  $\mathbb{Q}HS$ 's.*

*Idea of Proof.* For a quasihomogeneous singularity  $(X, p)$  the resolution graph is star-shaped and the sum in (\*) simplifies somewhat. Our expression for the holomorphic regularization  $R_M$  in (\*\*) is formally identical to the Poincaré series associated to the Universal Abelian cover of  $(X, p)$  introduced by W. Neumann. The proof of the Main Conjecture in this case relies on a formula for  $p_g(X, p)$  of Dolgachev-Pinkham, and on some ideas of Neumann and Zagier.

□