

Locally compact quantum groups. A von Neumann algebra approach.

A. Van Daele (*)

Abstract

In this paper, we give an alternative approach to the theory of locally compact quantum groups, as developed by Kustermans and Vaes in [K-V1] and [K-V2]. We start with a von Neumann algebra and a comultiplication on this von Neumann algebra. We assume that there exist faithful left and right Haar weights. Then we develop the theory within this von Neumann algebra setting.

In [K-V3] locally compact quantum groups are also studied in the von Neumann algebraic context. This approach is independent of the original C^* -algebraic approach in the sense that the earlier results are not used. However, the paper [K-V3] is not really independent because for many proofs, the reader is referred to the original paper where the C^* -version is developed.

In this paper, we give a self-contained approach. Moreover, at various points, we do things differently. We have a different treatment of the antipode. It is similar to the original treatment in [K-V2]. But together with the fact that we work in the von Neumann algebra framework, it allows us to use an idea from [S-V-Z2] to obtain the uniqueness of the Haar weights in an early stage. We take advantage of this fact when deriving the other main results in the theory. We also give a slightly different approach to duality. Finally, we collect, in a systematic way, several important formulas.

In an appendix, we indicate very briefly how the C^* -approach and the von Neumann algebra approach eventually yield the same objects. The passage from the von Neumann algebra setting to the C^* -algebra setting is more or less standard. For the other direction, we use a new method. It is based on the observation that the Haar weights on the C^* -algebra extend to weights on the double dual with central support and that all these supports are the same. Of course, we get the von Neumann algebra by cutting down the double dual with this unique support projection in the centre.

All together, we see that there are many advantages when we develop the theory of locally compact quantum groups in the von Neumann algebra framework, rather than in the C^* -algebra framework. It is not only simpler, the theory of weights on von Neumann algebras is better known and one needs very little to go from the C^* -algebras to the von Neumann algebras. Moreover, in many cases when constructing examples, the von Neumann algebra with the coproduct is constructed from the very beginning and the Haar weights are constructed as weights on this von Neumann algebra (using left Hilbert algebra theory).

This paper is written in a concise way. In many cases, only indications for the proofs of the results are given. This information should be enough for specialists to see that these results are correct. We will give more details in [VD8], a paper which will be expository, aimed at non-specialists. See also [VD6] for an 'expanded' version of the appendix.

February 2006 (Version 1.0)

(*) Department of Mathematics, K.U. Leuven, Celestijnenlaan 200B, B-3001 Heverlee (Belgium). E-mail: Alfons.VanDaele@wis.kuleuven.be

0. Introduction

Let M be a von Neumann algebra and Δ a comultiplication on M (see Definition 1.1 in Section 1 for a precise definition). The pair (M, Δ) is called a *locally compact quantum group* (in the von Neumann algebraic sense) if there exist faithful left and right Haar weights (see Definition 2.1 in Section 2). This definition is due to Kustermans and Vaes (see [K-V3]).

In their fundamental papers [K-V1] and [K-V2], Kustermans and Vaes develop the theory of locally compact quantum groups in the C^* -algebraic framework and in [K-V3], they show that both the original C^* -algebra approach and the von Neumann algebra approach give the same objects. There is indeed a standard procedure to go from a locally compact quantum group in the C^* -algebra setting to a locally compact quantum group in the von Neumann algebra sense (and vice versa).

In this paper, we present an alternative approach to the theory of locally compact quantum groups. The basic difference is that we develop the main theory in the framework of von Neumann algebras (and not in the C^* -algebraic setting as was done in [K-V1] and [K-V2]). It is well-known that the von Neumann algebra setting is, in general, simpler to work in. To begin with, the definition of a locally compact quantum group in the von Neumann algebra framework is already less complicated than in the C^* -algebra setting. Also the theory of weights on von Neumann algebras is better known than the theory of weights on C^* -algebras. One has to be a bit more careful with using the various topologies, but on the other hand, one does not have to worry about multiplier algebras.

We also describe a (relatively) quick way to go from a locally compact quantum group in the C^* -algebraic sense to a locally compact quantum group in the von Neumann algebraic sense. We do not need to develop the C^* -theory to do this. In some sense, this justifies our choice. Remark that the other direction, from von Neumann algebras to C^* -algebras, is the easier one (and standard).

However, the difference of this work with the other (earlier) approaches not only lies in the fact that we work in the von Neumann algebra framework. We also have a slightly different approach to construct the antipode (see Section 1). We do not use operator space techniques (as e.g. in [V-VD]). This in combination with the use of Connes' cocycle Radon Nikodym theorem (an idea that we found in earlier works by Stratila, Voiculescu and Zsido (see [S-V-Z1], [S-V-Z2] and [S-V-Z3])) allows us to obtain uniqueness of the Haar weights in an earlier stage of the development. This in turn will yield other simplifications.

Finally, our approach in this paper is also self-contained. In their paper on the von Neumann algebraic approach [K-V3], Kustermans and Vaes do not really use results from the earlier paper on the C^* -algebra approach [K-V2], but nevertheless, it is hard to read it without the first paper because of the fundamental references to this first paper. In fact, we also rely less on results from other papers (e.g. on weights on C^* -algebras or about manageability of multiplicative unitaries) as is done in the original works. Also for the proofs that are omitted, this is the case. Moreover, where possible, we avoid working with unbounded operators and weights (more than in the original papers) but we try to use bounded operators and normal linear functionals.

The *content of the paper* is as follows.

In *Section 1* we work with a von Neumann algebra and a comultiplication. We consider the antipode S , together with an involutive operator K on the Hilbert space that implements the antipode in the sense that, roughly speaking, $S(x)^* = KxK$ when $x \in \mathcal{D}(S)$. The right Haar weight is needed to construct this operator K and the left Haar weight is used to prove that it is densely defined. This last property is closely related with the right regular representation being unitary. Also in this section, we focus on various other densities. It makes this section longer than the others, but the reason for doing so is that these density results are closely related with the construction of the antipode and the operator K . Finally, in this section, we modify the definition of the antipode so that it becomes more tractable. We discuss its polar decomposition (with the scaling group (τ_t) and the unitary antipode R) and we prove the basic formulas about this modified antipode, needed further in the paper.

Our approach here is not so very different from the way this is done by Kustermans and Vaes in the sense that we use the same ideas. Among other things, we also use Kustermans' trick to prove that the right regular representation is unitary. On the other hand, we try to avoid using weights when possible.

In *Section 2* we give the main results. One of these results is the uniqueness of the Haar weights. The formulas involving the scaling group (τ_t) and the unitary antipode R , proven in Section 1, together with Connes' cocycle Radon Nikodym theorem, are used to show the uniqueness. Here, our approach is quite different from the original one in [K-V1] and [K-V2] and uses an idea found in [S-V-Z2]. From the uniqueness, and again using basic formulas involving the scaling group and the unitary antipode from Section 1, the main results are relatively easy to prove.

In *Section 3* we treat the dual. This is more or less standard. Our approach is again slightly different in the way we use the results obtained in Section 1 and in Section 2. Also, since we are basically only considering the von Neumann algebra version, the construction of the dual is somewhat simpler.

In *Section 4* we collect a set of formulas. The main ingredients are the various objects (the left and right Haar weights with their modular structures, the left and right regular representations, the antipode with the scaling group and the unitary antipode, the operators on the Hilbert space implementing these automorphism groups, ...), for the original pair (M, Δ) , as well as for the dual pair $(\hat{M}, \hat{\Delta})$. In fact, this section and these formulas can well be used as a fairly complete chapter needed to work with locally compact quantum groups.

We have chosen to discuss the procedure to pass from the C^* -algebraic locally compact quantum groups to the von Neumann algebraic ones in the *Appendix*. We do this because it is not really needed for the development of the theory as it is done in this paper. Here again, our approach is rather different from the original one. The main idea is to pass first to the double dual A^{**} of the C^* -algebra A . Then, it is quickly proven that the supports of the invariant weights (of the type used in this theory), are all the same central projection in the double dual. Cutting down the double dual by this central projection gives us the von Neumann algebra M . The coproduct, as well as the Haar weights, are obtained by

first extending the corresponding original objects to the double dual and then restricting them again to this von Neumann algebra M . The converse is more or less standard and of course, makes use of the results in the paper. A more or less independent treatment of the connection of the two approaches is found in an expanded version of this appendix, see [VD6].

We also use this appendix to say something more about the relation of our approach with the one by Masuda, Nakagami and Woronowicz (in [M-N] and [M-N-W]).

In this paper, we will not give full details. We give precise definitions and statements, but often we will only *sketch proofs*. We give sufficiently many details so that the reader should be 'convinced' about the result. In fact, any reader, familiar with the material, should be able to complete the proofs without too much effort. On the other hand, for the less experienced reader, we refer to a forthcoming paper *Notes on locally compact quantum groups* [VD8]. These notes are intended as lecture notes on the subject, for (young) researchers who want to learn about locally compact quantum groups. So, full details of the proofs of the results in this paper can be found there. This style of writing allows us to keep this paper relatively short, while on the other hand, we still are able to make it, to a great extent, self-contained. In the appendix, we give even less details because this is not so essential for the development here. In [VD6] is found an 'expanded' version of this appendix, but again, for more details, we refer to [VD8]. There is also the paper on weights on C^* -algebras [Q-VD] where more details can be found about weight theory on C^* -algebras as is necessary for understanding the locally compact quantum groups in the C^* -algebraic sense. Finally, let us also refer to a forthcoming set of notes on the theory of multiplier Hopf $(*)$ -algebras and algebraic quantum groups [VD7]. Much information about the purely algebraic theory can be found there and this can be helpful to understand the technically, far more difficult analytical theory. Also the original papers on the theory of multiplier Hopf $(*)$ -algebras [VD2] and [VD3], although certainly not necessary for understanding this paper, can be helpful.

We would like to emphasize the *importance of the original work* by Kustermans and Vaes, also for this alternative treatment. We do not really use results from their work, but certainly, we have been greatly inspired by their results and techniques. Without their pioneering work, this paper would not have been written. It is worthwhile mentioning that the PhD thesis of Vaes (for those who have access to this work) is easier to read than the original paper [K-V2]. Also the paper by Masuda, Nakagami and Woronowicz [M-N-W], treating independently the theory of locally compact quantum groups, has helped us to develop our new approach. Throughout the paper, we will not always repeat to refer to the original works, but we will do so where we feel this is appropriate.

Let us now finish this introduction with some *basic references* and *standard notations* used in this paper. We will also say something about the difference in conventions used in the field.

When \mathcal{H} is a Hilbert space, we will use $\mathcal{B}(\mathcal{H})$ to denote the von Neumann algebra of all bounded linear operators on \mathcal{H} . We use M_* for the space of normal linear functionals on a von Neumann algebra M and in particular $\mathcal{B}(\mathcal{H})_*$ for normal linear functionals on

$\mathcal{B}(\mathcal{H})$. When ω is a such a functional, we use $\bar{\omega}$ for the linear functional defined by $\bar{\omega}(x) = \omega(x^*)^- = \overline{\omega(x^*)}$ (where in all these cases, the $-$ stands for complex conjugation). On one occasion, we will also need the absolute value $|\omega|$ and the norm $\|\omega\|$ of a normal linear functional. If ξ and η are two vectors in the Hilbert space \mathcal{H} , we will write $\langle \cdot, \xi, \eta \rangle$ to denote the normal linear functional ω on $\mathcal{B}(\mathcal{H})$ given by $x \mapsto \langle x\xi, \eta \rangle$. In this case we have e.g. $\bar{\omega}(x) = \langle x\eta, \xi \rangle$ and, provided $\|\xi\| = \|\eta\| = 1$, that $|\omega|(x) = \langle x\xi, \xi \rangle$ and $\|\omega\| = 1$.

We refer to [S-Z], [T2] and [T3] for the theory of C*-algebras and von Neumann algebras. We will work with normal semi-finite weights on von Neumann algebras and with lower semi-continuous densely defined weights on C*-algebras. We will use the standard notations for the objects associated with such weights. If e.g. ψ is a normal semi-finite weight on a von Neumann algebra M , we will use \mathcal{N}_ψ for the left ideal of elements $x \in M$ such that $\psi(x^*x) < \infty$. Also \mathcal{M}_ψ will be the hereditary *-subalgebra $\mathcal{N}_\psi^* \mathcal{N}_\psi$ of M . We will use the G.N.S.-representation associated with such a weight. The Hilbert space will be denoted by \mathcal{H}_ψ while Λ_ψ is used for the canonical map from \mathcal{N}_ψ to the space \mathcal{H}_ψ . We will let the von Neumann algebra act directly on its G.N.S. space, i.e. we will drop the notation π_ψ . The modular operator on \mathcal{H}_ψ (in the case of a faithful weight) will be denoted by ∇ (and *not* by Δ because we reserve Δ for comultiplications). We will use (σ_t^ψ) for the modular automorphisms.

Again we refer to [T3] for the theory of weights on C*-algebras and von Neumann algebras, as well as for the modular theory and its relation with weights. See also [St].

We will be using various (continuous) one-parameter groups of automorphisms. We assume σ -weak continuity, but one can easily see that σ -weak continuity for one-parameter groups of *-automorphisms implies also continuity for the stronger operator topologies (like the σ -strong or even the σ -strong-* topology). We also have that the map $t \mapsto \omega \circ \alpha_t$ is continuous for any $\omega \in M_*$ with the norm topology on M_* when α is a continuous one-parameter group of automorphisms. We will be interested in analytical elements and the analytical generator $\alpha_{\frac{i}{2}}$ of such a one-parameter group α . A few things can be found in Chapter VIII of [T3], and a nice reference is also Appendix F in [M-N-W]. It seems to be better to first define the analytical generator for the action of \mathbb{R} on M_* , dual to the one-parameter group (because this is norm continuous), and define the analytical generator on M by taking the adjoint. Doing so, we get that the linear map $\alpha_{\frac{i}{2}}$ is closed for the σ -weak topology and that the analytical elements form a core with respect to the σ -strong-* topology.

When we write the tensor product of spaces, we will always mean completed tensor products. In the case of two Hilbert spaces, this is the Hilbert space tensor product. In the case of C*-algebras, it is understood to be the minimal C*-tensor product. Finally, for von Neumann algebras we take the usual von Neumann algebra (i.e. the spatial) tensor product.

Unfortunately, there are a number of different conventions used in this field by different authors/schools. In Hopf algebras e.g. it is common to endow the dual of a (finite-dimensional) Hopf algebra with a coproduct simply by dualizing the product whereas in the theory of locally compact quantum groups, usually the opposite coproduct on the dual

is taken. In the earlier works on Kac algebras (see [E-S]), the left regular representation is defined as the adjoint of what is commonly used now. Kustermans and Vaes work mainly with the left regular representation (as in the case of Kac algebras), whereas Baaĵ and Skandalis (in [B-S]) and Masuda, Nakagami and Woronowicz (in [M-N-W]) prefer the right regular representation as their starting point. Also a different convention in [M-N-W] is used for the polare decomposition of the antipode.

In this paper, we will mainly follow the conventions used by Kustermans and Vaes in their original papers. In a few occasions, mostly as a consequence of the difference in approach, we will choose slightly different conventions. In that case, we will clearly say so. It will only be the case in the process of obtaining the main results. In the formulation of the main results, we will be in accordance with the conventions in [K-V*].

Acknowledgements

Part of this work was done while I was on sabbatical in Trondheim (2002-2003). First of all, I would like to thank my colleague in Leuven, J. Quaegebeur, who did some of my teaching in Leuven and thus made it easier for me to go on sabbatical leave. It is also a pleasure to thank my colleagues of the University in Trondheim, M. Landstad and C. Skau, for their hospitality during my stay and for giving me the opportunity to talk about my work in their seminar. Part of this work was also done while visiting the University of Fukuoka (in 2004) and while visiting the University of Urbana-Champaign (in 2005). I would also like to thank A. Inoue, H. Kurose and Z.-J. Ruan for the hospitality during these (and earlier) visit(s).

I am also grateful to my coworkers in Leuven, especially J. Kustermans and S. Vaes, who have always been willing to share their competence in this field. This has been of great importance for my deeper understanding of the theory of locally compact quantum groups. Again, without this, the present paper would not have been written.

1. The antipode: construction and properties. Densities.

Let M be a von Neumann algebra and denote by $M \otimes M$ the von Neumann tensor product of M with itself. Recall the following definition which is the basic ingredient of this paper.

1.1 Definition Let Δ be a unital and normal $*$ -homomorphism from M to $M \otimes M$. Then Δ is called a *comultiplication* on M if $(\Delta \otimes \iota)\Delta = (\iota \otimes \Delta)\Delta$ (coassociativity), where ι is used to denote the identity map from M to itself.

The standard example comes from a locally compact group G . We take $M = L^\infty(G)$ and define Δ on M by $\Delta(f)(r, s) = f(rs)$ whenever $f \in L^\infty(G)$ and $r, s \in G$. We identify $M \otimes M$ with $L^\infty(G \times G)$.

A preliminary definition of the antipode and first properties

Let us now assume that M acts on the Hilbert space \mathcal{H} in standard form.

We first define the following subspace of the von Neumann algebra M .

1.2 Definition For an element $x \in M$ we say that $x \in \mathcal{D}_0$ if there is an element $x_1 \in M$ satisfying the following condition: For all $\varepsilon > 0$ and vectors $\xi_1, \xi_2, \dots, \xi_n, \eta_1, \eta_2, \dots, \eta_m$ in \mathcal{H} , there exist elements $p_1, p_2, \dots, p_m, q_1, q_2, \dots, q_m$ in M such that

$$\begin{aligned} \|x\xi_k \otimes \eta_k - \sum_j \Delta(p_j)(\xi_k \otimes q_j^* \eta_k)\| &< \varepsilon \\ \|x_1\xi_k \otimes \eta_k - \sum_j \Delta(q_j)(\xi_k \otimes p_j^* \eta_k)\| &< \varepsilon \end{aligned}$$

for all k .

We will see later that, because of forthcoming assumptions, we will have that $x_1 = 0$ if $x = 0$. Therefore, it would be possible to define a linear map S_0 on \mathcal{D}_0 by letting $S_0(x) = x_1^*$. For this map, we can prove the following properties:

- i) If $x \in \mathcal{D}_0$, then $S_0(x)^* \in \mathcal{D}_0$ and $S_0(S_0(x)^*)^* = x$.
- ii) If $x, y \in \mathcal{D}_0$, then $xy \in \mathcal{D}_0$ and $S_0(xy) = S_0(y)S_0(x)$.
- iii) The map $x \rightarrow S_0(x)^*$ is closed for the strong operator topology on M .

The properties i) and iii) are immediate consequences of the definition while ii) is obtained using a simple calculation.

This operator would be a candidate for the antipode (see the following remarks), but we will not define the antipode like this but rather through its polar decomposition (see the Definitions 1.22 and 1.23 later in this section).

1.3 Remarks i) The definition of S_0 above is inspired by a result in Hopf $(^*)$ -algebra theory. Indeed, if (H, Δ) is a Hopf algebra with antipode S and if $a \in H$, then using the Sweedler notation, we get

$$\begin{aligned} a \otimes 1 &= \sum a_{(1)} \otimes a_{(2)} S(a_{(3)}) \\ &= \sum \Delta(a_{(1)})(1 \otimes S(a_{(2)})). \end{aligned}$$

So, if we have a Hopf $(^*)$ -algebra, and if we write $\sum_j p_j \otimes q_j = \sum a_{(1)} \otimes S(a_{(2)})^*$, we get

$$\begin{aligned} \sum_j \Delta(q_j)(1 \otimes p_j^*) &= \sum \Delta(S(a_{(2)})^*)(1 \otimes a_{(1)}^*) \\ &= \sum S(a_{(3)})^* \otimes S(a_{(2)})^* a_{(1)}^* \\ &= S(a)^* \otimes 1. \end{aligned}$$

If we do not have a $*$ -structure, we have a similar formula, but we lose the symmetry.

ii) This formula can also be illustrated in the case $M = L^\infty(G)$ where G is a locally compact group. In this case we know that $S(f)(r) = f(r^{-1})$ when $f \in L^\infty(G)$ and $r \in G$. If we approximate

$$f(r) = f(rs \cdot s^{-1}) \simeq \sum_i p_i(rs) \overline{q_i(s^{-1})},$$

we get

$$\sum_i q_i(rs) \overline{p_i(s)} \simeq \overline{f(s \cdot (rs)^{-1})} = \overline{f(r^{-1})}.$$

iii) We have used the above idea in the construction of the antipode for Hopf C^* -algebras in [V-VD]. In fact, also the construction of the antipode in the paper [K-V2] uses this idea, but that is less obvious.

iv) The well-definedness of S_0 in the general case is a problem and it is also not clear whether or not there are even non-trivial elements in \mathcal{D}_0 . As we will see later in this section, the left and right Haar weights will be used to solve this problem.

iv) One of the nice aspects however of this definition of the antipode is that it does not depend on the possible choices of the left and the right Haar weights.

The involutive operator K implementing the antipode

In what follows, we will see how the existence of the Haar weights eventually leads to, not only the *well-definedness* of this preliminary antipode S_0 , but also the *density of the domain* \mathcal{D}_0 . But as we already mentioned, we will not define the antipode in this way. On the other hand, we will now do something similar and define a map like $S_0(\cdot)^*$, but on the Hilbert space level.

To do this, we will now assume the existence of a right Haar weight. We recall the definition (see e.g. [K-V3]).

1.4 Definition Let M be a von Neumann algebra and Δ a comultiplication on M (as in Definition 1.1). A *right Haar weight* on M is a faithful, normal semi-finite weight on M such that

$$\psi((\iota \otimes \omega)\Delta(x)) = \omega(1)\psi(x)$$

whenever $x \in M$, $x \geq 0$ and $\psi(x) < \infty$ and when $\omega \in M_*$ and $\omega \geq 0$ (right invariance).

We will now further in this section assume the existence of such a right Haar weight ψ . We will consider the G.N.S.-representation of M for ψ . Let \mathcal{N}_ψ be the set of elements $x \in M$ such that $\psi(x^*x) < \infty$. We will use Λ_ψ to denote the canonical map from the \mathcal{N}_ψ to \mathcal{H}_ψ . As usual, we extend ψ to the $*$ -subalgebra \mathcal{M}_ψ (defined as $\mathcal{N}_\psi^* \mathcal{N}_\psi$). We have $\langle \Lambda_\psi(x), \Lambda_\psi(y) \rangle = \psi(y^*x)$ for all $x, y \in \mathcal{N}_\psi$. We consider M as acting directly on \mathcal{H}_ψ (i.e. we drop the notation π_ψ) and so we will write $x\Lambda_\psi(y) = \Lambda_\psi(xy)$ when $x \in M$ and $y \in \mathcal{N}_\psi$. We refer to [St] and [T3] for details about weights and the G.N.S.-construction for weights.

Also, by now, the construction of the *right regular representation* has become standard. We recall it here and refer to e.g. [K-V2] and [K-V3] (and also [VD8]) for details.

1.5 Proposition There exists a bounded operator V from $\mathcal{H}_\psi \otimes \mathcal{H}$ to itself, characterized (and defined) by

$$((\iota \otimes \omega)V)\Lambda_\psi(x) = \Lambda_\psi((\iota \otimes \omega)\Delta(x))$$

whenever $x \in \mathcal{N}_\psi$ and $\omega \in \mathcal{B}(\mathcal{H})_*$. It has its 'second leg' in M , i.e. $V \in \mathcal{B}(\mathcal{H}_\psi) \otimes M$ and satisfies the following formulas:

- i) $V^*V = 1$ (i.e. V is an isometry),
- ii) $V(x \otimes 1) = \Delta(x)V$ for all $x \in M$,
- iii) $(\iota \otimes \Delta)V = V_{12}V_{13}$ (where we use the standard 'leg numbering' notation).

Roughly speaking we have $V(\Lambda_\psi(x) \otimes \xi) = \sum \Lambda_\psi(x_{(1)}) \otimes x_{(2)}\xi$ when we use the Sweedler notation $\Delta(x) = \sum x_{(1)} \otimes x_{(2)}$ for $x \in M$.

Recall that the invariance is used to get that $(\iota \otimes \omega)\Delta(x) \in \mathcal{N}_\psi$ when $x \in \mathcal{N}_\psi$, a result which is needed to define V as above. Also, it is the right invariance that implies that V is an isometry. It is known that in general, it seems impossible to show that V is a unitary without further assumptions. In our approach, we will get unitarity, in some sense, as a 'byproduct' of the further study of the antipode (see Proposition 1.15). Because we do not yet know that V is unitary, we need to formulate condition ii) as we have done and we can not (yet) write $\Delta(x) = V(x \otimes 1)V^*$. This will follow later.

It is easy to show that in the case $M = L^\infty(G)$, the operator V is indeed intimately related with the right regular representation of G on $L^2(G)$ (with the right Haar measure on G). The right Haar weight on $L^\infty(G)$ is obtained by integration with respect to this right Haar measure.

The *next step* is to construct the operator $S_0(\cdot)^*$ on the Hilbert space level, in this case, on \mathcal{H}_ψ . It will be denoted by K . The definition is very much as in Definition 1.2:

1.6 Definition Let $\xi \in \mathcal{H}_\psi$. We say that $\xi \in \mathcal{D}(K)$ if there is a vector $\xi_1 \in \mathcal{H}_\psi$ satisfying the following condition: For all $\varepsilon > 0$ and vectors $\eta_1, \eta_2, \dots, \eta_m$ in \mathcal{H}_ψ , there exist elements $p_1, p_2, \dots, p_m, q_1, q_2, \dots, q_m$ in \mathcal{N}_ψ such that

$$\begin{aligned} \|\xi \otimes \eta_k - V(\sum_j \Lambda_\psi(p_j) \otimes q_j^* \eta_k)\| &< \varepsilon \\ \|\xi_1 \otimes \eta_k - V(\sum_j \Lambda_\psi(q_j) \otimes p_j^* \eta_k)\| &< \varepsilon \end{aligned}$$

for all k .

Remark that this definition is indeed similar to Definition 1.2 because, roughly speaking, the operator V is the map $p \otimes q^* \mapsto \Delta(p)(1 \otimes q^*)$ on the Hilbert space level.

Again, we would like to define the operator K by $K\xi = \xi_1$ but we need the following result (which in this case, can be proven):

1.7 Lemma Let ξ and ξ_1 be as Definition 1.6 and assume $\xi = 0$. Then also $\xi_1 = 0$.

The idea of the proof is the following. Using $\xi = 0$ and the fact that V is isometric, we get a net of elements $\sum_j p_j \otimes q_j$ such that $\sum_j \Lambda_\psi(p_j) \otimes q_j^* \eta \rightarrow 0$ (using the first inequality) as well as that $\sum_j \Lambda_\psi(q_j) \otimes p_j^* \eta$ converges (by the second inequality) for all η . Standard techniques (about weights and the associated G.N.S.-representation) give that the maps involved are preclosed and hence also $\sum_j \Lambda_\psi(q_j) \otimes p_j^* \eta \rightarrow 0$ for all η . It follows that $\xi_1 = 0$. This argument is not fundamentally different from a similar argument in [K-V2]. For details, we refer to [VD8].

1.8 Definition If $\xi \in \mathcal{D}(K)$ and if ξ_1 is as in Definition 1.6, we set $K\xi = \xi_1$.

Remark that our operator K is essentially the operator G^* in the work of Kustermans and Vaes (as we will see later - cf. e.g. Remark 4.10 in Section 4). Therefore, it should not be a surprise that the techniques used above to define K and to show that it is well-defined are similar as those used in [K-V2]. Observe that we will not use the symbol G for this operator as it is used for a locally compact group.

Just as in the case of Definition 1.2, we get easily the following results:

1.9 Proposition i) If $\xi \in \mathcal{D}(K)$, then $K\xi \in \mathcal{D}(K)$ and $K(K\xi) = \xi$,
ii) K is a closed operator.

The counterpart of the other result for S_0 , namely that $S_0(xy) = S_0(y)S_0(x)$ when $x, y \in \mathcal{D}_0$, is the following:

1.10 Proposition Let $x \in \mathcal{D}_0$ and assume that x_1 is as in Definition 1.2. If $\xi \in \mathcal{D}(K)$ then $x\xi \in \mathcal{D}(K)$ and $Kx\xi = x_1K\xi$.

Again, the proof of this formula is straightforward and in fact of the same nature as the proof of the formula $S_0(xy)^* = S_0(x)^*S_0(y)^*$ when $x, y \in \mathcal{D}_0$.

As an *important consequence* of the above proposition, we find that, if $\mathcal{D}(K)$ is dense and if x and x_1 are as in Definition 1.2, then $x = 0$ will imply $x_1 = 0$. Indeed, it will follow that $x_1K\xi = 0$ for all $\xi \in \mathcal{K}$ and by Proposition 1.9 i) we have that the range of K is equal to $\mathcal{D}(K)$.

Density of the domain of the operator K

For the *following step*, we need a left Haar weight on M . It is used to produce (enough) elements in $\mathcal{D}(K)$ and in \mathcal{D}_0 . Recall that a *left Haar weight* is a faithful, normal semi-finite weight φ on M satisfying left invariance, i.e.

$$\varphi((\omega \otimes \iota)\Delta(x)) = \omega(1)\varphi(x)$$

whenever $x \in M$, $x \geq 0$ and $\varphi(x) < \infty$ and when $\omega \in M_*$ and $\omega \geq 0$. For a left Haar weight, we have the *left regular representation*. We use the G.N.S.-space \mathcal{H}_φ and $\Lambda_\varphi : \mathcal{N}_\varphi \rightarrow \mathcal{H}_\varphi$ for the associated canonical map. Again we let M act directly on \mathcal{H}_φ (i.e. we drop the notation π_φ as we did before with ψ). Later, we will identify the two Hilbert spaces \mathcal{H}_φ and \mathcal{H}_ψ in such a way that the actions of M are the same (see the end of Section 2).

The left regular representation is considered in the next proposition.

1.11 Proposition There is a bounded operator W on $\mathcal{H} \otimes \mathcal{H}_\varphi$, characterized (and defined) by

$$((\omega \otimes \iota)W^*)\Lambda_\varphi(x) = \Lambda_\varphi((\omega \otimes \iota)\Delta(x))$$

when $x \in \mathcal{N}_\varphi$ and $\omega \in \mathcal{B}(\mathcal{H})_*$. Now, the 'first leg' of W sits in M , that is $W \in M \otimes \mathcal{B}(\mathcal{H}_\varphi)$ and we have:

- i) $WW^* = 1$ (i.e. W is a co-isometry),
- ii) $(1 \otimes x)W = W\Delta(x)$ for all $x \in M$,
- iii) $(\Delta \otimes \iota)W = W_{13}W_{23}$.

Here, roughly speaking, we have $W^*(\xi \otimes \Lambda_\varphi(x)) = \sum x_{(1)}\xi \otimes \Lambda_\varphi(x_{(2)})$ when $\Delta(x) = \sum x_{(1)} \otimes x_{(2)}$ formally. Observe the difference in convention (using the adjoint) when compared with the right regular representation. The proof of this proposition however is completely similar as for the right regular representation.

In order to use W to construct elements in $\mathcal{D}(K)$ and in \mathcal{D}_0 , we need different steps. We formulate different lemmas as some of the results will be needed later. First we have the following:

1.12 Lemma Let $\omega \in \mathcal{B}(\mathcal{H}_\varphi)_*$ and $x = (\iota \otimes \omega)W$ and $x_1 = (\iota \otimes \bar{\omega})W$, then $x \in \mathcal{D}_0$ and x_1 satisfies the conditions as in Definition 1.2.

Proof (sketch): Assume that $\omega = \langle \cdot, \xi, \eta \rangle$. Take an orthonormal basis (ξ_j) in \mathcal{H}_φ . Define

$$\begin{aligned} p_j &= (\iota \otimes \langle \cdot, \xi_j, \eta \rangle)W \\ q_j &= (\iota \otimes \langle \cdot, \xi_j, \xi \rangle)W. \end{aligned}$$

Using the formula $(\Delta \otimes \iota)W = W_{13}W_{23}$ (Proposition 1.11), we find

$$\begin{aligned} \sum_j \Delta(p_j)(1 \otimes q_j^*) &= (\iota \otimes \iota \otimes \omega)(W_{13}W_{23}W_{23}^*) \\ &= (\iota \otimes \iota \otimes \omega)W_{13} \\ &= x \otimes 1 \end{aligned}$$

and similarly

$$\begin{aligned} \sum_j \Delta(q_j)(1 \otimes p_j^*) &= (\iota \otimes \iota \otimes \bar{\omega})(W_{13}W_{23}W_{23}^*) \\ &= (\iota \otimes \iota \otimes \bar{\omega})W_{13} \\ &= x_1 \otimes 1. \end{aligned}$$

The sums converge in the strong operator topology.

This gives the result for elements ω of the form $\langle \cdot, \xi, \eta \rangle$. Then it follows for all $\omega \in \mathcal{B}(\mathcal{H}_\varphi)_*$ by approximation.

Remark that only the essential properties of W are used in the above argument and that it is not necessary to have a left regular representation, associated to a left Haar weight. Only the conditions i) and iii) of Proposition 1.11 are needed.

Compare this lemma with Proposition 5.6 in [V-VD] where a similar argument is found. Observe again that one of the differences between this approach to the antipode and the one in [V-VD] lies in the fact that we avoid the use of operator space techniques here.

Later, we will combine this result with the property proven in Proposition 1.10 (cf. Proposition 1.16).

In a similar way, elements in the domain of K are constructed, but here, we have to be a bit more careful. First, we have the following lemma.

1.13 Lemma If $c \in \mathcal{N}_\psi$ and $\omega \in \mathcal{B}(\mathcal{H}_\varphi)_*$ we have $(\iota \otimes \omega(c \cdot))W \in \mathcal{N}_\psi$.

Proof: If we let $x = (\iota \otimes \omega(c \cdot))W$, we get

$$\begin{aligned} x^*x &\leq \|\omega\|(\iota \otimes |\omega|)(W^*(1 \otimes c^*)(1 \otimes c)W) \\ &= \|\omega\|(\iota \otimes |\omega|)(\Delta(c^*)W^*W\Delta(c)) \\ &\leq \|\omega\|(\iota \otimes |\omega|)(\Delta(c^*)\Delta(c)) \\ &= \|\omega\|(\iota \otimes |\omega|)(\Delta(c^*c)). \end{aligned}$$

As ψ is right invariant and $c \in \mathcal{N}_\psi$, we get also $x \in \mathcal{N}_\psi$.

Observe that we do not need that W is unitary. It is sufficient for this argument that $W^*W \leq 1$ and this is true for a co-isometry.

Now, the following result should not come as a surprise.

1.14 Lemma Let $c, d \in \mathcal{N}_\psi$ and $\omega \in \mathcal{B}(\mathcal{H}_\varphi)_*$ and define $\xi = \Lambda_\psi((\iota \otimes \omega(c \cdot d^*))W)$. Then $\xi \in \mathcal{D}(K)$ and $K\xi = \Lambda_\psi((\iota \otimes \bar{\omega}(d \cdot c^*))W)$.

Proof (sketch): The proof of this lemma is based on the same decomposition as in Lemma 1.12. Again we take $\omega = \langle \cdot, \xi', \eta' \rangle$ and now

$$\begin{aligned} p_j &= (\iota \otimes \langle \cdot, \xi_j, c^* \eta' \rangle)W \\ q_j &= (\iota \otimes \langle \cdot, \xi_j, d^* \xi' \rangle)W. \end{aligned}$$

By Lemma 1.13 we have $p_j, q_j \in \mathcal{N}_\psi$. This is necessary for the use of Definition 1.6.

Having these results, we are ready to show that the domain of K is dense. Simultaneously, we obtain that the right regular representation is unitary.

- 1.15 Proposition** i) V is unitary,
ii) $\mathcal{D}(K)$ is dense.

Proof (sketch): The proof of i) uses Kustermans' trick as in [K-V2]. Define

$$\mathcal{K} = \overline{\text{sp}}\{\Lambda_\psi((\iota \otimes \omega(c \cdot))W) \mid c \in \mathcal{N}_\psi, \omega \in \mathcal{B}(\mathcal{H}_\varphi)_*\}$$

where by $\overline{\text{sp}}$ we mean that we are taking the closed linear span. Consider V as acting on the space $\mathcal{H}_\psi \otimes \mathcal{H}_\varphi$ (by taking \mathcal{H}_φ for \mathcal{H}). Consider the notations of Lemma 1.14, but with $d = 1$. Using the same techniques, we find that $\xi \otimes \eta$ is approximated by finite sums of the form $\sum V(\Lambda_\psi(p_j) \otimes q_j^* \eta)$ for any $\eta \in \mathcal{H}_\varphi$. Because ξ , as well as all the elements $\Lambda_\psi(p_j)$ belong to \mathcal{K} , we find that $\mathcal{K} \otimes \mathcal{H}_\varphi \subseteq V(\mathcal{K} \otimes \mathcal{H}_\varphi)$.

On the other hand, we have the formula

$$(\iota \otimes \varphi)((\Delta(x^*)(1 \otimes y)) = (\iota \otimes \langle \cdot, \Lambda_\varphi(y), \Lambda_\varphi(x) \rangle)W$$

whenever $x, y \in \mathcal{N}_\varphi$. Using this result, and some other results about weights and careful approximation techniques, one can show that also

$$\mathcal{K} = \overline{\text{sp}}\{\Lambda_\psi((\iota \otimes \omega)\Delta(x)) \mid x \in \mathcal{N}_\psi, \omega \in M_*\}$$

and consequently that $V(\mathcal{H}_\psi \otimes \mathcal{H}_\varphi) \subseteq \mathcal{K} \otimes \mathcal{H}_\varphi$.

By a combination of the two results above and using that V is isometric, we get $\mathcal{K} = \mathcal{H}_\psi$. Therefore, V is unitary.

As we also have

$$\mathcal{K} = \overline{\text{sp}}\{\Lambda_\psi((\iota \otimes \omega(c \cdot d^*))W) \mid c, d \in \mathcal{N}_\psi, \omega \in \mathcal{B}(\mathcal{H}_\varphi)_*\},$$

we get from $\mathcal{K} = \mathcal{H}_\psi$ that $\mathcal{D}(K)$ is dense.

Compare the proof of this proposition with arguments found in Section 3.3 of [K-V2].

By symmetry, of course also the left regular representation W associated to any left Haar weight will be unitary. Observe that we now can rewrite the formulas ii) of Proposition 1.5 and ii) of Proposition 1.11 as $\Delta(x) = V(x \otimes 1)V^*$ and $\Delta(x) = W^*(1 \otimes x)W$ respectively.

The unitarity of the regular representations can also be proven in an other, perhaps shorter (still essentially the same) way, but because we also need the density of the domain $\mathcal{D}(K)$ of K , we have chosen to prove these results together as above.

It should not come as a surprise that the density of $\mathcal{D}(K)$ is essentially the same result as saying that the isometry V is in fact a unitary. Indeed, when $\mathcal{D}(K)$ is dense, it follows that for any $\xi \in \mathcal{H}_\psi$ and any $\eta \in \mathcal{H}$, the vector $\xi \otimes \eta$ can be approximated with elements in the range of V (cf. Definition 1.6). Roughly speaking, this says that the map $p \otimes q \mapsto \Delta(p)(1 \otimes q)$, considered on the Hilbert space level, has dense range. Later, at the end of this section, we will see that this map on $M \otimes M$ also has dense range. This in turn will be a consequence of the density of \mathcal{D}_0 (cf. Proposition 1.21 below). Remark that,

although there are similarities, the two density results are different because the topologies considered are different.

The antipode and its polar decomposition

We have now shown that the domain $\mathcal{D}(K)$ of the operator K is dense and as we mentioned already (see the remark following Proposition 1.10) it would now be possible to define the antipode as the map S_0 given by $S_0(x) = x_1^*$ (cf. a remark following Definition 1.2). It would still be necessary to show that also the domain \mathcal{D}_0 of S_0 is dense.

Eventually, we will see that indeed, the space \mathcal{D}_0 is dense (see Proposition 1.21 below). But first, we will construct the antipode by means of its polar decomposition. Some of the formulas needed to do this will also play an important role in the next section where we obtain the main results.

Let us first formulate a result that easily follows from combining Lemma 1.12 with Proposition 1.10.

1.16 Proposition For any $\xi \in \mathcal{D}(K)$ and $\omega \in \mathcal{B}(\mathcal{H}_\varphi)_*$, we have that $((\iota \otimes \omega)W)\xi \in \mathcal{D}(K)$ and

$$K((\iota \otimes \omega)W)\xi = ((\iota \otimes \bar{\omega})W)K\xi.$$

Next, we need a similar formula, but for the other leg of W . And because eventually, we will need all of this to obtain uniqueness of the Haar weights, we will work with two left Haar weights φ_1 and φ_2 . We will use the left regular representations for these two left Haar weights and we will use W_1 and W_2 to denote them. We will in what follows consider these operators as acting on the spaces $\mathcal{H}_\psi \otimes \mathcal{H}_{\varphi_1}$ and $\mathcal{H}_\psi \otimes \mathcal{H}_{\varphi_2}$ respectively.

We then have the following result.

1.17 Proposition Let T_r be the closure of the operator $\Lambda_{\varphi_1}(x) \mapsto \Lambda_{\varphi_2}(x^*)$ with $x \in \mathcal{N}_{\varphi_1} \cap \mathcal{N}_{\varphi_2}^*$. If $\xi \in \mathcal{D}(T_r)$ then $((\omega \otimes \iota)W_1^*)\xi \in \mathcal{D}(T_r)$ for all $\omega \in \mathcal{B}(\mathcal{H}_\psi)_*$ and

$$T_r((\omega \otimes \iota)W_1^*)\xi = ((\bar{\omega} \otimes \iota)W_2^*)T_r\xi.$$

Proof: Fix $\omega \in \mathcal{B}(\mathcal{H}_\psi)_*$. First, we prove the formula for $\xi = \Lambda_{\varphi_1}(x)$ with $x \in \mathcal{N}_{\varphi_1} \cap \mathcal{N}_{\varphi_2}^*$. We get

$$((\omega \otimes \iota)W_1^*)\Lambda_{\varphi_1}(x) = \Lambda_{\varphi_1}((\omega \otimes \iota)\Delta(x))$$

by the definition of W_1 . Then, because also $(\omega \otimes \iota)\Delta(x) \in \mathcal{N}_{\varphi_1} \cap \mathcal{N}_{\varphi_2}^*$, we get from the definition of T_r that

$$\begin{aligned} T_r(((\omega \otimes \iota)W_1^*)\Lambda_{\varphi_1}(x)) &= \Lambda_{\varphi_2}((\bar{\omega} \otimes \iota)\Delta(x^*)) \\ &= ((\bar{\omega} \otimes \iota)W_2^*)\Lambda_{\varphi_2}(x^*) \\ &= ((\bar{\omega} \otimes \iota)W_2^*)T_r\Lambda_{\varphi_1}(x). \end{aligned}$$

The general case follows because T_r is the closure of the map $\Lambda_{\varphi_1}(x) \mapsto \Lambda_{\varphi_2}(x^*)$ with $x \in \mathcal{N}_{\varphi_1} \cap \mathcal{N}_{\varphi_2}^*$.

Now, we will combine the above result with the similar formula for K , applied for both W_1 and W_2 . Compare with results in Section 5.2 of [K-V2].

1.18 Proposition With the notations as before, we have the equality

$$(K \otimes T_r)W_1 = W_2^*(K \otimes T_r).$$

Proof (sketch): Take vectors $\xi \in \mathcal{D}(K)$, $\xi' \in \mathcal{D}(K^*)$, $\eta \in \mathcal{D}(T_r)$ and $\eta' \in \mathcal{D}(T_r^*)$. Remember that $\xi, \xi' \in \mathcal{H}_\psi$ while $\eta \in \mathcal{H}_{\varphi_1}$ and $\eta' \in \mathcal{H}_{\varphi_2}$.

If we first use the formula for K and then the similar formula for T_r (from Proposition 1.17), we find, after a careful but straightforward calculation, that

$$\langle W_2(K\xi \otimes T_r\eta), \xi' \otimes \eta' \rangle = \langle W_1^*(\xi \otimes \eta), K^*\xi' \otimes T_r^*\eta' \rangle^-.$$

This implies the inclusion $W_2(K \otimes T_r) \subseteq (K \otimes T_r)W_1^*$.

On the other hand, if we first use the formula for T_r and then the formula for K , we find

$$\langle W_2^*(K\xi \otimes T_r\eta), \xi' \otimes \eta' \rangle = \langle W_1(\xi \otimes \eta), K^*\xi' \otimes T_r^*\eta' \rangle^-.$$

This implies the inclusion $W_2^*(K \otimes T_r) \subseteq (K \otimes T_r)W_1$.

If we take the first inclusion and multiply with W_2^* from the left and with W_1 from the right, we get $(K \otimes T_r)W_1 \subseteq W_2^*(K \otimes T_r)$ and if we combine this with the previous inclusion, we get the result.

This formula is very important for the further development in Section 2. Of course, we can also replace both W_1 and W_2 by W associated with any left Haar weight φ . We will use both cases in the next section when we show that Haar weights are unique. In this section, we will use it (with W) to construct the antipode and to prove some more density results as we announced.

In order to use our formula, we need to consider the polar decomposition of the operators involved.

1.19 Notation Let K be the operator on \mathcal{H}_ψ as defined in Definition 1.6 and Definition 1.8. Now, let T be the closure of the map $\Lambda_\varphi(x) \mapsto \Lambda_\varphi(x^*)$ where $x \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$. We use

$$K = IL^{\frac{1}{2}} \quad \text{and} \quad T = J\nabla^{\frac{1}{2}}$$

to denote the polar decompositions of these operators.

The properties of all these operators are well-known and easy consequences of the fact that K and T are conjugate linear and involutive. We have e.g. that $J\nabla J = \nabla^{-1}$ so that

$J\nabla^{it}J = \nabla^{it}$ (because J is conjugate linear). Similarly for the operators I and L . See e.g. Chapter VI in [T3].

Remember that, roughly speaking, our operator K coincides with the operator G^* in [K-V2] and therefore, that the operator L is essentially the operator N^{-1} in [K-V2] (see Section 5 in [K-V2]). See also Section 4, in particular Remark 4.10.

If we apply Proposition 1.18 to the case $\varphi_1 = \varphi_2 = \varphi$, we get $(K \otimes T)W = W^*(K \otimes T)$ where W is the left regular representation associated with φ . As a consequence of the uniqueness of the polar decomposition, we get the following result.

1.20 Proposition i) $(I \otimes J)W(I \otimes J) = W^*$,
ii) $(L^{it} \otimes \nabla^{it})W(L^{-it} \otimes \nabla^{-it}) = W$ for all $t \in \mathbb{R}$.

In the next section, we will use a similar formula ii), but for two weights and we will combine it with this formula ii) here to get uniqueness of the Haar weights.

We will now show in the next proposition that the left leg of W is dense in M and therefore, the above formulas will allow us to define maps $R : M \rightarrow M$ and $\tau_t : M \rightarrow M$ for all t by $R(x) = Ix^*I$ and $\tau_t(x) = L^{it}xL^{-it}$. These maps will give us the polar decomposition of the antipode (see Definition 1.23 below).

We first need the following observation. Denote by $(\sigma_t^\varphi)_{t \in \mathbb{R}}$ the modular automorphisms on M defined by $\sigma_t^\varphi(x) = \nabla^{it}x\nabla^{-it}$. Similarly, let us define the one-parameter group of automorphisms (τ_t) on $\mathcal{B}(\mathcal{H}_\psi)$ by $\tau_t = L^{it} \cdot L^{-it}$. Then it follows from the second formula in Proposition 1.20 and from $\Delta(x) = W^*(1 \otimes x)W$ for all $x \in M$ that $\Delta(\sigma_t^\varphi(x)) = (\tau_t \otimes \sigma_t^\varphi)\Delta(x)$ for all $x \in M$. From this formula, it follows that the space of slices, spanned by the elements $(\omega \otimes \iota)\Delta(x)$ with $x \in M$ and $\omega \in M_*$ will be left invariant by the modular automorphisms $(\sigma_t^\varphi)_{t \in \mathbb{R}}$. We will need this for the proof of the following proposition (see Proposition 1.4 in [K-V2]).

1.21 Proposition Let W be the left regular representation associated with some left Haar weight φ as before. Then the following three subspaces of M

- i) $\{(\iota \otimes \omega)W \mid \omega \in \mathcal{B}(\mathcal{H}_\varphi)_*\}$,
 - ii) $\text{sp}\{(\iota \otimes \omega)\Delta(x) \mid x \in M, \omega \in M_*\}$,
 - iii) $\text{sp}\{(\omega \otimes \iota)\Delta(x) \mid x \in M, \omega \in M_*\}$
- are σ -weakly dense in M .

Proof: We will only consider i) and ii) because the density in iii) will follow by symmetry. The spaces in i) and in ii) have the same closure. To show this, we use again the formula

$$(\iota \otimes \varphi)((\Delta(x^*)(1 \otimes y)) = (\iota \otimes \langle \cdot, \Lambda_\varphi(y), \Lambda_\varphi(x) \rangle)W$$

with $x, y \in \mathcal{N}_\varphi$.

Let us now denote by M_e the closure of the space in i). It follows from the fact that W satisfies the pentagon equation, that M_e is a subalgebra of M . Because M_e is

also the closure of the space in ii) and this is obviously self-adjoint, we get that M_e is a $*$ -subalgebra of M .

In the proof of Proposition 1.15, we have seen that the space

$$\text{sp}\{\Lambda_\psi((\iota \otimes \omega)\Delta(x)) \mid x \in \mathcal{N}_\psi, \omega \in M_*\}$$

is dense in \mathcal{H}_ψ . Standard approximation techniques give that also

$$\text{sp}\{\Lambda_\psi((\iota \otimes \omega)\Delta(x)) \mid x \in \mathcal{N}_\psi \cap \mathcal{N}_\psi^*, \omega \in M_*\}$$

is still dense in \mathcal{H}_ψ . This will imply that the space $\Lambda_\psi(\mathcal{N}_\psi \cap \mathcal{N}_\psi^* \cap M_e)$ is dense in $\Lambda_\psi(\mathcal{N}_\psi \cap \mathcal{N}_\psi^*)$.

We have seen in a remark preceding this proposition, that the space of slices in iii) is invariant by the modular automorphisms of φ . Similarly, the modular automorphism σ_t^ψ leaves M_e invariant. So the space $\Lambda_\psi(\mathcal{N}_\psi \cap \mathcal{N}_\psi^* \cap M_e)$ will be invariant under the modular unitaries ∇^{it} (where we use ∇ for the modular operator associated with the right Haar weight ψ). It follows that also $\Lambda_\psi(\mathcal{N}_\psi \cap \mathcal{N}_\psi^* \cap M_e)$ is dense in $\Lambda_\psi(\mathcal{N}_\psi \cap \mathcal{N}_\psi^*)$ with respect to the $\#$ -norm (cf. Section 1 in Chapter VI in [T3]). Then, from a result in Hilbert algebra theory (see Lemma 5.1 and the proof of Theorem 10.1 in [T1]), it will follow that also $\mathcal{N}_\psi \cap \mathcal{N}_\psi^* \cap M_e$ is dense in M . Therefore we have that $M_e = M$. This completes the proof of the proposition.

From this proposition and taking into account Lemma 1.12, it follows that the map $p \otimes q \mapsto \Delta(p)(1 \otimes q)$ has dense range in $M \otimes M$. Indeed, one can approximate elements of the form $x \otimes 1$ by linear combinations of elements of the form $\Delta(p)(1 \otimes q)$ when $x \in \mathcal{D}_0$ (by the very definition of \mathcal{D}_0) and as now this domain is shown to be dense, we can do this for any $x \in M$. By multiplying with elements of M in the second factor, we get the density of $\Delta(M)(1 \otimes M)$. By symmetry, we also have that $\Delta(M)(M \otimes 1)$ will be dense in $M \otimes M$. Also compare with the remark following Proposition 1.15.

We *finish this section* by the definition of the antipode S and its polar decomposition and by formulating an important property which will be frequently used in the next sections.

1.22 Definition Define $R : M \rightarrow M$ by $R(x) = Ix^*I$ and $\tau_t : M \rightarrow M$ by $\tau_t(x) = L^{it}xL^{-it}$.

It is a consequence of the density result i) in Proposition 1.21 that these maps do leave M invariant. We have that R and τ_t commute because I and L^{it} commute for all t . The automorphisms (τ_t) are called *the scaling automorphisms* whereas the anti-automorphism R is called *the unitary antipode*. Together they represent what is commonly referred to as *the polar decomposition of the antipode*:

1.23 Definition The antipode S is defined as the composition $R\tau_{-\frac{i}{2}}$ where $\tau_{-\frac{i}{2}}$ is the analytic generator associated with the one-parameter group (τ_t) in the point $-\frac{i}{2}$

Recall, as was already mentioned in the introduction, that the analytic generator is defined first on the predual M_* and then on M by taking the adjoint.

In the next sections, we will (essentially) no longer need the operators K , I and L (they will be replaced by other, more adapted operators, see Section 4). We will use the unitary antipode R and the scaling group (τ_t) . The main results in this section involving K , I and L can be restated solely in terms of R and τ_t . This is e.g. quite obvious for the two formulas in Proposition 1.20 (and for the similar result involving two left Haar weights, see a remark in the beginning of Section 2, before Proposition 2.2).

It is somewhat harder with Proposition 1.16, but the following result can be shown about the antipode as defined in 1.23.

1.24 Proposition For any $\omega \in \mathcal{B}(\mathcal{H}_\varphi)_*$ we have that $(\iota \otimes \omega)W \in \mathcal{D}(S)$ and $S((\iota \otimes \omega)W) = (\iota \otimes \omega)(W^*)$. The space of such elements is invariant under the scaling automorphisms (τ_t) and it is a core for S .

Indeed, formally, from Proposition 1.16, we get $KxK = x_1^*$ when $x = (\iota \otimes \omega)W$ and $x_1 = (\iota \otimes \omega)(W^*)$. Now

$$KxK = IL^{\frac{1}{2}}xL^{-\frac{1}{2}}I = R(\tau_{-\frac{i}{2}}(x))^*.$$

One has to be somewhat careful, but the argument can be made precise. Furthermore, because of the second formula in Proposition 1.20, the space of such elements $(\iota \otimes \omega)W$ will be invariant under the scaling automorphisms and as this space is dense in M , it will be a core for S . See e.g. [VD8] for details.

We will refer to this result when in the sequel, we write loosely $(S \otimes \iota)W = W^*$.

2. The main results about (M, Δ)

In Section 1, we have introduced the antipode and obtained some results about density. We also proved an important formula and some consequences of it. In this section, we will prove the main results about a locally compact quantum group and its related objects (such as the left and right Haar weights, the modular automorphism groups, the scaling group and the unitary antipode, ...). In the next section, we will treat the dual and in the fourth section we will prove the main results about the objects associated with a locally compact quantum group in relation with those of the dual.

But first, it is appropriate to formulate here the precise definition of a locally compact quantum group (in the von Neumann algebraic setting), see [K-V3].

2.1 Definition Let M be a von Neumann algebra. Let Δ be a comultiplication on M . The pair (M, Δ) is called a *locally compact quantum group* if there exist a left and a right Haar weight.

The correct definition of a comultiplication on a von Neumann algebra M was given in Section 1 (Definition 1.1). It is a unital normal $*$ -homomorphism from M to $M \otimes M$ satisfying coassociativity. A right Haar weight is a faithful normal semi-finite weight on M that is right invariant (again see Section 1, Definition 1.4). A left Haar weight is a faithful normal semi-finite weight on M that is left invariant.

In this section, (M, Δ) will be a locally compact quantum group (in the sense of the above definition) and ψ will be a right Haar weight and φ, φ_1 and φ_2 will denote left Haar weights. We will use the notations and results of the previous section.

A first major objective is to prove that Haar weights are unique. We consider the two left Haar weights φ_1 and φ_2 . We denote by $(u_t)_{t \in \mathbb{R}}$ the Connes' cocycle Radon Nikodym derivative of φ_1 w.r.t. φ_2 and we will show that u_t is a scalar multiple of 1 for each t . This will imply that the two weights are proportional.

Our starting point will be the formula in Proposition 1.18. Whereas in the previous section, we have used this formula for the case of one invariant weight, now we will also use it for the two different weights φ_1 and φ_2 .

Recall that we use W_1 and W_2 for the left regular representations associated with φ_1 and φ_2 respectively. And as in the previous section, denote by T_r the closure of the operator $\Lambda_{\varphi_1}(x) \mapsto \Lambda_{\varphi_2}(x^*)$ where $x \in \mathcal{N}_{\varphi_1} \cap \mathcal{N}_{\varphi_2}^*$. Let $T_r = J_r \nabla_r^{\frac{1}{2}}$ be the polar decomposition. Remark that ∇_r is an operator on \mathcal{H}_{φ_1} and that J_r maps \mathcal{H}_{φ_1} to \mathcal{H}_{φ_2} . From Proposition 1.18, we know that $(K \otimes T_r)W_1 = W_2^*(K \otimes T_r)$. Then, as in Proposition 1.20, it follows from the uniqueness of the polar decompositions that $L^{it} \otimes \nabla_r^{it}$ commutes with W_1 for all t . This result can be restated without the use of the operator L . We simply write e.g. $(\tau_t \otimes \iota)W_1 = (1 \otimes \nabla_r^{-it})W_1(1 \otimes \nabla_r^{it})$. We refer to a remark following Definition 1.23.

Let us also use T_1 for the closure of the operator $\Lambda_{\varphi_1}(x) \mapsto \Lambda_{\varphi_1}(x^*)$ with $x \in \mathcal{N}_{\varphi_1} \cap \mathcal{N}_{\varphi_1}^*$ and $T_1 = J_1 \nabla_1^{\frac{1}{2}}$ for its polar decomposition. It is known from the *relative modular theory* (see e.g. Section 3 in Chapter VIII of [T3]) that $u_t = \nabla_1^{it} \nabla_r^{-it}$. Then the following result follows easily.

2.2 Proposition For each $t \in \mathbb{R}$ we have $\Delta(u_t) = 1 \otimes u_t$.

Proof: We just saw that $(L^{it} \otimes \nabla_r^{it})W_1(L^{-it} \otimes \nabla_r^{-it}) = W_1$ for all $t \in \mathbb{R}$. If we combine this with the formula ii) in Proposition 1.20 of the previous section (for W_1), we get $(1 \otimes u_t)W_1(1 \otimes u_t^*) = W_1$. Because $W_1^*(1 \otimes x)W_1 = \Delta(x)$ for all $x \in M$, we get the result.

If the right Haar weight ψ is bounded, we can immediately apply it on the formula above and get $\psi(u_t)1 = \psi(1)u_t$ so that u_t must be a scalar multiple of the identity for all t . We will be able to conclude this, also in general, but then we need a finer argument.

Before we complete this argument, recall that we have defined $R : M \rightarrow M$ by $R(x) = Ix^*I$ and $\tau_t : M \rightarrow M$ by $\tau_t(x) = L^{it}xL^{-it}$ (see Definition 1.22). Again, we will also use $(\sigma_t^\varphi)_{t \in \mathbb{R}}$ to denote the modular automorphisms on M defined by $\sigma_t^\varphi(x) = \nabla^{it}x\nabla^{-it}$.

We get a first set of important formulas:

2.3 Theorem For all $x \in M$ and $t \in \mathbb{R}$ we have:

- i) $\Delta(\sigma_t^\varphi(x)) = (\tau_t \otimes \sigma_t^\varphi)\Delta(x)$,
- ii) $\Delta(\tau_t(x)) = (\tau_t \otimes \tau_t)\Delta(x)$,
- iii) $\Delta(R(x)) = (R \otimes R)\Delta'(x)$ where Δ' is obtained from Δ by applying the flip.

Proof: As we saw already, the first formula is an immediate consequence of the second formula in Proposition 1.20 because $\Delta(x) = W^*(1 \otimes x)W$ for all $x \in M$. The formulas ii) and iii) follow in a straightforward way from the definitions and the formulas in Proposition 1.20 and again $\Delta(x) = W^*(1 \otimes x)W$, combined also with $(\Delta \otimes \iota)W = W_{13}W_{23}$. In fact, formula ii) can also be obtained from i) using coassociativity and the density ii) in Proposition 1.21.

In order to complete the proof of the uniqueness of the left Haar weights, we need the following result.

2.4 Proposition If $x \in M$ and $\Delta(x) = 1 \otimes x$ or $\Delta(x) = x \otimes 1$, then x must be a scalar multiple of 1.

Proof: We will assume that $\Delta(x) = x \otimes 1$ and prove that x is a scalar multiple of 1. The other property (which is the one we really need) will follow by symmetry.

So assume that $x \in M$ and that $\Delta(x) = x \otimes 1$. Take a function f on \mathbb{R} of the type $f(t) = \exp(-p(t - q)^2)$ with $p, q \in \mathbb{R}$ and define

$$y = \int f(t)\sigma_t^\varphi(x) dt$$

$$z = \int f(t)\tau_t(x) dt.$$

Because $\Delta(x) = x \otimes 1$ and $\Delta(\sigma_t^\varphi(x)) = (\tau_t \otimes \sigma_t^\varphi)\Delta(x) = \tau_t(x) \otimes 1$, we get $\Delta(y) = z \otimes 1$. Take any $w \in \mathcal{M}_\varphi (= \mathcal{N}_\varphi^* \mathcal{N}_\varphi)$. Then, because y is analytic with respect to (σ_t^φ) , we have also $yw \in \mathcal{M}_\varphi$ (see e.g. Section 2 of Chapter VIII in [T3]). Therefore we can apply φ to the second leg of the equation $\Delta(yw) = (z \otimes 1)\Delta(w)$ and use left invariance of φ to get $\varphi(yw)1 = \varphi(w)z$. Because this holds for all $w \in \mathcal{M}_\varphi$, it follows from the faithfulness of φ that y and z are scalar multiples of the identity. Because this is true for all such functions f , this can only happen when x itself is a scalar multiple of 1.

Now we are ready to obtain the uniqueness of the Haar weights.

2.5 Theorem Any two left Haar weights on a locally compact quantum group are equal (up to a scalar). Similarly for right Haar weights.

Proof: Combining Proposition 2.2 where we have shown that $\Delta(u_t) = 1 \otimes u_t$ for all t , and the previous result, we find that u_t is a scalar multiple of 1 for all t . This implies that the two weights φ_1 and φ_2 are proportional. The result for right Haar weights follows by symmetry.

Because of this result, we will in what follows use (σ_t) for the modular automorphisms of a left Haar weight (in stead of (σ_t^φ) as we did before). We will use (σ'_t) for the modular automorphisms of a right Haar weight.

From the fact that R flips the coproduct, it follows that $\varphi \circ R$ is right invariant when φ is left invariant. From now on, we will assume that φ is *fixed left Haar weight* and we will take *the right Haar weight* ψ to be this composition $\varphi \circ R$.

Now, here is another couple of formulas.

2.6 Theorem For all $x \in M$ and $t \in \mathbb{R}$ we have:

- i) $R(\sigma_t(x)) = \sigma'_{-t}(R(x))$,
- ii) $\Delta(\sigma'_t(x)) = (\sigma'_t \otimes \tau_{-t})\Delta(x)$.

The first property is a consequence of $\psi = \varphi \circ R$ and the fact that R is a anti-homomorphism. It can be shown using e.g. the K.M.S. property of weights w.r.t. the modular automorphisms. Having i), clearly ii) will follow from i) and iii) in Theorem 2.3. We could have obtained the second formula of this theorem also by symmetry, but then we could not be sure that we had the same scaling group.

From the uniqueness of the Haar weights, we get easily that the scaling automorphisms (τ_t) leave the Haar weights relatively invariant:

2.7 Theorem There exists a strictly positive number ν so that $\varphi \circ \tau_t = \nu^{-t}\varphi$ and $\psi \circ \tau_t = \nu^{-t}\psi$ for all $t \in \mathbb{R}$.

Proof: It follows from $\Delta \circ \tau_t = (\tau_t \otimes \tau_t) \circ \Delta$ that $\varphi \circ \tau_t$ is also left invariant and so, by uniqueness, it is a scalar multiple of φ . As this is true for all τ_t , we get a strictly positive number ν such that $\varphi \circ \tau_t = \nu^{-t}\varphi$ for all t . Composing with R and using that R and τ_t commute, we get also $\psi \circ \tau_t = \nu^{-t}\psi$ for all t .

From the fact that φ is relatively invariant under τ_t , it follows that τ_t commutes with all the modular automorphisms $(\sigma_s)_{s \in \mathbb{R}}$ of the left Haar weight. Similarly, τ_t will commute with all the modular automorphisms $(\sigma'_s)_{s \in \mathbb{R}}$ of the right Haar weight. It is not so hard to get that then also the modular automorphisms of the left Haar weight commute with the modular automorphisms of the right Haar weight. Indeed, fix $s, t \in \mathbb{R}$ and denote $\gamma = \sigma_t \tau_{-t}$ and $\gamma' = \sigma'_s \tau_s$. From the formulas i) and ii) in Theorem 2.3 and ii) in Theorem 2.6 we get

$$\begin{aligned}\Delta(\gamma(x)) &= (\iota \otimes \gamma)\Delta(x) \\ \Delta(\gamma'(x)) &= (\gamma' \otimes \iota)\Delta(x)\end{aligned}$$

for all x . It follows that $\Delta(\gamma\gamma'(x)) = \Delta(\gamma'\gamma(x))$ for all x . This will imply $\gamma\gamma' = \gamma'\gamma$. And as τ and σ commute, as well as τ and σ' , we can conclude from this that also σ and σ' will commute. So, we get the following result.

2.8 Theorem All the automorphism groups σ , σ' and τ mutually commute.

Now, we will show that ψ is relatively invariant w.r.t. the modular automorphism group σ and that φ is relatively invariant w.r.t. the modular automorphism group σ' , with (essentially) the same scaling factor ν . Observe that also this result would imply that σ and σ' commute. However, we will use that σ and σ' commute to prove the following theorem.

2.9 Theorem We have $\psi \circ \sigma_t = \nu^{-t}\psi$ and $\varphi \circ \sigma'_t = \nu^t\varphi$.

Proof: Take $x \in \mathcal{M}_\psi$ and consider the formula $\Delta(\sigma_t(x)) = (\tau_t \otimes \sigma_t)\Delta(x)$ (Theorem 2.3.i). If we also have that $\sigma_t(x) \in \mathcal{M}_\psi$, we can use $\psi \circ \tau_t = \nu^{-t}\psi$ (Theorem 2.7), and we find that $\psi(\sigma_t(x)) = \nu^{-t}\psi(x)$ by invariance. Because the weights $\psi \circ \sigma_t$ and $\nu^{-t}\psi$ have the same modular automorphism group (because σ and σ' commute), it will follow (from Proposition 3.16 of Chapter VIII in [T3]) that these weights are the same if we can show that the *-subalgebra $\mathcal{M}_\psi \cap \sigma_{-t}(\mathcal{M}_\psi)$ is dense. This is what we will do now.

Take any $x \in \mathcal{N}_\psi$ and take a function f on \mathbb{R} as in the proof of Proposition 2.4 to define $y = \int f(s)\sigma'_s(x)ds$. We know that y is still in \mathcal{N}_ψ and analytic with respect to σ' . Because σ commutes with σ' , also $\sigma_{-t}(y)$ will be analytic and so, as before, $\mathcal{N}_\psi\sigma_{-t}(y) \subseteq \mathcal{N}_\psi$. On the other hand, $\mathcal{N}_\psi\sigma_{-t}(y) \subseteq \sigma_{-t}(\mathcal{N}_\psi)$ because $y \in \mathcal{N}_\psi$ and \mathcal{N}_ψ is a left ideal. So, we have produced elements in the intersection of \mathcal{N}_ψ and $\sigma_{-t}(\mathcal{N}_\psi)$. It is not hard to conclude that this intersection is dense, as well as the intersection $\mathcal{M}_\psi \cap \sigma_{-t}(\mathcal{M}_\psi)$. This completes the proof of the first part of the theorem.

The second formula is obtained from the first one by using i) of Theorem 2.6.

Remark that the proof of this result is different from the original one. In [K-V2], first a stronger form of right invariance for ψ is needed.

Now we can add one more relation of the type proven in Theorem 2.3 and Theorem 2.6:

2.10 Theorem For all $x \in M$ we have $\Delta(\tau_t(x)) = (\sigma_t \otimes \sigma'_{-t})\Delta(x)$.

Proof: Because of the relative invariance of φ under both τ_t and σ'_t , we have one-parameter groups of unitaries (v_t) and (w_t) on \mathcal{H}_φ satisfying

$$\begin{aligned} v_t\Lambda_\varphi(x) &= \nu^{\frac{1}{2}t}\Lambda_\varphi(\tau_t(x)) \\ w_t\Lambda_\varphi(x) &= \nu^{-\frac{1}{2}t}\Lambda_\varphi(\sigma'_t(x)) \end{aligned}$$

for all $x \in \mathcal{N}_\varphi$ and all t . When W is the left regular representation as before, it follows from the equations $\Delta \circ \tau_t = (\tau_t \otimes \tau_t)\Delta$ and $\Delta \circ \sigma'_t = (\sigma'_t \otimes \tau_{-t})\Delta$ that

$$\begin{aligned} (1 \otimes v_t^*)W^*(1 \otimes v_t) &= (\tau_t \otimes \iota)W^* \\ (1 \otimes v_t)W^*(1 \otimes w_t) &= (\sigma'_t \otimes \iota)W^*. \end{aligned}$$

If we apply $R \otimes J(\cdot)^*J$ to the second equation, we get because J commutes with w_t and v_t that

$$(1 \otimes w_t^*)W^*(1 \otimes v_t^*) = (\sigma_{-t} \otimes \iota)W^*.$$

If we combine all of this with the formula $(\Delta \otimes \iota)W = W_{13}W_{23}$ as we did before, we can conclude that $\Delta \circ \tau_t = (\sigma_t \otimes \sigma'_{-t})\Delta$ on the left leg of W . As this leg is dense, we get the result.

The unitary groups (v_t) and (w_t) , defined in the proof of this theorem, will play an important role further. This will be seen in Section 4 where we will recall the definitions.

From the fact that ψ is relatively invariant under the modular automorphisms of φ , we also get the following important result.

2.11 Theorem There exists a unique, non-singular, positive self-adjoint operator δ , affiliated with M such that $\psi = \varphi(\delta^{\frac{1}{2}} \cdot \delta^{\frac{1}{2}})$. This operator satisfies $\sigma_t(\delta) = \nu^t \delta$ and $\sigma'_t(\delta) = \nu^{-t} \delta$. It is invariant under the automorphisms (τ_t) and $R(\delta) = \delta^{-1}$. We also have the relation $\sigma'_t(x) = \delta^{it} \sigma_t(x) \delta^{-it}$.

First remark that we have formulated the above results in terms of the unbounded operator δ . It is quite obvious what is meant by this for all the formulas in the formulation, except for the first one. This will have to be interpreted with the use of the Connes' cocycle Radon Nikodym derivative as we will see in the proof. We refer to Corollary 3.6 in Section VIII of [T3] and to [V1] for more details. In fact, when possible, we will rather consider the equivalent formulas in terms of the unitary operators (δ^{it}) in order to avoid this kind of (technical) difficulties.

Proof: Consider the weights φ and $\psi = \varphi \circ R$ and consider the cocycle Radon Nikodym derivative. We will write $u_t = (D\psi : D\varphi)_t$ for all t . For any automorphism α of M , we have

$$(D\psi \circ \alpha : D\varphi \circ \alpha)_t = \alpha^{-1}(D\psi : D\varphi)_t$$

and if we apply this with $\alpha = \sigma_s$ and if we use that $\varphi \circ \sigma_s = \varphi$ and $\psi \circ \sigma_s = \nu^{-s} \psi$, we get $\sigma_s(u_t) = \nu^{ist} u_t$ for all t and all s . Because $u_{s+t} = u_s \sigma_s(u_t)$ we easily calculate that $(\nu^{-\frac{1}{2}it^2} u_t)$ is a one-parameter group of unitaries in M . Therefore, there exists a non-singular positive self-adjoint operator δ , affiliated with M , such that $u_t = \nu^{\frac{1}{2}it^2} \delta^{it}$ for all t . This gives us the element δ such that, at least formally, $\psi = \varphi(\delta^{\frac{1}{2}} \cdot \delta^{\frac{1}{2}})$. Takesaki (Chapter VIII in [T3]) treats this situation in the case $\nu = 1$ whereas Vaes [V1] considers the general case.

Because ψ and φ are scaled with the same factor by the automorphisms τ_s we must have that u_t and hence also δ is invariant under τ_s . Because $\psi = \varphi \circ R$ and $\varphi \circ R = \psi$ we will get that $R(u_t) = u_{-t}$ and so $R(\delta) = \delta^{-1}$. Because $\sigma_t(u_s) = \nu^{ist} u_s$ we obtain $\sigma_t(\delta) = \nu^t \delta$ and finally, if we apply R to this last formula, we get the other one $\sigma'_t(\delta) = \nu^{-t} \delta$.

The last statement is standard.

One also has the formula $\Delta(\delta) = \delta \otimes \delta$ but this is not so easy to get. One possible way to prove this formula can be found in the original work [K-V2]. Another proof is found

in [M-N-W]. We will still give an other argument but for this, we need some more results and so we postpone the proof of this formula (see Remark 4.16 in Section 4).

From now on, we will identify \mathcal{H}_ψ with \mathcal{H}_φ in the obvious way, namely by defining Λ_ψ from \mathcal{N}_ψ to \mathcal{H}_φ by $\Lambda_\psi(x) = \Lambda_\varphi(x\delta^{\frac{1}{2}})$ when $x \in \mathcal{N}_\psi$. This is possible because of the above theorem. The element $x\delta^{\frac{1}{2}}$ is considered to be the closure of the operator, first defined on the domain of $\delta^{\frac{1}{2}}$. This will be bounded and in \mathcal{N}_ψ for $x \in \mathcal{N}_\varphi$. Finally, observe that this identification is compatible with the actions of M on both spaces. So, we are not creating any conflict with these conventions.

3. The dual $(\hat{M}, \hat{\Delta})$

In this section, *we start with a locally compact quantum group (M, Δ)* as in Definition 2.1 and we will construct the dual $(\hat{M}, \hat{\Delta})$. We will also show that, repeating the procedure, gives the original pair (M, Δ) .

We take a left invariant Haar weight φ and the right Haar weight ψ obtained from it by composing with the unitary antipode R (see a remark before Theorem 2.6 in the previous section). We will, as explained also at the end of the previous section, identify \mathcal{H}_ψ with \mathcal{H}_φ by defining Λ_ψ from \mathcal{N}_ψ to \mathcal{H}_φ by $\Lambda_\psi(x) = \Lambda_\varphi(x\delta^{\frac{1}{2}})$ when $x \in \mathcal{N}_\psi$. Observe again that this is compatible with the actions of M on both spaces. Moreover, because we have uniqueness of the Haar weights, it is no longer necessary to use the subscript φ when considering the Hilbert space. So, from now on, we will simply use \mathcal{H} for \mathcal{H}_φ . In other words, we will be working solely within the Hilbert space \mathcal{H}_φ and denote it by \mathcal{H} .

We consider the left regular representation W of (M, Δ) associated with the left Haar weight as in Proposition 1.11. With the above conventions, we get that W acts on $\mathcal{H} \otimes \mathcal{H}$ and that $W \in M \otimes \mathcal{B}(\mathcal{H})$. Then also, W satisfies the pentagon equation $W_{12}W_{13}W_{23} = W_{23}W_{12}$.

We will make use of the antipode S and its polar decomposition $S = R\tau_{-\frac{i}{2}}$. We will also use that the space of elements of the form $(\iota \otimes \omega)W$ with $\omega \in \mathcal{B}(\mathcal{H})_*$ is invariant under the scaling group (τ_t) , that it is a core for S and that $S((\iota \otimes \omega)W) = (\iota \otimes \omega)(W^*)$. We will refer to this property when we write formally $(S \otimes \iota)W = W^*$. For details, see Section 1, Proposition 1.24.

The dual von Neumann algebra and the dual coproduct

First the underlying von Neumann algebra \hat{M} is defined.

3.1 Definition Let \hat{M} be the σ -weak closure of the subspace $\{(\omega \otimes \iota)W \mid \omega \in M_*\}$ of $\mathcal{B}(\mathcal{H})$.

With this definition, the dual von Neumann algebra \hat{M} also acts on the space \mathcal{H} (which is \mathcal{H}_φ). In some sense, this means that, from the very beginning, we identify the 'spaces' $L^2(G)$ and $L^2(\hat{G})$, thus not using the 'Fourier transform' explicitly. We come back to this remark later in this section when we construct the dual left Haar weight.

It follows from the fact that W is a multiplicative unitary, that the subspace C of $\mathcal{B}(\mathcal{H})$, defined as $\{(\omega \otimes \iota)W \mid \omega \in M_*\}$, is a subalgebra of $\mathcal{B}(\mathcal{H})$. In order to have that its closure is a von Neumann algebra, we will use the following result (which will also be needed further in this section).

3.2 Lemma Consider the one-parameter group of automorphisms (τ_t) on M and its dual on M_* . If $\omega \in M_*$ is analytic and if ω_1 is defined as $\bar{\omega} \circ \tau_{-\frac{i}{2}} \circ R$, then

$$((\omega \otimes \iota)W)^* = (\omega_1 \otimes \iota)W.$$

Conversely, if $\omega, \omega_1 \in M_*$ and satisfy the above equation, then $\omega_1(x) = \omega(S(x)^*)^- = \bar{\omega}(S(x))$ for all $x \in \mathcal{D}(S)$.

Proof: Recall that formally, $(S \otimes \iota)W = W^*$ and that $S = R\tau_{-\frac{i}{2}}$. Now, it is not so hard to show rigorously that the first statement of the lemma is correct.

Conversely, let $\omega, \omega_1 \in M_*$ and assume that $((\omega \otimes \iota)W)^* = (\omega_1 \otimes \iota)W$. If $x = (\iota \otimes \rho)W$ with $\rho \in \mathcal{B}(\mathcal{H})_*$, then

$$\begin{aligned} \omega_1(x) &= \rho((\omega_1 \otimes \iota)W) \\ &= \bar{\rho}((\omega \otimes \iota)W)^- \\ &= \omega((\iota \otimes \bar{\rho})W)^- \\ &= \omega(S(x)^*)^-. \end{aligned}$$

Now, because the 'left leg' of W is a core for S , we will also get the desired formula for all $x \in \mathcal{D}(S)$.

Now the following (standard) result can be shown.

3.3 Proposition \hat{M} is a von Neumann algebra and $W \in M \otimes \hat{M}$.

Proof: Again consider the subspace C defined by $\{(\omega \otimes \iota)W \mid \omega \in M_*\}$ (without taking the closure). By the first statement of the lemma, and using that such analytic elements are dense, we get that the closure \hat{M} is self-adjoint. It acts non-degenerately on \mathcal{H} because W is a unitary. Therefore, we have that \hat{M} will be a von Neumann algebra on \mathcal{H} .

The second statement follows from the commutation theorem for tensor products of von Neumann algebras.

We see from the above that also the norm closure of the space C is a C^* -algebra, acting non-degenerately on the Hilbert space \mathcal{H} . It is denoted by \hat{A} and it will be considered

further in the Appendix where we treat the relation between the von Neumann algebra approach and the C*-algebra approach.

Remark that it is not immediate that these subalgebras \hat{A} and \hat{M} are self-adjoint. Usually, either regularity (cf. [B-S]) or manageability (cf. [W2]) of the multiplicative unitary W is used. Here we use a result which is closely related to manageability, but (in some sense) also weaker than manageability. As explained already in the introduction, we avoid the use of manageability.

Finally, we also want to mention the following. Because $\Delta(x) = W(1 \otimes x)W^*$ for $x \in M$, it follows from Proposition 2.4 that $M \cap \hat{M}' = \mathbb{C}1$. Indeed, if $x \in M \cap \hat{M}'$, then $W(1 \otimes x) = (1 \otimes x)W$ because $x \in \hat{M}'$ and so $\Delta(x) = 1 \otimes x$. By Proposition 2.4 we get $x \in \mathbb{C}1$.

Now, we proceed to define the coproduct $\hat{\Delta}$ on \hat{M} . We first formulate the following result.

3.4 Lemma We have $W(y \otimes 1)W^* \in \hat{M} \otimes \hat{M}$ for all $y \in \hat{M}$.

This result is standard and easy to prove. It follows from the definition of \hat{M} , the formula $W_{23}W_{12}W_{23}^* = W_{12}W_{13}$ and the fact that $W \in M \otimes \hat{M}$.

Now, it is easy to define the coproduct $\hat{\Delta}$. As is common in the theory of locally compact quantum groups, we use the flip in the following definition.

3.5 Definition We let $\hat{\Delta}(y) = \chi(W(y \otimes 1)W^*)$ for $y \in \hat{M}$ where χ is the flip on the tensor product $\hat{M} \otimes \hat{M}$.

It is clear, using the previous lemma, that $\hat{\Delta}$ is a unital and normal *-homomorphism from \hat{M} to $\hat{M} \otimes \hat{M}$. The coassociativity follows in the standard way from the pentagon equation.

Construction of the left Haar weight $\hat{\varphi}$ on $(\hat{M}, \hat{\Delta})$

The next step is the construction of the Haar weights on the dual $(\hat{M}, \hat{\Delta})$. We will first construct the left Haar weight $\hat{\varphi}$ and later discuss the existence of the right Haar weight $\hat{\psi}$. We will follow the (more or less) standard procedure.

We begin with the construction of the map $\hat{\Lambda}$ (which later will be closed and give the map $\Lambda_{\hat{\varphi}}$ associated with the dual left Haar weight $\hat{\varphi}$). The definition looks somewhat strange although it is well-known in the theory of Kac algebras. Let us give here a short motivation using the algebraic theory of multiplier Hopf algebras with integrals [VD3] (for more details about this motivation, we refer to [VD8]).

In [VD3], the Fourier transform \hat{a} of an element a is defined as the linear functional $\omega = \varphi(\cdot a)$. Now, it is well-known that the multiplicative unitary W is essentially the duality (see e.g. [VD1] for the Hopf algebra case and [D-VD1] for the case of algebraic quantum groups). So, formally, we have $\hat{a} = (\omega \otimes \iota)W$ when $\omega = \varphi(\cdot a)$. As we remarked already before, the 'spaces' $L^2(G)$ and $L^2(\hat{G})$ are identified which means (again formally) that we want $\Lambda_{\hat{\varphi}}(\hat{a}) = \Lambda_{\varphi}(a)$. This formula is rewritten as

$$\langle \Lambda_{\hat{\varphi}}(\hat{a}), \Lambda_{\varphi}(x) \rangle = \langle \Lambda_{\varphi}(a), \Lambda_{\varphi}(x) \rangle = \varphi(x^*a) = \omega(x^*)$$

whenever $x \in \mathcal{N}_\varphi$.

So, the following definition is not a surprise.

3.6 Definition Define a subspace $\hat{\mathcal{N}}$ of \hat{M} and a linear map $\hat{\Lambda} : \hat{\mathcal{N}} \rightarrow \mathcal{H}$ as follows. We say that an element y of \hat{M} belongs to $\hat{\mathcal{N}}$ if there exists a linear functional $\omega \in M_*$ such that $y = (\omega \otimes \iota)W$ and a vector $\xi \in \mathcal{H}$ such that $\langle \xi, \Lambda_\varphi(x) \rangle = \omega(x^*)$ for all $x \in \mathcal{N}_\varphi$. Then we set $\hat{\Lambda}(y) = \xi$.

As before, Λ_φ is the canonical map from \mathcal{N}_φ to \mathcal{H} . Remark that ω is uniquely determined by y because the left leg of W is dense in M (cf. Proposition 1.21) and that ξ is completely determined by ω because $\Lambda_\varphi(\mathcal{N}_\varphi)$ is dense in \mathcal{H} . It is also clear that $\hat{\mathcal{N}}$ is a subspace and that $\hat{\Lambda}$ is linear and injective.

In the next lemma, we will show that there are enough elements in this space $\hat{\mathcal{N}}$. We will need *right bounded* vectors in \mathcal{H} . These are elements $\eta \in \mathcal{H}$ such that there exist a bounded operator, denoted by $\pi'(\eta)$, satisfying $\pi'(\eta)\Lambda_\varphi(x) = x\eta$ for all $x \in \mathcal{N}_\varphi$. Such vectors form a dense subspace and the space of operators $\pi'(\eta)$ with η right bounded, is dense in the commutant M' of M (see e.g. Chapter VI in [T3]).

3.7 Lemma Let $\xi, \eta \in \mathcal{H}$ and assume that η is right bounded. Let $\omega = \langle \cdot, \eta \rangle$ and $y = (\omega \otimes \iota)W$. Then $y \in \hat{\mathcal{N}}$ and $\hat{\Lambda}(y) = \pi'(\eta)^*\xi$. In particular, the space $\hat{\Lambda}(\hat{\mathcal{N}})$ is dense in \mathcal{H} and also $\hat{\mathcal{N}}$ is σ -weakly dense in \hat{M} .

The proof is easy and straightforward.

We also have the following.

3.8 Lemma Let $\omega, \omega_1 \in M_*$ and $y = (\omega \otimes \iota)W$ and $y_1 = (\omega_1 \otimes \iota)W$. If $y \in \hat{\mathcal{N}}$, then also $y_1 y \in \hat{\mathcal{N}}$ and $\hat{\Lambda}(y_1 y) = y_1 \hat{\Lambda}(y)$.

Proof: For any $x \in \mathcal{N}_\varphi$ we have

$$\begin{aligned}
\langle y_1 \hat{\Lambda}(y), \Lambda_\varphi(x) \rangle &= \langle \hat{\Lambda}(y), y_1^* \Lambda_\varphi(x) \rangle \\
&= \langle \hat{\Lambda}(y), ((\overline{\omega_1} \otimes \iota)(W^*)) \Lambda_\varphi(x) \rangle \\
&= \langle \hat{\Lambda}(y), \Lambda_\varphi((\overline{\omega_1} \otimes \iota)\Delta(x)) \rangle \\
&= \omega((\overline{\omega_1} \otimes \iota)\Delta(x))^* \\
&= \omega((\omega_1 \otimes \iota)\Delta(x^*)) \\
&= (\omega_1 \omega)(x^*)
\end{aligned}$$

where $\omega_1 \omega$ is defined as usual by $(\omega_1 \omega)(a) = (\omega_1 \otimes \omega)\Delta(a)$ for all $a \in M$. Remark that we have used the definition of W^* as given in Proposition 1.11. Finally, it is easy to see that $y_1 y = ((\omega_1 \omega) \otimes \iota)W$ using the pentagon equation and the formula $\Delta(a) = W^*(1 \otimes a)W$ for $a \in M$.

The left Haar weight $\hat{\varphi}$ will be obtained by constructing a left Hilbert algebra. Definition 3.6 and the Lemmas 3.7 and 3.8 clearly provide the first step. We need one more lemma before we can come to the main part of the construction.

3.9 Lemma Consider the set $\hat{\mathcal{N}}_0$ of elements $y \in \hat{\mathcal{N}}$ so that y^* has the form $(\omega_1 \otimes \iota)W$ for some $\omega_1 \in M_*$. Then $\hat{\mathcal{N}}_0$ is still σ -weakly dense in \hat{M} and also $\hat{\Lambda}(\hat{\mathcal{N}}_0)$ is still dense in \mathcal{H} .

Proof: Take $\omega = \langle \cdot, \xi, \eta \rangle$ with $\xi, \eta \in \mathcal{H}$ and assume that η is right bounded. We know from Lemma 3.7 that y , defined as $(\omega \otimes \iota)W$, is in $\hat{\mathcal{N}}$ and that $\hat{\Lambda}(y) = \pi'(\eta)^*\xi$. Now, because of Lemma 3.2, we need such elements with ω analytic with respect to (τ_t) .

To construct such elements, define a one-parameter group of unitaries (v_t) on \mathcal{H} (as in the proof of Theorem 2.10) by $v_t \Lambda_\varphi(x) = \nu^{\frac{1}{2}t} \Lambda_\varphi(\tau_t(x))$ when $x \in \mathcal{N}_\varphi$. It is clear that $\tau_t(x) = v_t x v_t^*$ for all $x \in M$. It will be possible to take the vector ξ above so that it is analytic with respect to (v_t) . And because every v_t will map right bounded elements to right bounded elements, we will also be able to take η analytic and still right bounded. Then ω will be analytic and of the required form. This will give us the result of the lemma.

Now, we come to the main step in the construction of our left Hilbert algebra.

3.10 Proposition Let $\mathfrak{A} = \hat{\Lambda}(\hat{\mathcal{N}} \cup \hat{\mathcal{N}}^*)$. We can equip \mathfrak{A} with the $*$ -algebra structure inherited from $\hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$. If we denote y by $\pi(\xi)$ when $y \in \hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$ and $\xi = \hat{\Lambda}(y)$, then we have

- i) both \mathfrak{A} and \mathfrak{A}^2 are dense in \mathcal{H} ,
- ii) $\pi(\xi)$ is continuous for all $\xi \in \mathfrak{A}$,
- iii) π is a $*$ -representation of \mathfrak{A} .

Proof: It follows from the definition of $\hat{\mathcal{N}}$ and Lemma 3.8 that $\hat{\mathcal{N}}$ is a subalgebra of \hat{M} and so $\hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$ is a $*$ -subalgebra of \hat{M} . Because $\hat{\Lambda}$ is injective, we can equip \mathfrak{A} with the $*$ -algebra structure of $\hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$.

To prove i), we use the previous lemma. Indeed, if we take y, y_1 in the set $\hat{\mathcal{N}}_0$, we see that both y^*y_1 and y_1^*y will be in $\hat{\mathcal{N}}$ and this will provide us with enough elements in $\hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$. And because this set is dense in M_* and also $\hat{\Lambda}(\hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*)$ is dense, we get i).

Statements ii) and iii) are immediate consequences of the notations.

There is now only one point missing for \mathfrak{A} to be a left Hilbert algebra. It is also needed that the $*$ -operation in \mathfrak{A} , usually denoted by $\xi \mapsto \xi^\sharp$, is preclosed. This will be a consequence of the following lemma.

3.11 Lemma If $y \in \hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$ then

$$\langle \hat{\Lambda}(y^*), \Lambda_\varphi(a) \rangle = \langle \hat{\Lambda}(y), \Lambda_\varphi(S(a^*)) \rangle^-$$

whenever $a \in \mathcal{N}_\varphi$, $a^* \in \mathcal{D}(S)$ and $S(a^*) \in \mathcal{N}_\varphi$.

Proof: Take $y \in \hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$ and let ω, ω_1 be in M_* so that $y = (\omega \otimes \iota)W$ and $y^* = (\omega_1 \otimes \iota)W$. From Lemma 3.2 we know that $\omega_1(x) = \omega(S(x)^*)^-$ for all $x \in \mathcal{D}(S)$. If now a is as in the formulation of the lemma, it will follow from the definition of $\hat{\Lambda}$ that

$$\begin{aligned} \langle \hat{\Lambda}(y^*), \Lambda_\varphi(a) \rangle &= \omega_1(a^*) = \omega(S(a^*)^*)^- \\ &= \langle \hat{\Lambda}(y), \Lambda_\varphi(S(a^*)) \rangle^- \end{aligned}$$

and the result will follow.

So, in order to prove that the involution in \mathfrak{A} is preclosed, we just need to argue that there are enough elements a as in the lemma. This is the content of the next lemma.

3.12 Lemma The set of elements $\Lambda_\varphi(a)$ with $a \in \mathcal{N}_\varphi$ such that also $a^* \in \mathcal{D}(S)$ and $S(a^*) \in \mathcal{N}_\varphi$ is dense in \mathcal{H} .

Proof: Take $a \in \mathcal{N}_\varphi$. Formally, we have

$$\begin{aligned} S(a^*) &= R\tau_{-\frac{i}{2}}(a^*) = R(\tau_{\frac{i}{2}}(a))^* \\ &= R(\tau_{\frac{i}{2}}(a)\delta^{\frac{1}{2}})^*\delta^{\frac{1}{2}}. \end{aligned}$$

For such an element to be again in \mathcal{N}_φ , we need that $\tau_{\frac{i}{2}}(a)\delta^{\frac{1}{2}}$ is well-defined and in \mathcal{N}_φ . We need two results to obtain such elements. First observe that φ is relatively invariant w.r.t. the automorphisms (τ_t) and so also \mathcal{N}_φ is invariant and standard techniques allow to produce elements $a \in \mathcal{N}_\varphi$ that are analytic w.r.t. (τ_t) . Next we know that the elements δ^{is} are analytic w.r.t. the modular automorphisms (σ_t) and so $\mathcal{N}_\varphi\delta^{is} \subseteq \mathcal{N}_\varphi$ for all s . This will allow us to produce elements $a \in \mathcal{N}_\varphi$ such that $a\delta^{\frac{1}{2}}$ is well-defined and still in \mathcal{N}_φ . The two techniques together will give us enough of the desired elements.

We know from Section 1 that the operator K is essentially the map $\Lambda_\psi(x) \mapsto \Lambda_\psi(S(x)^*)$. A simple (formal) argument then shows that the map $\Lambda_\varphi(x) \mapsto \Lambda_\varphi(S(x^*))$ is essentially the operator K^* . So, we expect that the map $\hat{\Lambda}(y) \mapsto \hat{\Lambda}(y^*)$ will be nothing else but the operator K . It does not seem to be easy to prove this result exactly. In the next section, we will find a way around this problem by (in some sense) 'redefining' these maps (see Remark 4.10). We will then also take up again the argument that we sketch here in the proof of Lemma 3.12 (see the proof of Theorem 4.11).

For the moment, there is no need to get this more precise result. Indeed, if we combine all the previous results, we find that \mathfrak{A} is a left Hilbert algebra and this is what we need. Then we can use the general procedure to construct a faithful normal semi-finite weight from a left Hilbert algebra (see Chapter VII in [T3]) and we will arrive at the following theorem.

3.13 Theorem There exists a normal faithful semi-finite weight $\hat{\varphi}$ on \hat{M} such that the G.N.S.-representation can be realized in \mathcal{H} , satisfying $\hat{\mathcal{N}} \subseteq \mathcal{N}_{\hat{\varphi}}$ and such that the canonical map $\Lambda_{\hat{\varphi}}$ is the closure of $\hat{\Lambda}$ on $\hat{\mathcal{N}}$.

Proof: We have the left Hilbert algebra $\hat{\Lambda}(\hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*)$ sitting inside \mathcal{H} . The canonical weight $\hat{\varphi}$ associated to this left Hilbert algebra has the property that elements in $\hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$ belong to $\mathcal{N}_{\hat{\varphi}}$ and that $\hat{\varphi}(y^*y) = \langle \hat{\Lambda}(y), \hat{\Lambda}(y) \rangle$ for such elements. The rest follows from standard Hilbert algebra theory and the construction method of the associated weight.

The left invariance of $\hat{\varphi}$ and the associated regular representation

Now, we need to show that $\hat{\varphi}$ is left invariant on the pair $(\hat{M}, \hat{\Delta})$. When this is done, we can easily construct the right Haar weight on $(\hat{M}, \hat{\Delta})$. Indeed, as in the proof of Theorem 2.3, we will have that $\hat{\Delta}(\hat{R}(y)) = \chi(\hat{R} \otimes \hat{R})\hat{\Delta}(y)$ whenever $y \in \hat{M}$ where as before χ denotes the flip and where here \hat{R} is defined on \hat{M} by $\hat{R}(y) = Jy^*J$. Recall that J is the modular conjugation associated with φ as defined in Notation 1.19. Therefore, a right Haar weight can be constructed from the left Haar weight on \hat{M} by composing it with this map \hat{R} . We will show later, in Section 4, that the use of the notation \hat{R} is justified (see Proposition 4.12).

This will give us that the pair $(\hat{M}, \hat{\Delta})$ is again a locally compact quantum group. In order to show that repeating the procedure will bring us back to the original locally compact quantum group (M, Δ) , we will prove that the left regular representation of the dual is nothing else but the unitary $\Sigma W^* \Sigma$ where as is common, Σ denotes the flip operator on the tensor product $\mathcal{H} \otimes \mathcal{H}$.

We will prove this result first because it can be used to show that the dual left Haar weight $\hat{\varphi}$ is indeed left invariant. In other words, the main result left to prove is the following.

3.14 Proposition Define the unitary $\hat{W} = \Sigma W^* \Sigma$ on $\mathcal{H} \otimes \mathcal{H}$. Then $(\omega \otimes \iota)\hat{\Delta}(y) \in \mathcal{N}_{\hat{\varphi}}$ and

$$((\omega \otimes \iota)\hat{W}^*)\Lambda_{\hat{\varphi}}(y) = \Lambda_{\hat{\varphi}}((\omega \otimes \iota)\hat{\Delta}(y))$$

whenever $y \in \mathcal{N}_{\hat{\varphi}}$ and $\omega \in \mathcal{B}(\mathcal{H})_*$.

Proof: First take $y \in \hat{\mathcal{N}}$ and let $\omega \in M_*$ be such that $y = (\omega \otimes \iota)W$. Take any $\rho \in \hat{M}_*$ and put $y_1 = (\rho \otimes \iota)\hat{\Delta}(y)$. Since $\hat{\Delta}(y) = \chi(W(y \otimes 1)W^*)$ (cf. Definition 3.5), it follows from a straightforward calculation (using the pentagon equation) that $y_1 = (\omega(\cdot c) \otimes \iota)W$ where $c = (\iota \otimes \rho)W$. Then, when $a \in \mathcal{N}_{\hat{\varphi}}$, we have

$$\begin{aligned} \omega(a^*c) &= \langle \hat{\Lambda}(y), c^* \Lambda_{\varphi}(a) \rangle \\ &= \langle c \hat{\Lambda}(y), \Lambda_{\varphi}(a) \rangle \end{aligned}$$

and it follows that $y_1 \in \hat{\mathcal{N}}$ and that $\hat{\Lambda}(y_1) = c \hat{\Lambda}(y)$. This precisely means that

$$\begin{aligned} \hat{\Lambda}((\rho \otimes \iota)\hat{\Delta}(y)) &= ((\iota \otimes \rho)W)\hat{\Lambda}(y) \\ &= ((\rho \otimes \iota)\hat{W}^*)\hat{\Lambda}(y). \end{aligned}$$

This proves the result when $y \in \hat{\mathcal{N}}$. The general case follows because $\Lambda_{\hat{\varphi}}$ on $\mathcal{N}_{\hat{\varphi}}$ is the closure of $\hat{\Lambda}$ on $\hat{\mathcal{N}}$.

Now, it is not hard to prove left invariance of $\hat{\varphi}$.

3.15 Proposition The weight $\hat{\varphi}$, as constructed in Proposition 3.13 is left invariant on $(\hat{M}, \hat{\Delta})$.

Proof: Take $y \in \mathcal{N}_{\hat{\varphi}}$. Take also a vector $\xi \in \mathcal{H}$ and let $\omega = \langle \cdot, \xi, \xi \rangle$. Consider an orthonormal basis (ξ_i) in \mathcal{H} . Then we have

$$(\omega \otimes \iota)\hat{\Delta}(y^*y) = \sum y_i^*y_i$$

where $y_i = (\langle \cdot, \xi, \xi_i \rangle \otimes \iota)\hat{\Delta}(y)$. We know from the previous proposition that $y_i \in \mathcal{N}_{\hat{\varphi}}$ and that $\Lambda_{\hat{\varphi}}(y_i) = z_i\Lambda_{\hat{\varphi}}(y)$ where $z_i = (\langle \cdot, \xi, \xi_i \rangle \otimes \iota)\hat{W}^*$. And because

$$\sum z_i^*z_i = (\langle \cdot, \xi, \xi \rangle \otimes \iota)(\hat{W}\hat{W}^*) = \omega(1)1,$$

it follows that

$$\begin{aligned} \hat{\varphi}((\omega \otimes \iota)\hat{\Delta}(y^*y)) &= \sum \hat{\varphi}(y_i^*y_i) \\ &= \sum \langle z_i^*z_i\Lambda_{\hat{\varphi}}(y), \Lambda_{\hat{\varphi}}(y) \rangle \\ &= \omega(1)\hat{\varphi}(y^*y). \end{aligned}$$

This proves invariance.

Observe that the invariance is proven by first constructing the candidate for the left regular representation and using that this is a unitary. This is a standard technique (e.g. see the construction of the Haar weight in [VD5]).

Now we are almost ready for the *main result*. We just need that there is also a right invariant Haar weight. This will follow from the next proposition (a result that was already announced earlier).

3.16 Proposition Define \hat{R} on \hat{M} by $\hat{R}(y) = Jy^*J$, where as before, J is the modular conjugation associated with the weight φ on M (cf. Notation 1.19). Then \hat{R} is an involutive *-anti-automorphism of \hat{M} that flips the coproduct $\hat{\Delta}$.

This all essentially follows from the formula i) in Proposition 1.20 (compare also with the proof of iii) in Theorem 2.3).

As an immediate consequence, we get that the weight $\hat{\psi}$, defined as the composition of $\hat{\varphi}$ with \hat{R} , will be a right invariant weight. Therefore, we have completed the proof of the following, main result.

3.17 Theorem The pair $(\hat{M}, \hat{\Delta})$ (as constructed in 3.1 and 3.5), is a locally compact quantum group (in the sense of Definition 2.1).

In the next section we will argue that the involutive $*$ -anti-automorphism \hat{R} is indeed the unitary antipode on the dual and that this notation is consistent. In fact, we will obtain more formulas in the next section relating the objects of the original quantum group (M, Δ) with those of the dual $(\hat{M}, \hat{\Delta})$.

The bidual

We finish this section with some remark about biduality. We have the following result.

3.18 Theorem The dual of $(\hat{M}, \hat{\Delta})$ is again (M, Δ) .

This follows from the fact that the regular representation \hat{W} of the dual coincides with $\Sigma W^* \Sigma$ (cf. Proposition 3.14).

Let us now make a comparison with the theory of (multiplier) Hopf algebras. There, the dual is usually equipped with the coproduct, dual to the product and not with the opposite coproduct as we have done here. It is obvious that in that case, the dual of the dual is the original algebra with the original coproduct. If, as is done here in the operator algebra approach, the dual is equipped with the opposite coproduct, then one might expect that the dual of the dual will yield the original algebra but with both the opposite product and the opposite coproduct. That this is not seen here is simply a result of the treatment. There is no problem as the unitary antipode is a map that converts the product to the opposite product and the coproduct to the opposite coproduct.

4. A collection of formulas

In this section, we collect many of the *formulas relating the various objects* associated with a locally compact quantum group and its dual. We will not give all the possible relations (as there are many), but the most important ones. Other equalities can easily be obtained from the ones that we prove. Again, it should be mentioned that we do not get really new results but that some of the results are proven in a slightly other fashion than in the original papers by Kustermans and Vaes. Also, the formulas are organized in another (perhaps more systematic) way.

Fix a locally compact quantum group (M, Δ) and consider the dual $(\hat{M}, \hat{\Delta})$ as constructed in the previous section. In this section, we will freely use the definitions and notations of the previous sections. When appropriate, we will explicitly recall the necessary notions and results.

We will do so with the following definition and notations (some of them introduced already in the proof of Theorem 2.10).

4.1 Definition Define continuous one-parameter groups of unitaries (u_t) , (v_t) and (w_t) on \mathcal{H} by

$$\begin{aligned} u_t \Lambda_\varphi(x) &= \Lambda_\varphi(\sigma_t(x)) \\ v_t \Lambda_\varphi(x) &= \nu^{\frac{1}{2}t} \Lambda_\varphi(\tau_t(x)) \\ w_t \Lambda_\varphi(x) &= \nu^{-\frac{1}{2}t} \Lambda_\varphi(\sigma'_t(x)) \end{aligned}$$

when $x \in \mathcal{N}_\varphi$.

The relative invariance of φ with respect to (τ_t) and (σ'_t) respectively is used to justify the definitions of (v_t) and (w_t) . Recall from Theorem 2.7 and Theorem 2.9 that $\varphi \circ \tau_t = \nu^{-t}\varphi$ and $\varphi \circ \sigma'_t = \nu^t\varphi$ (where ν is the scaling constant). Of course, $u_t = \nabla^{it}$ for all t . We have just introduced this notation in order to have some more symmetry. In what follows, we will use either of these two notations for this one-parameter group.

4.2 Remarks i) We have that $v_t = P^{it}$ where P is the operator defined in [K-V2, Definition 6.9]. Because of the special role of this operator (see further), and because we want to be as close as possible to the notations used in the papers [K-V*], we will further in this section use P^{it} as well as v_t (whatever is more convenient), just as in the case of ∇^{it} and u_t .

ii) It can be verified that $w_t \Lambda_\psi(x) = \Lambda_\psi(\sigma'_t(x))$ for all $x \in \mathcal{N}_\psi$. Therefore, $w_t = \mathbb{V}^{it}$ where \mathbb{V} is the modular operator associated with the right Haar weight ψ on M . Also in this case, we will use \mathbb{V}^{it} as well as w_t .

In the following proposition, we formulate a first relation involving some of these operators.

4.3 Proposition We have $\mathbb{V}^{it} = \delta^{it} (J\delta^{it}J) \nabla^{it}$ for all t .

This result follows because $\sigma'_t(x) = \delta^{it}\sigma(x)\delta^{-it}$ for all $x \in M$ and $\sigma_s(\delta^{it}) = \nu^{ist}\delta^{it}$ (see Theorem 2.11). Observe that δ^{it} and $J\delta^{it}J$ commute and that also ∇^{it} commutes with the product $\delta^{it} (J\delta^{it}J)$.

From the definitions of these unitaries, and using that the automorphism groups involved mutually commute (cf. Theorem 2.8), we also get the following formulas.

4.4 Proposition i) All the unitaries (u_t) , (v_t) and (w_t) mutually commute and they all also commute with the modular conjugation J .

ii) We have

$$\begin{aligned} \sigma_t(x) &= u_t x u_t^* \\ \sigma'_t(x) &= w_t x w_t^* \\ \tau_t(x) &= v_t x v_t^* \end{aligned}$$

for all $x \in M$ and $t \in \mathbb{R}$.

We will now introduce some more operators of the same type as in Definition 4.1. Later, we will justify the notations used in this definition.

4.5 Definition Define a conjugate linear, involutive operator \hat{J} and a non-singular, positive self-adjoint operator $\hat{\nabla}$ on \mathcal{H} by

$$\begin{aligned}\hat{J}\Lambda_\varphi(x) &= \Lambda_\varphi(R(x)^*\delta^{\frac{1}{2}}) \\ \hat{\nabla}^{it}\Lambda_\varphi(x) &= \Lambda_\varphi(\tau_t(x)\delta^{-it})\end{aligned}$$

whenever $x \in \mathcal{N}_\varphi$ and $t \in \mathbb{R}$.

Observe that $R(x)^* \in \mathcal{N}_\psi$ when $x \in \mathcal{N}_\varphi$ because $\psi = \varphi \circ R$. Then $R(x)^*\delta^{\frac{1}{2}}$ is well-defined and belongs to \mathcal{N}_φ (as we have seen before, at the end of Section 2). Also $\tau_t(x)\delta^{-it} \in \mathcal{N}_\varphi$ whenever $x \in \mathcal{N}_\varphi$. Therefore the definitions above are possible.

To show that \hat{J} is isometric we use

$$\varphi(\delta^{\frac{1}{2}}R(x)R(x)^*\delta^{\frac{1}{2}}) = \psi(R(x^*x)) = \varphi(x^*x)$$

when $x \in \mathcal{N}_\varphi$. Because $R(\delta) = \delta^{-1}$, we will have that $\hat{J}^2 = 1$.

To show that the map $\Lambda_\varphi(x) \mapsto \Lambda_\varphi(\tau_t(x)\delta^{-it})$ is isometric and that we get indeed a one-parameter group of unitaries, we can either make a straightforward calculation or use the formula that we obtain in Proposition 4.6 below.

It is also straightforward to show that \hat{J} and $\hat{\nabla}^{it}$ commute for all $t \in \mathbb{R}$.

From the definition, we immediately get the following relation.

4.6 Proposition We have $\hat{\nabla}^{it} = J\delta^{it}J P^{it}$ for all t

This formula is an easy consequence of the definitions and again of the fact that $\sigma_s(\delta^{it}) = \nu^{ist}\delta^{it}$. Observe that also here the operators $J\delta^{it}J$ and P^{it} commute.

4.7 Proposition We have $R(x) = \hat{J}x^*\hat{J}$ and $\tau_t(x) = \hat{\nabla}^{it}x\hat{\nabla}^{-it}$ for all x and all t .

Again, these two formulas are easy consequences of the definitions. Remark that here we get another one-parameter group of unitaries that implements the scaling group τ_t (compare with Proposition 4.4).

Recall that we also had the formulas $R(x) = Ix^*I$ and $\tau_t(x) = L^{it}xL^{-it}$ (see Definition 1.22). Indeed, there are reasons to believe that we have $I = \hat{J}$ and $L = \hat{\nabla}$. We will come back to this problem later (cf. Remark 4.10).

Now, we will prove some new relations.

4.8 Proposition We have $\nabla^{it} = \hat{J}\nabla^{-it}\hat{J}$ for all t .

The result follows easily from the fact that $R(\sigma_t(x)) = \sigma'_{-t}(R(x))$ for all $x \in M$ (cf. Theorem 2.6) and $\sigma_t(\delta) = \nu^t \delta$ (Theorem 2.11).

This is one useful formula involving the operator \hat{J} . Another one is the following relation between the left regular representation W associated with φ and the right regular representation V associated with ψ on M .

4.9 Proposition We have

$$V = (\hat{J} \otimes \hat{J})\Sigma W^* \Sigma(\hat{J} \otimes \hat{J}).$$

Recall that here Σ denotes the flip on $\mathcal{H} \otimes \mathcal{H}$.

The proof is straightforward. Formally, we can write, with the Sweedler notation $\Delta(x) = x_{(1)} \otimes x_{(2)}$ (without the summation sign because we are using this symbol for something else here) and using that R flips the coproduct:

$$\begin{aligned} V(\hat{J} \otimes \hat{J})(\Lambda_\varphi(x) \otimes \xi) &= V(\Lambda_\varphi(R(x)^* \delta^{\frac{1}{2}}) \otimes \hat{J}\xi) \\ &= V(\Lambda_\psi(R(x)^*) \otimes \hat{J}\xi) \\ &= \Lambda_\psi(R(x_{(2)})^*) \otimes R(x_{(1)})^* \hat{J}\xi \\ &= \Lambda_\varphi(R(x_{(2)})^* \delta^{\frac{1}{2}}) \otimes \hat{J}x_{(1)}\xi \\ &= (\hat{J} \otimes \hat{J})(\Lambda_\varphi(x_{(2)}) \otimes x_{(1)}\xi) \end{aligned}$$

when $x \in \mathcal{N}_\varphi$ and $\xi \in \mathcal{H}$. We see that indeed $(\hat{J} \otimes \hat{J})V(\hat{J} \otimes \hat{J}) = \Sigma W^* \Sigma$.

As we mentioned already, we will show later that \hat{J} is the modular conjugation associated with $\hat{\varphi}$ on \hat{M} and therefore we will get $\hat{J}\hat{M}\hat{J} = \hat{M}'$ and $V \in \hat{M}' \otimes M$.

It would be possible to include more relations at this point, but we will postpone this. First we will show that indeed \hat{J} and $\hat{\nabla}$ are the modular conjugation and the modular operator of $\hat{\varphi}$.

In Section 1, we have mentioned that formally, $K\Lambda_\psi(x) = \Lambda_\psi(S(x)^*)$ for well chosen elements $x \in \mathcal{N}_\psi$. From the proof of Lemma 3.12 we also expect that $K^*\Lambda_\varphi(x) = \Lambda_\varphi(S(x^*))$ for certain elements $x \in \mathcal{N}_\varphi$. We have mentioned these formulas only for a better understanding and motivation. We did not use these formulas in any argument.

Now, as promised, we want to look at these formulas in a correct way. Remember that we used $K = IL^{\frac{1}{2}}$ to denote the polar decomposition of K and that $K^* = IL^{-\frac{1}{2}}$. As we have seen in the previous section (see Lemma 3.11), we expect K to be the closure of the map $\Lambda_\varphi(y) \mapsto \Lambda_\varphi(y^*)$ for $y \in \mathcal{N}_\varphi \cap \mathcal{N}_\varphi^*$.

We will prove a closely related result. It is not quite the same as we explain in the following important remark.

4.10 Remark i) Formally, we have

$$\begin{aligned} \hat{J}\hat{\nabla}^{\frac{1}{2}}\Lambda_\varphi(x) &= \hat{J}\Lambda_\varphi(\tau_{-\frac{i}{2}}(x)\delta^{-\frac{1}{2}}) \\ &= \Lambda_\varphi(R(\tau_{-\frac{i}{2}}(x))^* R(\delta^{-\frac{1}{2}})^* \delta^{\frac{1}{2}}) \\ &= \Lambda_\varphi(S(x)^* \delta) \end{aligned}$$

and

$$\begin{aligned}
\hat{J}\hat{\nabla}^{-\frac{1}{2}}\Lambda_\varphi(x) &= \hat{J}\Lambda_\varphi(\tau_{\frac{i}{2}}(x)\delta^{\frac{1}{2}}) \\
&= \Lambda_\varphi(R(\tau_{\frac{i}{2}}(x))^*R(\delta^{\frac{1}{2}})^*\delta^{\frac{1}{2}}) \\
&= \Lambda_\varphi(S(x^*))
\end{aligned}$$

for appropriate elements $x \in \mathcal{N}_\varphi$. So, we indeed expect that $K = \hat{J}\hat{\nabla}^{\frac{1}{2}}$ and so $I = \hat{J}$ and $L = \hat{\nabla}$. However, although this result is likely to be true, it is not clear how to prove it.

ii) We will not worry about this question (in this paper). After all, we only have used I and L^{it} in the formulas $R(x) = Ix^*I$ and $\tau_t(x) = L^{it}xL^{-it}$ and these formulas remain true when I and L are replaced by \hat{J} and $\hat{\nabla}$ (cf. Proposition 4.7). In other words, we can safely replace I by \hat{J} and L by $\hat{\nabla}$ in the relevant formulas.

iii) We look at this problem closer in the notes [VD8]. See also [K-V3] but remark again that the operator K here should be compared with the operator G^* (see e.g. the introduction and Corollary 2.9 in [K-V3]).

Now, we come to the following important result.

4.11 Theorem Denote by \hat{T} the closure of the conjugate linear map $\Lambda_{\hat{\varphi}}(y) \mapsto \Lambda_{\hat{\varphi}}(y^*)$ where $y \in \mathcal{N}_{\hat{\varphi}} \cap \mathcal{N}_{\hat{\varphi}}^*$. Then, the polar decomposition of \hat{T} is given by $\hat{T} = \hat{J}\hat{\nabla}^{\frac{1}{2}}$ (with the operators \hat{J} and $\hat{\nabla}$ as in Definition 4.5).

Proof: From Lemma 3.11 we know that

$$\langle \Lambda_{\hat{\varphi}}(y^*), \Lambda_\varphi(a) \rangle = \langle \Lambda_{\hat{\varphi}}(y), \Lambda_\varphi(S(a^*)) \rangle^-$$

where $y \in \hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$ and when $a \in \mathcal{N}_\varphi$ is an element such that also $a^* \in \mathcal{D}(S)$ and $S(a^*)$ is still in \mathcal{N}_φ . It follows that \hat{T} is contained in the adjoint of the map $\Lambda_\varphi(a) \mapsto \Lambda_\varphi(S(a^*))$ with a as above.

As we have seen already in the proof of Lemma 3.12, we can relatively easily produce such elements $a \in \mathcal{N}_\varphi$ by requiring that a is analytic both with respect to the automorphism group τ , as well as with respect to multiplication from the right with the unitary group (δ^{it}) . Moreover the space of elements $\Lambda_\varphi(a)$ with such elements a will be left invariant by the operators $\hat{\nabla}^{it}$. All of this will imply that $\Lambda_{\hat{\varphi}}(y) \in \mathcal{D}(\hat{J}\hat{\nabla}^{\frac{1}{2}})$ and that $\hat{J}\hat{\nabla}^{\frac{1}{2}}\Lambda_{\hat{\varphi}}(y) = \Lambda_{\hat{\varphi}}(y^*)$ whenever $y \in \hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*$. Now, a straightforward calculation shows that $\Lambda_{\hat{\varphi}}(\hat{\mathcal{N}} \cap \hat{\mathcal{N}}^*)$ is invariant under the unitaries $\hat{\nabla}^{it}$. Then the result follows.

So, we get as expected, that \hat{J} is the modular conjugation and $\hat{\nabla}$ the modular operator associated with the dual left Haar weight $\hat{\varphi}$.

As a first consequence of this result, we get e.g. that $M \cap \hat{M} = \mathbb{C}1$. Indeed, just before Lemma 3.4 in the previous section, we saw that $M \cap \hat{M}' = \mathbb{C}1$. And because $\hat{J}M\hat{J} = M$ and $\hat{J}\hat{M}\hat{J} = \hat{M}'$, we get that also $M \cap \hat{M} = \mathbb{C}1$.

A second important consequence is the following result (see Proposition 2.1 in [K-V3]).

4.12 Proposition The unitary antipode \hat{R} on \hat{M} is given by $\hat{R}(y) = Jy^*J$. The scaling group $\hat{\tau}$ is given by $\hat{\tau}_t(y) = \nabla^{it}y\nabla^{-it}$ for all $t \in \mathbb{R}$ whenever $y \in \hat{M}$.

This result can be proven in two ways. One argument uses duality as follows. We know that the unitary antipode R on M is given by $R(x) = \hat{J}x^*\hat{J}$ and the scaling group by $\tau_t(x) = \hat{\nabla}^{it}x\hat{\nabla}^{-it}$ (cf. Proposition 4.7). From Theorem 4.11, we know that $\hat{T} = \hat{J}\hat{\nabla}^{\frac{1}{2}}$ is the polar decomposition of the dual operator \hat{T} . Hence, because $J\nabla^{\frac{1}{2}}$ is the polar decomposition of the operator T associated with the left Haar weight φ on M , we will get the formulas in the proposition by duality. A second argument would be possible by using the formulas

$$(\hat{J} \otimes J)W(\hat{J} \otimes J) = W^* \quad \text{and} \quad (\hat{\nabla}^{it} \otimes \nabla^{it})W(\hat{\nabla}^{-it} \otimes \nabla^{-it}) = W$$

and the fact that the right leg of W is dense in \hat{M} (and related things). Remark that these formulas are found in Proposition 1.20 with the operators I and L in the place of \hat{J} and $\hat{\nabla}$ but, as we explained in Remark 4.10 ii), they will still be correct.

Next, we have another important consequence.

4.13 Proposition The scaling constant $\hat{\nu}$ of the dual is ν^{-1} . Furthermore, we also have

$$P^{it}\Lambda_{\hat{\varphi}}(y) = \nu^{-\frac{1}{2}t}\Lambda_{\hat{\varphi}}(\hat{\tau}_t(y))$$

for all $y \in \hat{M}$.

Proof: The proof of this result will use the basic formula $(\tau_t \otimes \hat{\tau}_t)W = W$ for all t . Start with an element $y \in \hat{\mathcal{N}}$ with $y = (\omega \otimes \iota)W$ and $\omega \in M_*$ so that $\omega(a^*) = \langle \hat{\Lambda}(y), \Lambda_{\varphi}(a) \rangle$ for all $a \in \mathcal{N}_{\varphi}$. Then

$$\begin{aligned} \langle P^{it}\hat{\Lambda}(y), \Lambda_{\varphi}(a) \rangle &= \langle \hat{\Lambda}(y), P^{-it}\Lambda_{\varphi}(a) \rangle \\ &= \nu^{-\frac{1}{2}t}\langle \hat{\Lambda}(y), \Lambda_{\varphi}(\tau_{-t}(a)) \rangle \\ &= \nu^{-\frac{1}{2}t}\omega(\tau_{-t}(a^*)) \end{aligned}$$

for all $a \in \mathcal{N}_{\varphi}$. Now we have $((\omega \circ \tau_{-t}) \otimes \iota)W = \hat{\tau}_t(y)$ by the basic formula above. Hence, we see that $\hat{\tau}_t(y) \in \hat{\mathcal{N}}$ and that $P^{it}\hat{\Lambda}(y) = \nu^{-\frac{1}{2}t}\hat{\Lambda}(\hat{\tau}_t(y))$. Then this formula (as in the formulation of the proposition) holds for all $y \in \mathcal{N}_{\hat{\varphi}}$.

From this formula (and because P^{it} is a unitary operator), it follows that $\hat{\varphi} \circ \hat{\tau}_t = \nu^t \hat{\varphi}$ for all t and therefore $\hat{\nu} = \nu^{-1}$.

We see that (P^{it}) is, in some sense, a self-dual group of unitaries. One can write $\hat{P} = P$ when \hat{P} would be defined for the dual as P is defined for the original pair (M, Δ) . So, this one-parameter group of unitaries implements the scaling group both on M and on \hat{M} . In combination with the formula $(S \otimes \iota)W = W^*$, it will give that W is a manageable

multiplicative unitary (in the sense of [W2]). As we have already mentioned however, we will not use this property as such.

We will now use a similar technique as in the proof of Proposition 4.13 to obtain more relations. Recall that in the above proof, we used the basic formula $(\tau_t \otimes \hat{\tau}_t)W = W$. First, we will now formulate again some other results of this type.

Recall that formally, we have $W^*(\xi \otimes \Lambda_\varphi(x)) = \sum x_{(1)}\xi \otimes \Lambda_\varphi(x_{(2)})$ (when we use $\Delta(x) = \sum x_{(1)} \otimes x_{(2)}$) with $x \in \mathcal{N}_\varphi$ and $\xi \in \mathcal{H}$. Then, from the definitions of (u_t) , (v_t) and (w_t) and the formulas

$$\begin{aligned}\Delta(\sigma_t(x)) &= (\tau_t \otimes \sigma_t)\Delta(x) \\ \Delta(\sigma'_t(x)) &= (\sigma'_t \otimes \tau_{-t})\Delta(x) \\ \Delta(\tau_t(x)) &= (\tau_t \otimes \tau_t)\Delta(x) \\ &= (\sigma_t \otimes \sigma'_{-t})\Delta(x)\end{aligned}$$

for all $x \in M$ and $t \in \mathbb{R}$, we get the following relations (see also the proof of Theorem 2.10).

4.14 Lemma We have

$$\begin{aligned}(\tau_t \otimes \iota)W &= (1 \otimes u_t^*)W(1 \otimes u_t) \\ (\sigma'_t \otimes \iota)W &= (1 \otimes w_t^*)W(1 \otimes v_t^*) \\ (\tau_t \otimes \iota)W &= (1 \otimes v_t^*)W(1 \otimes v_t) \\ (\sigma_t \otimes \iota)W &= (1 \otimes v_t^*)W(1 \otimes w_t^*)\end{aligned}$$

for all $t \in \mathbb{R}$.

Each of the four formulas comes from one of the above relations, in the same order. The first and the third formula do not give anything new. This is simply the fact that $(\tau_t \otimes \hat{\tau}_t)W = W$, combined with the knowledge that both u_t and v_t implement $\hat{\tau}_t$ on \hat{M} .

However, using all these results, in combination with the definition of the dual map $\hat{\Lambda}$, as in the proof of Proposition 4.13, we obtain the following result about the modular element $\hat{\delta}$, relating the left and the right Haar weights on the dual $(\hat{M}, \hat{\Delta})$.

4.15 Proposition We have $\hat{\delta}^{it} = v_t^* w_t^*$ where v_t and w_t are defined as in Definition 4.1. Furthermore $\hat{\Delta}(\hat{\delta})^{it} = \hat{\delta}^{it} \otimes \hat{\delta}^{it}$ for all t .

Proof: Because we have

$$\begin{aligned}(\sigma_t \otimes \iota)W &= (1 \otimes v_t^*)W(1 \otimes w_t^*) \\ &= (1 \otimes v_t^*)W(1 \otimes v_t)(1 \otimes v_t^* w_t^*) \\ &= ((\tau_t \otimes \iota)W)(1 \otimes v_t^* w_t^*),\end{aligned}$$

we see that $w_t v_t \in \hat{M}$ for all t .

Now, if we use the same type of argument as in the proof of the previous proposition, we get $\hat{\tau}_t(y)v_t w_t \in \hat{\mathcal{N}}$ when $y \in \hat{\mathcal{N}}$ and

$$\nabla^{it} \hat{\Lambda}(y) = \hat{\Lambda}(\hat{\tau}_t(y)v_t w_t).$$

If we compare this with the earlier and similar formula

$$\hat{\nabla}^{it} \Lambda_\varphi(y) = \Lambda_\varphi(\tau_t(x)\delta^{-it})$$

for $x \in \mathcal{N}_\varphi$, we see that we must have $\hat{\delta}^{it} = v_t^* w_t^*$ for all t (by duality). Recall that v_t and w_t commute with each other.

From the formula $(\sigma'_t \otimes \iota)W = (1 \otimes w_t^*)W(1 \otimes v_t^*)$ we get

$$(w_t \otimes w_t)W(w_t^* \otimes v_t) = W$$

and from the formula $(\tau_t \otimes \iota)W = (1 \otimes v_t^*)W(1 \otimes v_t)$ we find

$$(v_t \otimes v_t)W(v_t^* \otimes v_t^*) = W$$

and combining these two results, we obtain

$$(\hat{\delta}^{-it} \otimes \hat{\delta}^{-it})W(\hat{\delta}^{it} \otimes 1) = W$$

proving that $\hat{\Delta}(\hat{\delta}^{it}) = \hat{\delta}^{it} \otimes \hat{\delta}^{it}$.

4.16 Remark By duality, we also will get $\Delta(\delta^{it}) = \delta^{it} \otimes \delta^{it}$ for all t .

This seems to be a strange (and certainly not an obvious) way to prove this basic formula. One could expect a more direct proof (e.g. along the lines of the proof of this formula for algebraic quantum groups in [VD3]). However, an attempt to do this turns out to become quite involved (see Section 7 in [K-V2]). In [M-N-W], this formula is proven in a more elegant way, but also uses results, not only about (M, Δ) itself, but also about the dual $(\hat{M}, \hat{\Delta})$ (see Proposition 6.12 in [M-N-W]).

Now, we are ready to collect the main formulas and relations. First, we have a number of formulas that express some of the operators in terms of the others.

4.17 Theorem We have the following formulas for the four modular operators:

$$\begin{aligned} \nabla^{it} &= (\hat{J}\hat{\delta}^{it}\hat{J})P^{it} \\ \hat{\nabla}^{it} &= (J\delta^{it}J)P^{it} \\ \nabla^{it} &= \hat{\delta}^{-it}P^{-it} \quad (= \hat{J}\nabla^{-it}\hat{J}) \\ \hat{\nabla}^{it} &= \delta^{-it}P^{-it} \quad (= J\hat{\nabla}^{-it}J). \end{aligned}$$

The second formula was obtained in Proposition 4.6. The first one follows e.g. by duality. From 4.8 we get the third formula while the last one follows again by duality.

We also have a number of *commutation rules*. We know that P^{it} commutes with J , \hat{J} , δ and $\hat{\delta}$ (and as a consequence of the above relations, also with all these modular operators). We also know that J commutes with $\hat{\delta}^{it}$ (because $\hat{R}(\hat{\delta}) = \hat{\delta}^{-1}$) and that \hat{J} commutes with δ^{it} (because $R(\delta) = \delta^{-1}$). The non-trivial commutation rules are formulated in the next two theorems.

4.18 Theorem We have

$$\begin{aligned}\hat{J}\nabla^{-it}\hat{J} &= \delta^{it}(J\delta^{it}J)\nabla^{it} \\ J\hat{\nabla}^{-it}J &= \hat{\delta}^{it}(\hat{J}\hat{\delta}^{it}\hat{J})\hat{\nabla}^{it}.\end{aligned}$$

The first formula is a combination of the formulas in Proposition 4.3 and 4.8. The second one comes with duality.

Next, we get the following two basic commutation rules.

4.19 Theorem We have $\hat{\delta}^{is}\delta^{it} = \nu^{-ist}\delta^{it}\hat{\delta}^{is}$ for all $s, t \in \mathbb{R}$. Also $\hat{J}J = \nu^{\frac{1}{4}}J\hat{J}$.

Proof: The first formula in this theorem can be obtained from the fact that $\sigma_s(\delta^{it}) = \nu^{ist}\delta^{it}$, in combination with the first formula in the previous theorem. To prove the second result, one can calculate explicitly the operator $\hat{J}T\hat{J}$ and use the uniqueness of the polar decomposition of the operator T .

The uniqueness of the polar decomposition of T , as used above, will also give the first formula of Theorem 4.18. On the other hand, it would also be possible to use this formula to give a more direct proof of the commutation rule between \hat{J} and J .

If we combine the formulas of Theorem 4.17 with the commutation rules in Theorem 4.18 and Theorem 4.19, we get most (if not all) of the other commutation rules. There is no need to include them all here (see e.g. Proposition 2.13 in [K-V3]).

Let us now *finish this section* with another formula, expressing the operators P^{it} in terms of the other operators. It is an analytic version of Radford's formula for the 4th power of the antipode, proven in the case of a finite-dimensional Hopf algebra in [R] and extended to the case of algebraic quantum groups in [D-VD2].

4.20 Theorem We have

$$P^{-2it} = \delta^{it}(J\delta^{it}J)\hat{\delta}^{it}(\hat{J}\hat{\delta}^{it}\hat{J})$$

for all t .

Proof: From Theorem 4.18 we get

$$\hat{J}\nabla^{-it}\hat{J} = \delta^{it}(J\delta^{it}J)\nabla^{it}$$

and if we replace in this formula ∇^{it} by $(\hat{J}\hat{\delta}^{it}\hat{J})P^{it}$ two times, we arrive at the desired result.

This is indeed one of the possible analytical versions of the formula, valid in the theory of finite-dimensional Hopf algebras

$$S^4(a) = \delta^{-1}(\hat{\delta} \triangleright a \triangleleft \hat{\delta}^{-1})\delta$$

where \triangleright and \triangleleft are used to denote the canonical left and right actions of the dual algebra. Remark that $P^{2it}\Lambda_\varphi(a) = \nu^t\Lambda_\varphi(\tau_{2t}(a))$ and that $\tau_{-i} = S^2$. Also $\delta^{it}J\delta^{it}J\Lambda_\varphi(a) = \nu^{\frac{1}{2}t}\Lambda_\varphi(\delta^{it}a\delta^{-it})$. To explain the last part of the formula, one should observe that the operator $\hat{\delta}^{it}$ is a convolution operator on M , but on the Hilbert space level and similarly for $\hat{J}\hat{\delta}^{it}\hat{J}$. The first one is 'left' convolution and the second one is 'right' convolution.

We have now the essential formulas, all formulated in these four last theorems. Also the commutation rules with the left regular representation, as formulated in Lemma 4.14, are essentially taken care of.

One can also draw certain interesting conclusions from these formulas. If e.g. the left and the right Haar weight on M coincide, that is when $\delta = 1$, it will follow from the commutation rules in Theorem 4.19 that the scaling constant ν has to be one. If both δ and $\hat{\delta}$ are trivial, it will follow from the last theorem that also $P = 1$ and that the scaling automorphisms are trivial. From Theorem 4.17, it will follow that all the modular automorphisms are trivial in this case and so that all Haar weights must be traces. Surely, there are other arguments for these statements, but it is nice to see how they follow from all these formulas.

Appendix: Other approaches to locally compact quantum groups

In this appendix, we will relate the von Neumann algebra approach (in particular, as it is treated in this paper) with the C*-algebra approach. We will consider the papers by Kustermans and Vaes [K-V1] and [K-V2], as well as the paper by Masuda, Nakagami and Woronowicz [M-N-W]. We will also briefly make a comparison with the earlier paper by Masuda and Nakagami [M-N].

Such a comparison is certainly interesting. In fact, the relation with (the equivalent) C*-algebra approach is not only interesting, it is also an important feature of the theory. Nevertheless, because of the scope of this paper, we will be very brief here. In [VD6], some more details are already found, but we refer to forthcoming papers for all the details about the material in this appendix (see [Q-VD] for results about weights on C*-algebras and [VD8] for the theory of locally compact quantum groups).

Now, we start with a C*-algebra A and a comultiplication on A . Recall the definition:

A.1 Definition A *comultiplication on a C^* -algebra A* is a non-degenerate $*$ -homomorphism Δ from A to the multiplier algebra $M(A \otimes A)$ of the minimal C^* -tensor product $A \otimes A$ of A with itself, satisfying coassociativity $(\Delta \otimes \iota)\Delta = (\iota \otimes \Delta)\Delta$. It is also assumed that slices $(\omega \otimes \iota)\Delta(a)$ and $(\iota \otimes \omega)\Delta(a)$ belong to A for all $a \in A$ and $\omega \in A^*$.

Non-degenerate here means that $\Delta(A)(A \otimes A)$ is dense in $A \otimes A$. Because of this condition, the $*$ -homomorphisms $\Delta \otimes \iota$ and $\iota \otimes \Delta$ have unique extensions (still denoted in the same way) to unital $*$ -homomorphisms from $M(A \otimes A)$ to $M(A \otimes A \otimes A)$. Therefore, coassociativity, as formulated above, has a meaning. Slice maps are defined from $M(A \otimes A)$ to $M(A)$. So, in general these slices $(\omega \otimes \iota)\Delta(a)$ and $(\iota \otimes \omega)\Delta(a)$ belong to $M(A)$. In [M-N-W], it is assumed that $\Delta(A)(1 \otimes A)$ and $\Delta(A)(A \otimes 1)$ are subsets of $A \otimes A$. Because any $\omega \in A^*$ is of the form $\rho(a \cdot)$ (and of the form $\rho(\cdot a)$) with $\rho \in A^*$ and $a \in A$, the latter conditions will imply the conditions in the definition above. In fact, it is shown in the theory that also these stronger conditions are valid. See e.g. a remark following Proposition 1.21 in Section 1 of this paper.

The following result is not so difficult to obtain.

A.2 Proposition Let (M, Δ) be a locally compact quantum group (as in Definition 2.1 of this paper) with left Haar weight φ . Let W be the left regular representation (as introduced in Section 1) and define A to be the norm closure of the space of slices $\{(\iota \otimes \omega)W \mid \omega \in \mathcal{B}(\mathcal{H}_\varphi)_*\}$. Then A is a C^* -algebra, it is a σ -weakly dense subalgebra of M and the restriction of Δ to A is a comultiplication on A .

It is easy to see (and a standard result about multiplicative unitaries) that A is a subalgebra. And just as in the case of the dual (see Proposition 3.3 in this paper), one can show that it is a $*$ -subalgebra. As the left leg of W is dense in M , it follows that also A is dense in M . To show that Δ maps A into $M(A \otimes A)$ is again a standard result for multiplicative unitaries (see [B-S]). See also the work on manageable multiplicative unitaries ([W2] and [S-W]). Also here, we refer to [VD8] for a detailed and independent approach.

Later in this section (see Definition A.10), we will recall the definition of a locally compact quantum group in the C^* -algebra context, as given by Kustermans and Vaes in [K-V1] and [K-V2]. First, we look at some results about (invariant) weights on C^* -algebras (with a comultiplication). The first one is about the kind of weights that is used in this theory.

A.3 Proposition Let A be a C^* -algebra. Let φ be a densely defined, lower semi-continuous weight on A . Denote by $\tilde{\varphi}$ the normal weight on the double dual A^{**} that extends φ . Denote by e the support projection of $\tilde{\varphi}$ in A^{**} . Then φ is approximately K.M.S. (see Definition 1.34 in [K-V2]) if and only if e is a central projection in A^{**} .

Recall that a faithful, densely defined lower semi-continuous weight on a C^* -algebra is approximately K.M.S. if it remains faithful when extended to $\pi_\varphi(A)''$. This last property essentially means that the support of π_φ in A^{**} (which is a central projection), coincides with the support projection of $\tilde{\varphi}$. So, the above result should not come as a surprise. We

refer to [Q-VD] for details; see also [VD8]. Further in this appendix, we will speak about a weight on a C^* -algebra with central support, or shortly call it a *central weight*.

Invariant weights on C^* -algebras with a comultiplication are defined as usual:

A.4 Definition Let A be a C^* -algebra with a comultiplication Δ (as in Definition A.1). A weight φ is called *left invariant* if $\varphi((\omega \otimes \iota)\Delta(a)) = \|\omega\|\varphi(a)$ whenever $a \in A$, $a \geq 0$ and $\varphi(a) < \infty$ and when $\omega \in A^*$ and $\omega \geq 0$. Similarly, a *right invariant* weight is defined.

Again, there is the following result.

A.5 Proposition Let (M, Δ) be a locally compact quantum group. The restriction of the left Haar weight φ to the C^* -subalgebra A of M , defined as in Proposition A.2, is a faithful, densely defined, lower semi-continuous central and left invariant weight. Similarly for the right Haar weight.

Proof: It is quite obvious that these restrictions are faithful, lower semi-continuous and invariant weights. They are central because they are restrictions of faithful weights to a C^* -subalgebra of the von Neumann algebra M . To show that the right invariant weight ψ is still densely defined on the C^* -algebra A , one can use the result in Lemma 1.13 of Section 1. To show that also the left invariant weight φ is still densely defined on the C^* -algebra, one can use that the unitary antipode R leaves the C^* -algebra invariant (a result that follows from the formula $(\hat{J} \otimes J)W(\hat{J} \otimes J) = W^*$ and the definition of R).

Using the formulas in the proof of Theorem 2.10 (or equivalently, the ones in Lemma 4.14), we see that the C^* -algebra A is invariant under the modular automorphism groups σ and σ' . Therefore, the restrictions of the Haar weights φ and ψ are KMS-weights (and so certainly central). Observe also that the C^* -algebra is not only invariant under the unitary antipode, but also under the scaling automorphisms τ , again see Lemma 4.14 (or equivalent earlier results).

We will now consider two important results. They will enable us to go quickly from the C^* -algebra setting to the von Neumann algebra framework.

A.6 Proposition Let A be a C^* -algebra with a comultiplication Δ . Assume that φ is a densely defined lower semi-continuous central weight on A . Consider the extension $\tilde{\varphi}$ on A^{**} of φ . Also extend Δ to a normal and unital $*$ -homomorphism $\tilde{\Delta} : A^{**} \rightarrow A^{**} \otimes A^{**}$ (the von Neumann algebra tensor product). Then $\tilde{\varphi}$ is still left invariant.

One uses left Hilbert algebra theory to show that the G.N.S.-space of the extension is the same as the original one. This result is not completely trivial but is essentially proven in Section 2 of Chapter VII on weights in [T3]. Then it is shown that the extension is still invariant. The way this is done, is similar as the invariance of the dual weight is proven in

Proposition 3.15. Essentially, the argument used to prove that left invariance of φ implies the formula $WW^* = 1$ for the left regular representation W , is 'reversed'.

A.7 Proposition As in Proposition A.6, assume that A is a C^* -algebra with a comultiplication Δ . Now assume that φ and ψ are non-trivial, densely defined, lower semi-continuous and central weights on A such that φ is left invariant and ψ is right invariant. Then the supports of φ and ψ in A^{**} are equal.

Proof: Denote by e and f the supports of $\tilde{\varphi}$ and $\tilde{\psi}$ resp. By assumption, they are central projections in A^{**} . By the left invariance of $\tilde{\varphi}$, we get $\tilde{\varphi}((\omega \otimes \iota)\tilde{\Delta}(1-e)) = 0$ for all $\omega \in A^*$ with $\omega \geq 0$. This implies that $(\tilde{\Delta}(1-e))(1 \otimes e) = 0$. Because e is central, we also get $(\tilde{\Delta}(x^*(1-e)x))(1 \otimes e) = 0$ for all $x \in A^{**}$ satisfying $\tilde{\psi}(x^*x) < \infty$. If we apply $\tilde{\psi}$, we get by using right invariance, that $\tilde{\psi}(x^*(1-e)x)e = 0$. So, $\tilde{\psi}(x^*(1-e)x) = 0$ because e is non-zero. Then we get $f(1-e) = 0$. A similar argument will give $e(1-f) = 0$ and therefore $e = f$.

Compare this result with Theorem 3.8 in [K-V2].

It follows immediately from this result that all invariant weights have the same support. This implies that we have a single von Neumann algebra. Indeed, we can consider the associated von Neumann algebra M , defined as $A^{**}e$, where e is the support of the non-trivial, densely defined, lower semi-continuous, central invariant weights.

All the previous results lead to the following which is the main theorem of this appendix.

A.8 Theorem Let A be a C^* -algebra with a comultiplication Δ . Assume that there exist faithful, densely defined lower semi-continuous weights φ and ψ , with central support and resp. left and right invariant. Let $M = A^{**}e$ where e is the support of these weights. Consider the extension $\tilde{\Delta}$ of Δ to A^{**} (as in the proof of Proposition A.6). Then restrict to M and define $\Delta_0(x) = \tilde{\Delta}(x)(e \otimes e)$ for $x \in M$. This is a comultiplication on the von Neumann algebra M . The restrictions to M of the extensions $\tilde{\varphi}$ and $\tilde{\psi}$ (as in Proposition A.3) are a left and a right Haar weight on (M, Δ_0) (as in Definition 1.4 and further in Section 1). So, the pair (M, Δ_0) is a locally compact quantum group in the sense of Definition 2.1.

Remark that $(\tilde{\Delta}(1-e))(1 \otimes e) = 0$ as we saw in the proof of Proposition A.6. Then $(\tilde{\Delta}(e))(1 \otimes e) = 1 \otimes e$ and so $\Delta_0(e) = e \otimes e$. This guarantees that the comultiplication Δ_0 on M is unital. This is also needed to show that Δ_0 is still coassociative.

So, roughly speaking, Theorem A.7 says that starting with a C^* -algebra with a comultiplication and nice invariant weights, we can associate a locally compact quantum group (in the von Neumann algebraic sense). If we combine Proposition A.2 with Proposition A.5, we see that also conversely, given a locally compact quantum group, we can associate a pair of a C^* -algebra and a comultiplication with nice invariant weights.

What happens when we perform these two operations, one after the other? Do we get back the original?

First, start with a locally compact quantum group (M, Δ) . Consider the C^* -algebra A as in Proposition A.2 and restrict Δ as in Proposition A.7. Then it is rather straightforward to show that the construction in Theorem A.7 will yield the original pair (M, Δ) .

On the other hand, take a C^* -algebra A with a comultiplication Δ and nice invariant weights. Consider the pair (M, Δ_0) as in Theorem A.7. We have the following lemma.

A.9 Lemma Let W be the left regular representation for the pair (M, Δ_0) . Then the norm closure of the space

$$\{(\iota \otimes \omega)W \mid \omega \in \mathcal{B}(\mathcal{H}_\varphi)_*\}$$

is the same as the norm closure of the space

$$\text{sp}\{(\iota \otimes \omega)\Delta(a) \mid a \in A, \omega \in A^*\}.$$

This result is essentially found along with the proof of Proposition 1.21 in Section 1.

It is not clear whether or not, this space will be all of A . So, in order to recover the original C^* -algebra, we need the extra *density conditions* as they are found in the original definition of a locally compact quantum group in the C^* -algebra setting (as given by Kustermans and Vaes in [K-V2]). We recall the definition here.

A.10 Definition Let A be a C^* -algebra and Δ a comultiplication on A (as in Definition A.1). Assume that the spaces

$$\begin{aligned} &\text{sp}\{(\iota \otimes \omega)\Delta(a) \mid a \in A, \omega \in A^*\} \\ &\text{sp}\{(\omega \otimes \iota)\Delta(a) \mid a \in A, \omega \in A^*\} \end{aligned}$$

are (norm) dense in A . Assume that there exist faithful, densely defined lower semi-continuous weights φ and ψ on A , both with central support and resp. left and right invariant. Then (A, Δ) is called a locally compact quantum group in the C^* -algebraic sense.

It is only for such a pair (A, Δ) that we have a complete equivalence of the C^* -algebraic and von Neumann algebraic setting. If the density conditions are not fulfilled, in some sense, the C^* -algebra might be too big. As we see from the previous discussion, we can pass to a smaller C^* -algebra, invariant under the comultiplication, satisfying the conditions of Definition A.10. However, it should be remarked that at present, there are no (obvious) examples of this phenomenon.

Next, one needs to prove properties for a locally compact quantum group (A, Δ) with a C^* -algebra A and a comultiplication Δ as in Definition A.10. These properties will follow from the ones proven for a locally compact quantum group (M, Δ) as defined in Definition 2.1. We have already given some indications (about the stronger density conditions, the invariance of the C^* -algebra under the modular automorphisms, the scaling automorphisms,

the unitary antipode, ...). But more results need to be considered. One can e.g. show relatively easy that $\delta^{it} \in M(A)$ for all t where δ is the modular element from Theorem 2.11. There is also the construction of the dual \hat{A} . It can be obtained either by applying the procedure in this appendix and find \hat{A} from \hat{M} (as in A.2 and A.5) or directly as the norm closure of the space $\{(\omega \otimes \iota)W \mid \omega \in M_*\}$ (as indicated in a remark following Proposition 3.3 - compare also with the result in Lemma A.9).

So far about the comparison of our approach in this paper, with the C*-algebraic approach by Kustermans and Vaes. Let us now also compare (briefly) with the approach of Masuda, Nakagami and Woronowicz.

There is first the original work by Masuda and Nakagami [M-N] where locally compact quantum groups are studied in the von Neumann algebra framework. Then, there there is the more recent work by these two authors and Woronowicz [M-N-W] where locally compact quantum groups are studied in the C*-algebra setting. We will not say anything more about the first paper as in some sense, the second one can be seen as an improvement of the first one. We refer to the introduction of [M-N-W] for a comparison of the two papers.

The main difference between the set of axioms in [M-N-W] and those formulated by Kustermans and Vaes is that in the first case, the antipode and its polar decomposition are assumed whereas in [K-V2], the antipode is constructed and its properties are proven. The same holds for our approach.

On the other hand, for this approach, as we have seen, we need to assume the existence of both a left and a right Haar weight. This is not the case in the approach of [M-N-W] where only a left Haar weight φ is assumed. However, the unitary antipode R (see Definition 1.5 in [M-N-W]) is part of the axioms and from all the axioms, it is not so hard to obtain that R flips the coproduct (see Proposition 2.6 in [M-N-W]). Then the composition $\varphi \circ R$ gives a right Haar weight.

Moreover, in examples, it is often rather easy to see what this unitary antipode should be and to verify that it flips the coproduct. Hence, usually, only one Haar weight is constructed explicitly while the other one is obtained by composing it with this candidate for the unitary antipode. Therefore it is in general more easy to verify the axioms of Kustermans and Vaes.

Another (minor) difference is that in [K-V2] weaker density conditions are needed (see earlier). Also a stronger invariance condition is proven by Kustermans and Vaes.

It is well-known that Masuda, Nakagami and Woronowicz started with their work on locally compact quantum groups, earlier than Kustermans and Vaes but that it took many years before their results were published. To my opinion, it is clear that the approach of Kustermans and Vaes, and also our approach in this paper, is better than their approach. The axioms are more complicated and this does not really help to get the main results in a quicker way. On the other hand, the contribution of Masuda, Nakagami and Woronowicz is of great importance to the theory and their paper contains interesting material and nice independent results.

Let us finish this discussion by pointing out that it was in fact Kirchberg, at a conference in Copenhagen, in 1991, who first formulated the idea of generalizing the axioms of Kac

algebras and thereby considering the polar decomposition of the antipode ([K]). As far as I know, his work was never published.

References

- [A] E. Abe: *Hopf algebras*. Cambridge University Press (1977).
- [B-S] S. Baaĳ & G. Skandalis: *Unitaires multiplicatifs et dualité pour les produits croisés de C^* -algèbres*. Ann. Scient. Ec. Norm. Sup., 4ème série, **26** (1993), 425-488.
- [D-VD1] L. Delvaux & A. Van Daele: *The Drinfel'd double versus the Heisenberg double for an algebraic quantum group*. J. Pure & Appl. Alg. 190 (2004), 59–84.
- [D-VD2] L. Delvaux & A. Van Daele: *Traces on (group-cograded) multiplier Hopf algebras*. Preprint K.U.Leuven and Universiteit Hasselt (2005).
- [E-S] M. Enock & J.-M. Schwartz: *Kac algebras and duality for locally compact groups*. Springer (1992).
- [K] E. Kirchberg: Lecture at the conference 'Invariance in Operator Algebras', Copenhagen, August 1992.
- [K-V1] J. Kustermans & S. Vaes: *A simple definition for locally compact quantum groups*. C.R. Acad. Sci., Paris, Sér. I **328** (10) (1999), 871–876.
- [K-V2] J. Kustermans & S. Vaes: *Locally compact quantum groups*. Ann. Sci. Éc. Norm. Sup. **33** (2000), 837–934.
- [K-V3] J. Kustermans & S. Vaes: *Locally compact quantum groups in the von Neumann algebra setting*. Math. Scand. **92** (2003), 68–92.
- [K-V4] J. Kustermans & S. Vaes: *The operator algebra approach to quantum groups*. Proc. Natl. Acad. Sci. USA **97** (2) (2000), 547–552.
- [K-^{*}] J. Kustermans, S. Vaes, L. Vainerman, A. Van Daele and S.L. Woronowicz: *Locally compact quantum groups*. Lecture Notes School/Conference on Noncommutative Geometry and Quantum groups, Warsaw, 2001. Banach Centre Publications, to appear.
- [M-VD1] A. Maes & A. Van Daele: *Notes on Compact Quantum Groups*. Nieuw Archief voor Wiskunde, Vierde serie **16** (1998), 73–112.
- [M-VD2] A. Maes & A. Van Daele: *The multiplicative unitary as a basis for duality*. Preprint K.U. Leuven (2002), 37p. (#math.OA/0205284).
- [M-N] M. Masuda & Y. Nakagami: *A von Neumann algebra framework for the duality of quantum groups*. Publ. RIMS Kyoto **30** (1994), 799–850.
- [M-N-W] T. Masuda, Y. Nakagami & S.L. Woronowicz: *A C^* -algebraic framework for the quantum groups*. Int. J. of Math. **14** (2003), 903–1001.

- [Q-VD] J. Quaegebeur & A. Van Daele: *Weights on C^* -algebras*. Preprint K.U.Leuven. In preparation.
- [R] D.E. Radford: *The order of the antipode of a finite-dimensional Hopf algebra is finite*. Amer. J. Math. **98** (1976), 333–355.
- [S-W] P. Soltan & S.L. Woronowicz: *A remark on manageable multiplicative unitaries*. Letters on Math. Phys. **57** (2001), 239–252.
- [St] S. Stratila: *Modular Theory in Operator Algebras*. Abacus Press, Turnbridge Wells, 1981.
- [S-V-Z1] S. Stratila, D. Voiculescu & L. Zsidó: *Sur les produits croisés*. Comptes Rendus Acad. Sc. Paris **282** (1976), ??–??
- [S-V-Z2] S. Stratila, D. Voiculescu & L. Zsidó: *On crossed products I*. Rev. Roumaine Math. Pure et Appl. **21** (1976), 1411–1449.
- [S-V-Z3] S. Stratila, D. Voiculescu & L. Zsidó: *On crossed products II*. Rev. Roumaine Math. Pure et Appl. **22** (1977), 83–117.
- [S-Z] S. Stratila & L. Zsidó: *Lectures on von Neumann algebras*. Abacus Press, Tunbridge Wells, 1979.
- [Sw] M. Sweedler: *Hopf algebras*. Benjamin, New-York (1969).
- [T1] M. Takesaki: *Tomita's theory of modular Hilbert algebras and applications*. Springer Lecture Notes in Mathematics 128. Springer (1970).
- [T2] M. Takesaki: *Theory of Operator Algebras I*. Springer-Verlag, New York (1979).
- [T3] M. Takesaki: *Theory of Operator Algebras II*. Springer-Verlag, New York (2001).
- [Tay] D.C. Taylor: *The strict topology for double centralizer algebras*. Trans. Am. Math. Soc. **150** (1970), 633–643.
- [V1] S. Vaes: *A Radon-Nikodym theorem for von Neumann algebras*. J. Operator Theory **46** (2001), 477–489.
- [V2] S. Vaes: *Locally compact quantum groups*. PhD. Thesis K.U.Leuven (2001).
- [V-VD] S. Vaes & A. Van Daele: *Hopf C^* -algebras*. Proc. London Math. Soc. **82** (2001), 337–384.
- [VD1] A. Van Daele: *Dual pairs of Hopf $*$ -algebras*. Lecture notes of a talk at the University of Orléans. Preprint K.U.Leuven (1991).
- [VD2] A. Van Daele: *Multiplier Hopf algebras*. Trans. Amer. Math. Soc. **342** (1994), 917–932.
- [VD3] A. Van Daele: *An algebraic framework for group duality*. Adv. in Math. **140** (1998), 323–366.
- [VD4] A. Van Daele: *Quantum groups with invariant integrals*. Proc. Natl. Acad. Sci. USA **97** (2000), 541–546.
- [VD5] A. Van Daele: *The Haar measure on some locally compact quantum groups*. Preprint K.U. Leuven (2001)., math.OA/0109004.

- [VD6] A. Van Daele: *Locally compact quantum groups: The von Neumann algebra versus the C^* -algebra approach*. Preprint K.U. Leuven (17p) (2005). To appear in The Bulletin of Kerala Mathematics Association.
- [VD7] A. Van Daele: *Notes on Multiplier Hopf $(^*-)algebras$ and algebraic quantum groups*. Preprint K.U. Leuven. In preparation.
- [VD8] A. Van Daele: *Notes on Locally Compact Quantum Groups*. Preprint K.U. Leuven. In preparation.
- [VH] L. Vanheeswijck: *Duality in the theory of crossed products*. Math. Scand. **44** (1979), 313–329.
- [W1] S.L. Woronowicz: *Compact Quantum Groups*. Quantum symmetries/Symmétries quantiques. Proceedings of the Les Houches summer school 1995, North-Holland, Amsterdam (1998), 845–884.
- [W2] S.L. Woronowicz: *From multiplicative unitaries to quantum groups*. Int. J. Math. **7** (1996), 127–149.