

Perverse equivalences and cacti


Iva Halacheva
(Northeastern University)

**Workshop on Lie Theory and Integrable Systems in
Symplectic and Poisson Geometry (Fields Institute)**

June 6, 2020

Overview

Let \mathfrak{g} be a simply-laced Kac–Moody Lie algebra.



Categorical \mathfrak{g} -action I

Warm-up case: $\mathfrak{g} = \mathfrak{sl}_2$

Suppose $\mathfrak{sl}_2 = \mathbb{C}\{e, f, h\} \curvearrowright V = \bigoplus_{n \in \mathbb{Z}} V_n$, an integrable \mathfrak{sl}_2 -rep.

$$e : V_n \rightarrow V_{n+2}, \quad f : V_n \rightarrow V_{n-2}, \quad (ef - fe)|_{V_n} = n \text{Id}_{V_n} \quad \forall n \in \mathbb{Z}.$$

Categorified \mathfrak{sl}_2 -action (Chuang–Rouquier, Khovanov–Lauda):

- an abelian category $\mathcal{C} = \bigoplus_n \mathcal{C}_n$ (with $K_0(\mathcal{C}_n) = V_n$)
- exact endofunctors E, F of \mathcal{C} , $E : \mathcal{C}_n \rightarrow \mathcal{C}_{n+2}, F : \mathcal{C}_n \rightarrow \mathcal{C}_{n-2}$
- natural transformations

$$\begin{aligned} \epsilon : EF \rightarrow I, \quad \eta : I \rightarrow FE & \quad (\text{unit and counit of adjunction}) \\ X : E \rightarrow E, \quad T : E^2 \rightarrow E^2 \end{aligned}$$

such that...

Categorical \mathfrak{g} -action II

- For $n \geq 0$ (analogously for $n < 0$), we have an isomorphism:

$$(\sigma, \epsilon, \epsilon \circ X|_F, \dots, \epsilon \circ X^{n-1}|_F) : EF|_{C_n} \xrightarrow{\cong} FE|_{C_n} \oplus I_{C_n}^{\oplus n},$$

where σ is composed of η , T , and ϵ .

- The natural transformations X , T give an action of the nil affine Hecke algebra H_n on E^n .

Example The adjoint representation of \mathfrak{sl}_2 :

$$(C_2 = \mathbb{C} - \text{mod}) \begin{array}{c} \xrightarrow{\text{Ind}} \\ \xleftarrow{\text{Res}} \end{array} (C_0 = \mathbb{C}[x]/x^2 - \text{mod}) \begin{array}{c} \xrightarrow{\text{Res}} \\ \xleftarrow{\text{Ind}} \end{array} (C_{-2} = \mathbb{C} - \text{mod})$$

Categorical \mathfrak{g} -action III

More generally:

The 2-**category** $\mathcal{U}_{\mathfrak{g}}$ categorifies (Lusztig's idempotent form) $\dot{U}_{\mathfrak{g}}$:

- objects are elements λ of the \mathfrak{g} -weight lattice.
- 1-morphisms are generated by $E_i : \lambda \rightarrow \lambda + \alpha_i, F_i : \lambda \rightarrow \lambda - \alpha_i$.
- 2-morphisms are generated by

$$\begin{aligned} X_i &= \begin{array}{c} \uparrow \\ \bullet \\ \downarrow \end{array}_i : E_i \rightarrow E_i, & X_i &= \begin{array}{c} \downarrow \\ \bullet \\ \uparrow \end{array}_i : F_i \rightarrow F_i, \\ T_{ij} &= \begin{array}{c} \nearrow \searrow \\ i \quad j \end{array} : E_i E_j \rightarrow E_j E_i, & T_{ij} &= \begin{array}{c} \searrow \nearrow \\ i \quad j \end{array} : F_i F_j \rightarrow F_j F_i \\ \cap^i &: E_i F_i \rightarrow I, & \cup^i &: F_i E_i \rightarrow I, & \cup^i &: I \rightarrow F_i E_i, & \cap^i &: I \rightarrow E_i F_i \end{aligned}$$

+KLR and further relations.

A **categorical \mathfrak{g} -representation** is a 2-functor $\mathcal{U}_{\mathfrak{g}} \rightarrow \mathcal{K}$ to an appropriate 2-category.

Note: A graded version, $\mathcal{U}_{q\mathfrak{g}}$, categorifies $\dot{U}_{q\mathfrak{g}}$.

The Rickard complex I

$\mathfrak{g} = \mathfrak{sl}_2$: Let $s = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} = \exp(-f)\exp(e)\exp(-f) \in SL_2$.

Then s restricts to an isomorphism on weight spaces, of the form

$$s|_{V_{-n}} = \sum_k (-1)^k e^{(n+k)} f^{(k)}$$

Rickard complex: Consider the complex of functors $\Theta = \bigoplus_n \Theta_n$,

$$\Theta_n : \text{Comp}(C_{-n}) \rightarrow \text{Comp}(C_n)$$

$$\Theta_n = (\dots \rightarrow E^{(n+2)}F^{(2)} \rightarrow E^{(n+1)}F^{(1)} \rightarrow \underline{E}^{(n)})$$

- $E^{(n)} \subseteq E^n, F^{(n)} \subseteq F^n$ defined using the H_n -action
- $E^{(n+k)}F^{(k)} \rightarrow E^{(n+k-1)}F^{(k-1)}$ comes from adjunction

The Rickard complex II

Theorem (Chuang-Rouquier '08)

Θ induces a self-equivalence on $D^b(C)$ and, by restriction, an equivalence $D^b(C_{-n}) \xrightarrow{\cong} D^b(C_n)$. Furthermore, $[\Theta] = s$.

Example: Let $R = \mathbb{C}[x]/x^2$. For the adjoint $s\mathbb{I}_2$ -representation and $N \in \mathcal{C}_0 = R\text{-mod}$,

$$\begin{aligned}\Theta_0 : D^b(R\text{-mod}) &\rightarrow D^b(R\text{-mod}) \\ N &\mapsto (R \otimes N \xrightarrow{\text{act}} \underline{N})\end{aligned}$$

Perverse equivalences

Suppose that $\mathcal{A}, \mathcal{A}'$ are abelian categories, $F : D^b(\mathcal{A}) \xrightarrow{\cong} D^b(\mathcal{A}')$, we have filtrations $\mathcal{A}_\bullet, \mathcal{A}'_\bullet$ by Serre subcategories

$$0 = \mathcal{A}_{-1} \subset \mathcal{A}_0 \subset \mathcal{A}_1 \subset \dots \subset \mathcal{A}_r = \mathcal{A}$$

$$0 = \mathcal{A}'_{-1} \subset \mathcal{A}'_0 \subset \mathcal{A}'_1 \subset \dots \subset \mathcal{A}'_r = \mathcal{A}'$$

and a perversity function $p : \{0, \dots, r\} \rightarrow \mathbb{Z}$.

Definition

F is **perverse** with respect to $(\mathcal{A}_\bullet, \mathcal{A}'_\bullet, p)$ if:

- 1 $F[-p(i)]$ restricts to an equivalence $D_{\mathcal{A}_i}^b(\mathcal{A}) \xrightarrow{\cong} D_{\mathcal{A}'_i}^b(\mathcal{A}')$.
- 2 The induced $D_{\mathcal{A}_i}^b(\mathcal{A})/D_{\mathcal{A}_{i-1}}^b(\mathcal{A}) \xrightarrow{\cong} D_{\mathcal{A}'_i}^b(\mathcal{A}')/D_{\mathcal{A}'_{i-1}}^b(\mathcal{A}')$ equivalence induces an equivalence

$$\mathcal{A}_i/\mathcal{A}_{i-1} \xrightarrow{\cong} \mathcal{A}'_i/\mathcal{A}'_{i-1}.$$

Perversity of the Rickard complexes

Consider C (all of whose objects have finite composition series) endowed with an \mathfrak{sl}_2 -categorical action. Let S be the set of simple objects, and consider the filtrations:

$$S_i = \{V \in S : F^{i+1} S = 0\} \quad \text{and} \quad S'_i = \{V \in S : E^{i+1} V = 0\}.$$

Proposition (Chuang–Rouquier)

The equivalence $\Theta : D^b(C) \xrightarrow{\cong} D^b(C)$ is perverse with respect to $(S_\bullet, S'_\bullet, p = \text{Id})$.

For a reduced word $w = s_{i_1} \dots s_{i_k}$ and weight μ , consider the composition $\Theta_\mu^w = \Theta_{s_{i_2} \dots s_{i_r}(\mu)}^{s_{i_1}} \circ \dots \circ \Theta_\mu^{s_{i_k}}$.

Theorem 1 (H–Licata–Losev–Yacobi)

Let $w_0 \in W$ be the longest element, and μ a weight of C . Then $\Theta_\mu^{w_0} : D^b(C_\mu) \rightarrow D^b(C_{w_0(\mu)})$ is a perverse equivalence.

Braid and cactus groups

The **braid group** $B_{\mathfrak{g}}$ is generated by $\sigma_i, i \in I$, with relations:

$$\begin{aligned}\sigma_i \sigma_j \sigma_i &= \sigma_j \sigma_i \sigma_j && \text{if } i \text{ and } j \text{ are connected,} \\ \sigma_i \sigma_j &= \sigma_j \sigma_i && \text{if } i \text{ and } j \text{ are not connected.}\end{aligned}$$

The **cactus group** $C_{\mathfrak{g}}$ is generated by $c_J, J \subseteq I$, with relations:


$$\begin{aligned}c_J^2 &= 1 && \forall J \subset I \\ c_J c_K &= c_K c_J && \forall J \cup K \subset I \text{ not connected} \\ c_J c_K &= c_{*J(K)} c_J && \forall K \subset J \subset I\end{aligned}$$

Where $\forall j \in J, \alpha_j$ simple root, $\alpha_{*J(j)} = -w_0^J \alpha_j$.

Proposition (Cautis–Kamnitzer '10)

The Rickard complexes satisfy the braid relations.

A cactus group action



Theorem 2 (H-Licata-Losev-Yacobi)

Let $\theta_J : \text{Irr}(C) \rightarrow \text{Irr}(C)$ denote the bijection induced from $\Theta_{w_0^J}$.
 Then the map $c_J \mapsto \theta_J$ defines a cactus group action $C_{\mathfrak{g}} \curvearrowright \text{Irr}(C)$
 which coincides with the combinatorial action of the cactus group
 on crystals (via Schützenberger involutions).

The End

Thank you!