

Matroid base polytope decomposition

J.L. Ramírez Alfonsín

(join work with V. Chatelain)

I3M, Université Montpellier 2

Definitions

A matroid $M = (E, \mathcal{I})$ is a finite ground set $E = \{1, \dots, n\}$ together with a collection $\mathcal{I} \subseteq 2^E$, known as independent sets satisfying

- if $I \in \mathcal{I}$ and $J \subseteq I$ then $J \in \mathcal{I}$
- if $I, J \in \mathcal{I}$ and $|J| > |I|$, then there exist an element $z \in J \setminus I$ such that $I \cup \{z\} \in \mathcal{I}$.

A base is any maximal independent set. The collection of bases \mathcal{B} satisfy the base exchange axiom

if $B_1, B_2 \in \mathcal{B}$ and $e \in B_1 \setminus B_2$ then there exist $f \in B_2 \setminus B_1$ such that $(B_1 - e) + f \in \mathcal{B}$.

Remark : All bases have the same cardinality, say r . We say that matroid $M = (E, \mathcal{B})$ has rank $r = r(M)$.

Definitions

A **matroid** $M = (E, \mathcal{I})$ is a finite ground set $E = \{1, \dots, n\}$ together with a collection $\mathcal{I} \subseteq 2^E$, known as **independent sets** satisfying

- if $I \in \mathcal{I}$ and $J \subseteq I$ then $J \in \mathcal{I}$
- if $I, J \in \mathcal{I}$ and $|J| > |I|$, then there exist an element $z \in J \setminus I$ such that $I \cup \{z\} \in \mathcal{I}$.

A **base** is any maximal independent set. The collection of bases \mathcal{B} satisfy the **base exchange axiom**

if $B_1, B_2 \in \mathcal{B}$ and $e \in B_1 \setminus B_2$ then there exist $f \in B_2 \setminus B_1$ such that $(B_1 - e) + f \in \mathcal{B}$.

Remark : All bases have the same cardinality, say r . We say that matroid $M = (E, \mathcal{B})$ has **rank** $r = r(M)$.

Definitions

A **matroid** $M = (E, \mathcal{I})$ is a finite ground set $E = \{1, \dots, n\}$ together with a collection $\mathcal{I} \subseteq 2^E$, known as **independent sets** satisfying

- if $I \in \mathcal{I}$ and $J \subseteq I$ then $J \in \mathcal{I}$
- if $I, J \in \mathcal{I}$ and $|J| > |I|$, then there exist an element $z \in J \setminus I$ such that $I \cup \{z\} \in \mathcal{I}$.

A **base** is any maximal independent set. The collection of bases \mathcal{B} satisfy the **base exchange axiom**

if $B_1, B_2 \in \mathcal{B}$ and $e \in B_1 \setminus B_2$ then there exist $f \in B_2 \setminus B_1$ such that $(B_1 - e) + f \in \mathcal{B}$.

Remark : All bases have the same cardinality, say r . We say that matroid $M = (E, \mathcal{B})$ has **rank** $r = r(M)$.

Examples

- Uniform matroids $U_{n,r}$ given by $E = \{1, \dots, n\}$ and $\mathcal{I} = \{I \subseteq E : |I| \leq r\}$.

- Linear matroids Let \mathbb{F} be a field, $A \in \mathbb{F}^{m \times n}$ an $(m \times n)$ -matrix over \mathbb{F} . Let $E = \{1, \dots, n\}$ be the index set of the columns of A . $I \subseteq E$ is independent if the columns indexed by I are linearly independent.

A matroid is said to be representable over \mathbb{F} if it can be expressed as linear matroid with matrix A and linear independence taken over \mathbb{F} .

- Graphic matroid Let $G = (V, E)$ be an undirected graph. Matroid $M = (E, \mathcal{I})$ where $\mathcal{I} = \{F \subseteq E : F \text{ is acyclic}\}$.

Examples

- Uniform matroids $U_{n,r}$ given by $E = \{1, \dots, n\}$ and $\mathcal{I} = \{I \subseteq E : |I| \leq r\}$.
- Linear matroids Let \mathbb{F} be a field, $A \in \mathbb{F}^{m \times n}$ an $(m \times n)$ -matrix over \mathbb{F} . Let $E = \{1, \dots, n\}$ be the index set of the columns of A . $I \subseteq E$ is independent if the columns indexed by I are linearly independent.

A matroid is said to be representable over \mathbb{F} if it can be expressed as linear matroid with matrix A and linear independence taken over \mathbb{F} .

- Graphic matroid Let $G = (V, E)$ be an undirected graph. Matroid $M = (E, \mathcal{I})$ where $\mathcal{I} = \{F \subseteq E : F \text{ is acyclic}\}$.

Examples

- Uniform matroids $U_{n,r}$ given by $E = \{1, \dots, n\}$ and $\mathcal{I} = \{I \subseteq E : |I| \leq r\}$.
- Linear matroids Let \mathbb{F} be a field, $A \in \mathbb{F}^{m \times n}$ an $(m \times n)$ -matrix over \mathbb{F} . Let $E = \{1, \dots, n\}$ be the index set of the columns of A . $I \subseteq E$ is independent if the columns indexed by I are linearly independent.

A matroid is said to be **representable** over \mathbb{F} if it can be expressed as linear matroid with matrix A and linear independence taken over \mathbb{F} .

- Graphic matroid Let $G = (V, E)$ be an undirected graph. Matroid $M = (E, \mathcal{I})$ where $\mathcal{I} = \{F \subseteq E : F \text{ is acyclic}\}$.

Examples

- Uniform matroids $U_{n,r}$ given by $E = \{1, \dots, n\}$ and $\mathcal{I} = \{I \subseteq E : |I| \leq r\}$.
- Linear matroids Let \mathbb{F} be a field, $A \in \mathbb{F}^{m \times n}$ an $(m \times n)$ -matrix over \mathbb{F} . Let $E = \{1, \dots, n\}$ be the index set of the columns of A . $I \subseteq E$ is independent if the columns indexed by I are linearly independent.

A matroid is said to be **representable** over \mathbb{F} if it can be expressed as linear matroid with matrix A and linear independence taken over \mathbb{F} .

- Graphic matroid Let $G = (V, E)$ be an undirected graph. Matroid $M = (E, \mathcal{I})$ where $\mathcal{I} = \{F \subseteq E : F \text{ is acyclic}\}$.

Applications

- Graph theory
- Combinatorial optimisation (via greedy characterization)
- Knot theory (Jones's polynomial)
- Hyperplane arrangements (via oriented matroids)
- Polytopes (tilings, convexity, Ehrhart polynomial)
- Rigidity
- Cryptography (secret sharing)

Applications

- Graph theory
- Combinatorial optimisation (via greedy characterization)
- Knot theory (Jones's polynomial)
- Hyperplane arrangements (via oriented matroids)
- Polytopes (tilings, convexity, Ehrhart polynomial)
- Rigidity
- Cryptography (secret sharing)

Let $P(M)$ the **base polytope** of M defined as the convex hull of the incidence vectors of bases of M , that is,

$$P(M) := \text{conv} \left\{ \sum_{i \in B} e_i : B \in \mathcal{B}(M) \right\}$$

where e_i denote the standard unit vector of \mathbb{R}^n
(these polytopes were first studied by J. Edmonds in the seventies).

Remark

- (a) $P(M)$ is a polytope of dimension at most $n - 1$.
- (b) $P(M)$ is a face of the **indepent polytope** of M defined as the convex hull of the incidence vectors of the independent sets of M .

Let $P(M)$ the **base polytope** of M defined as the convex hull of the incidence vectors of bases of M , that is,

$$P(M) := \text{conv} \left\{ \sum_{i \in B} e_i : B \in \mathcal{B}(M) \right\}$$

where e_i denote the standard unit vector of \mathbb{R}^n


(these polytopes were first studied by J. Edmonds in the seventies).

Remark

(a) $P(M)$ is a polytope of dimension at most $n - 1$.

(b) $P(M)$ is a face of the **indepent polytope** of M defined as the convex hull of the incidence vectors of the independent sets of M .

Exemple : $P(U_{4,2})$



A decomposition of $P(M)$ is a decomposition

$$P(M) = \bigcup_{i=1}^t P(M_i)$$

where each $P(M_i)$ is also a base matroid polytope for some M_i , and for each $1 \leq i \neq j \leq t$, the intersection $P(M_i) \cap P(M_j)$ is a face of both $P(M_i)$ and $P(M_j)$.

$P(M)$ is said decomposable if it has a matroid base polytope decomposition with $t \geq 2$, and indecomposable otherwise.

A decomposition is called hyperplane split if $t = 2$.

A decomposition of $P(M)$ is a decomposition

$$P(M) = \bigcup_{i=1}^t P(M_i)$$

where each $P(M_i)$ is also a base matroid polytope for some M_i , and for each $1 \leq i \neq j \leq t$, the intersection $P(M_i) \cap P(M_j)$ is a face of both $P(M_i)$ and $P(M_j)$.

$P(M)$ is said **decomposable** if it has a matroid base polytope decomposition with $t \geq 2$, and **indecomposable** otherwise.

A decomposition is called **hyperplane split** if $t = 2$.

Applications

(L. Lafforgue - Fields medal 2002) General method of compactification of the fine Schubert cell of the Grassmannian. It is proved that such compactification exists if the $P(M)$ is indecomposable.

Remark : Lafforgue's work implies that for a matroid M represented by vectors in \mathbb{F}^r , if $P(M)$ is indecomposable, then M will be **rigid**, that is, M will have only finitely many realizations, up to scaling and the action of $GL(r, \mathbb{F})$.

(Hacking, Keel and Tevelev) Compactification of the moduli space of hyperplane arrangements

(Speyer) Tropical linear spaces

(Ardila, Fink and Rincon) There exist matroid functions behave like *valuations* on the associated matroid base polytope decomposition.

Applications

(L. Lafforgue - Fields medal 2002) General method of compactification of the fine Schubert cell of the Grassmannian. It is proved that such compactification exists if the $P(M)$ is indecomposable.

Remark : Lafforgue's work implies that for a matroid M represented by vectors in \mathbb{F}^r , if $P(M)$ is indecomposable, then M will be rigid, that is, M will have only finitely many realizations, up to scaling and the action of $GL(r, \mathbb{F})$.

(Hacking, Keel and Tevelev) Compactification of the moduli space of hyperplane arrangements

(Speyer) Tropical linear spaces

(Ardila, Fink and Rincon) There exist matroid functions behave like *valuations* on the associated matroid base polytope decomposition.

Known results

Theorem (Kapranov 1993) Any decomposition of a rank 2 matroid can be achieved by a sequence of hyperplane splits.

Theorem (Billera, Jia and Reiner 2009)

- Found five rank 3 matroids on 6 elements for which the corresponding polytopes are indecomposable.
- Gave a rank 3 matroid on 6 elements having a 3-decomposition but cannot be obtained via hyperplane splits.

Known results

Theorem (Kapranov 1993) Any decomposition of a rank 2 matroid can be achieved by a sequence of hyperplane splits.

Theorem (Billera, Jia and Reiner 2009)

- Found five rank 3 matroids on 6 elements for which the corresponding polytopes are indecomposable.
- Gave a rank 3 matroid on 6 elements having a 3-decomposition but cannot be obtained via hyperplane splits.

Combinatorial decomposition

A base decomposition of a matroid M is a decomposition

$$\mathcal{B}(M) = \bigcup_{i=1}^t \mathcal{B}(M_i)$$

where $\mathcal{B}(M_k)$, $1 \leq k \leq t$ and $\mathcal{B}(M_i) \cap \mathcal{B}(M_j)$, $1 \leq i \neq j \leq t$ are collections of bases of matroids.

M is called *combinatorial decomposable* if it has a base decomposition.

A decomposition is *nontrivial* if $\mathcal{B}(M_i) \neq \mathcal{B}(M)$ for all i .

Combinatorial decomposition

A base decomposition of a matroid M is a decomposition

$$\mathcal{B}(M) = \bigcup_{i=1}^t \mathcal{B}(M_i)$$

where $\mathcal{B}(M_k)$, $1 \leq k \leq t$ and $\mathcal{B}(M_i) \cap \mathcal{B}(M_j)$, $1 \leq i \neq j \leq t$ are collections of bases of matroids.

M is called **combinatorial decomposable** if it has a base decomposition.

A decomposition is *nontrivial* if $\mathcal{B}(M_i) \neq \mathcal{B}(M)$ for all i .

- If $P(M)$ is decomposable then M is clearly combinatorial decomposable.

- A combinatorial decomposition of M could not yield to a decomposition of $P(M)$.

Example

$\mathcal{B}(M) = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{2, 4\}, \{3, 4\}\}$ has the following combinatorial decomposition

$\mathcal{B}(M_1) = \{\{1, 2\}, \{2, 3\}, \{2, 4\}\}$ and

$\mathcal{B}(M_2) = \{\{1, 3\}, \{2, 3\}, \{3, 4\}\}$

We verify that $\mathcal{B}(M_1), \mathcal{B}(M_2)$ and $\mathcal{B}(M_1) \cap \mathcal{B}(M_2) = \{2, 3\}$ are collections of bases of matroids.

However, $P(M_1)$ and $P(M_2)$ do not decompose $P(M)$.

- If $P(M)$ is decomposable then M is clearly combinatorial decomposable.
- A combinatorial decomposition of M could not yield to a decomposition of $P(M)$.


Example

$\mathcal{B}(M) = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{2, 4\}, \{3, 4\}\}$ has the following combinatorial decomposition

$$\mathcal{B}(M_1) = \{\{1, 2\}, \{2, 3\}, \{2, 4\}\} \text{ and}$$
$$\mathcal{B}(M_2) = \{\{1, 3\}, \{2, 3\}, \{3, 4\}\}$$

We verify that $\mathcal{B}(M_1), \mathcal{B}(M_2)$ and $\mathcal{B}(M_1) \cap \mathcal{B}(M_2) = \{2, 3\}$ are collections of bases of matroids.

However, $P(M_1)$ and $P(M_2)$ do not decompose $P(M)$.



Proposition Let P be d -polytope with set of vertices X . Let H be a hyperplane such that $F = H \cap P \neq \emptyset$ and non-supporting P . So, H partition X into X_1 and X_2 with $X_1 \cap X_2 = W$. Then, for each edge $[u, v]$ of P we have that $\{u, v\} \subset X_i$ for either $i = 1$ or 2 if and only if $F = \text{conv}(W)$.

Corollary Suppose that H divides P in two polytopes P_1 and P_2 . Then, $F = \text{conv}(W)$ if and only if $P_i = \text{conv}(X_i)$, $i = 1, 2$.

Proposition Let P be d -polytope with set of vertices X . Let H be a hyperplane such that $F = H \cap P \neq \emptyset$ and non-supporting P . So, H partition X into X_1 and X_2 with $X_1 \cap X_2 = W$. Then, for each edge $[u, v]$ of P we have that $\{u, v\} \subset X_i$ for either $i = 1$ or 2 if and only if $F = \text{conv}(W)$.

Corollary Suppose that H divides P in two polytopes P_1 and P_2 . Then, $F = \text{conv}(W)$ if and only if $P_i = \text{conv}(X_i)$, $i = 1, 2$.

Let (E_1, E_2) be a partition of E and let $r_i > 1$, $i = 1, 2$ be the rank of $M|_{E_i}$. We say that (E_1, E_2) is a **good partition** if there exist integers $0 < a_1 < r_1$ et $0 < a_2 < r_2$ such that :

(P1) $r_1 + r_2 = r + a_1 + a_2$ and

(P2) for any $X \in \mathcal{I}(M|_{E_1})$ with $|X| \leq r_1 - a_1$ and
for any $Y \in \mathcal{I}(M|_{E_2})$ with $|Y| \leq r_2 - a_2$
we have that $X \cup Y \in \mathcal{I}(M)$.

Lemma Let (E_1, E_2) be a good partition of E . Let

$$\mathcal{B}(M_1) = \{B \in \mathcal{B}(M) : |B \cap E_1| \leq r_1 - a_1\}$$

$$\mathcal{B}(M_2) = \{B \in \mathcal{B}(M) : |B \cap E_2| \leq r_2 - a_2\}$$

where r_i is the rank $M|_{E_i}$, $i = 1, 2$ and a_1, a_2 verify (P1) and (P2).

Then, $\mathcal{B}(M_1)$ and $\mathcal{B}(M_2)$ are collections of bases of matroids, say M_1 et M_2 .

Let (E_1, E_2) be a partition of E and let $r_i > 1$, $i = 1, 2$ be the rank of $M|_{E_i}$. We say that (E_1, E_2) is a **good partition** if there exist integers $0 < a_1 < r_1$ et $0 < a_2 < r_2$ such that :

(P1) $r_1 + r_2 = r + a_1 + a_2$ and

(P2) for any $X \in \mathcal{I}(M|_{E_1})$ with $|X| \leq r_1 - a_1$ and
for any $Y \in \mathcal{I}(M|_{E_2})$ with $|Y| \leq r_2 - a_2$
we have that $X \cup Y \in \mathcal{I}(M)$.

Lemma Let (E_1, E_2) be a good partition of E . Let

$$\mathcal{B}(M_1) = \{B \in \mathcal{B}(M) : |B \cap E_1| \leq r_1 - a_1\}$$

$$\mathcal{B}(M_2) = \{B \in \mathcal{B}(M) : |B \cap E_2| \leq r_2 - a_2\}$$

where r_i is the rank $M|_{E_i}$, $i = 1, 2$ and a_1, a_2 verify (P1) and (P2).

Then, $\mathcal{B}(M_1)$ and $\mathcal{B}(M_2)$ are collections of bases of matroids, say M_1 et M_2 .

Theorem (Chatelain and R.A. 2011) Let $M = (E, \mathcal{B})$ be a matroid and let (E_1, E_2) a good partition of E . Then, $P(M) = P(M_1) \cup P(M_2)$ is a nontrivial hyperplane split where M_1 et M_2 are the matroids of lemma above.

We say that two hyperplane splits $P(M_1) \cup P(M_2)$ and $P(M'_1) \cup P(M'_2)$ of $P(M)$ are equivalent if $P(M_i)$ is combinatorially equivalent to $P(M'_i)$, $i = 1, 2$. They are different otherwise.

Corollary (Chatelain and R.A. 2011) Let $n \geq r + 2 \geq 4$ be integers and let $h(U_{n,r})$ be the number of different hyperplane splits of $P(U_{n,r})$. Then,

$$h(U_{n,r}) \geq \left\lfloor \frac{n}{2} \right\rfloor - 1.$$

Theorem (Chatelain and R.A. 2011) Let $M = (E, \mathcal{B})$ be a matroid and let (E_1, E_2) a good partition of E . Then, $P(M) = P(M_1) \cup P(M_2)$ is a nontrivial hyperplane split where M_1 et M_2 are the matroids of lemma above.

We say that two hyperplane splits $P(M_1) \cup P(M_2)$ and $P(M'_1) \cup P(M'_2)$ of $P(M)$ are **equivalent** if $P(M_i)$ is **combinatorially equivalent** to $P(M'_i)$, $i = 1, 2$. They are **different** otherwise.

Corollary (Chatelain and R.A. 2011) Let $n \geq r + 2 \geq 4$ be integers and let $h(U_{n,r})$ be the number of different hyperplane splits of $P(U_{n,r})$. Then,

$$h(U_{n,r}) \geq \left\lfloor \frac{n}{2} \right\rfloor - 1.$$

Theorem (Chatelain and R.A. 2011) Let $M = (E, \mathcal{B})$ be a matroid and let (E_1, E_2) a good partition of E . Then, $P(M) = P(M_1) \cup P(M_2)$ is a nontrivial hyperplane split where M_1 et M_2 are the matroids of lemma above.

We say that two hyperplane splits $P(M_1) \cup P(M_2)$ and $P(M'_1) \cup P(M'_2)$ of $P(M)$ are equivalent if $P(M_i)$ is combinatorially equivalent to $P(M'_i)$, $i = 1, 2$. They are different otherwise.

Corollary (Chatelain and R.A. 2011) Let $n \geq r + 2 \geq 4$ be integers and let $h(U_{n,r})$ be the number of different hyperplane splits of $P(U_{n,r})$. Then,

$$h(U_{n,r}) \geq \left\lfloor \frac{n}{2} \right\rfloor - 1.$$

Example Let us consider $U_{4,2}$. Then, $E_1 = \{1, 2\}$ and $E_2 = \{3, 4\}$ is a good partition (and thus $r_1 = r_2 = 2$) with $a_1 = a_2 = 1$.

We have $\mathcal{B}(M_1) = \{\{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}\}$,

$\mathcal{B}(M_2) = \{\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}\}$ and


$\mathcal{B}(M_1) \cap \mathcal{B}(M_2) = \{\{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}\}$.

Example Let us consider $U_{4,2}$. Then, $E_1 = \{1, 2\}$ and $E_2 = \{3, 4\}$ is a good partition (and thus $r_1 = r_2 = 2$) with $a_1 = a_2 = 1$.

We have $\mathcal{B}(M_1) = \{\{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}, \{3, 4\}\}$,

$\mathcal{B}(M_2) = \{\{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}\}$ and

$\mathcal{B}(M_1) \cap \mathcal{B}(M_2) = \{\{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}\}$.




Lattice path matroids


Let $m = 3$ and $r = 4$ and let $M[Q, P]$ be the transversal matroid on $\{1, \dots, 7\}$ with presentation $(N_i : i \in \{1, \dots, 4\})$ where $N_1 = [1, 2, 3, 4]$, $N_2 = [3, 4, 5]$, $N_3 = [5, 6]$ and $N_4 = [7]$.

Lattice path matroids


Let $m = 3$ and $r = 4$ and let $M[Q, P]$ be the transversal matroid on $\{1, \dots, 7\}$ with presentation $(N_i : i \in \{1, \dots, 4\})$ where $N_1 = [1, 2, 3, 4]$, $N_2 = [3, 4, 5]$, $N_3 = [5, 6]$ and $N_4 = [7]$.




Example Transversal matroids (a) M_1 , (b) M_2 et (c) $M_1 \cap M_2$.



(a)



(b)



(c)

Theorem (Chatelain and R.A. 2011) Let $M_1 = (E_1, \mathcal{B})$ and $M_2 = (E_2, \mathcal{B})$ be two matroids of ranks r_1 and r_2 respectively where $E_1 \cap E_2 = \emptyset$. Then, $P(M_1 \oplus M_2)$ has a nontrivial hyperplane split if and only if either $P(M_1)$ or $P(M_2)$ has a nontrivial hyperplane split.


Remark The class of lattice path matroids are closed under direct sum.

Theorem (Chatelain and R.A. 2011) Let $M_1 = (E_1, \mathcal{B})$ and $M_2 = (E_2, \mathcal{B})$ be two matroids of ranks r_1 and r_2 respectively where $E_1 \cap E_2 = \emptyset$. Then, $P(M_1 \oplus M_2)$ has a nontrivial hyperplane split if and only if either $P(M_1)$ or $P(M_2)$ has a nontrivial hyperplane split.

Remark The class of lattice path matroids are closed under direct sum.

Theorem (Chatelain and R.A. 2011) Let $M_1 = (E_1, \mathcal{B})$ and $M_2 = (E_2, \mathcal{B})$ be two matroids of ranks r_1 and r_2 respectively where $E_1 \cap E_2 = \emptyset$. Then, $P(M_1 \oplus M_2)$ has a nontrivial hyperplane split if and only if either $P(M_1)$ or $P(M_2)$ has a nontrivial hyperplane split.

Remark The class of lattice path matroids are closed under direct sum.



Binary matroids

A matroid is called **binary** if it is representable over \mathbb{F}_2 .

Let $G(M)$ be the **base graph** of a matroid M ($G(M)$ is the 1-skeleton of $P(M)$).


Theorem (Maurer 1976) If x, y are two vertices at distance two then the neighbours of x and y form either a square or a pyramid or an octahedron.

Binary matroids

A matroid is called **binary** if it is representable over \mathbb{F}_2 .

Let $G(M)$ be the **base graph** of a matroid M ($G(M)$ is the 1-skeleton of $P(M)$).

Theorem (Maurer 1976) If x, y are two vertices at distance two then the neighbours of x and y form either a square or a pyramid or an octahedron.



Lemma Let $M = (E, \mathcal{B})$ a binary matroid and let $\mathcal{B}_1 \subset \mathcal{B}$ such that \mathcal{B}_1 is the collection of bases of a matroid, says M_1 . If $X \in \mathcal{B}_1$ and all the neighbours of X are elements of \mathcal{B}_1 then $\mathcal{B}_1 = \mathcal{B}$.

Theorem (Chatelain and R.A. 2011) Let M be a binary matroid. Then, $P(M)$ do not have a nontrivial hyperplane split.

Lemma Let $M = (E, \mathcal{B})$ a binary matroid and let $\mathcal{B}_1 \subset \mathcal{B}$ such that \mathcal{B}_1 is the collection of bases of a matroid, says M_1 . If $X \in \mathcal{B}_1$ and all the neighbours of X are elements of \mathcal{B}_1 then $\mathcal{B}_1 = \mathcal{B}$.

Theorem (Chatelain and R.A. 2011) Let M be a binary matroid. Then, $P(M)$ do not have a nontrivial hyperplane split.

Corollary Let M be a binary matroid. If $G(M)$ contains a vertex X having exactly d neighbours where $d = \dim(P(M))$ then $P(M)$ is indecomposable.

Remark The d -dimensional hypercube is the base graph of a binary matroid.

Corollary (Chatelain and R.A. 2011) Let $P(M)$ be the base matroid polytope of a matroid M having as 1-skeleton the hypercube. Then, $P(M)$ is indecomposable.

Corollary Let M be a binary matroid. If $G(M)$ contains a vertex X having exactly d neighbours where $d = \dim(P(M))$ then $P(M)$ is indecomposable.

Remark The d -dimensional hypercube is the base graph of a binary matroid.

Corollary (Chatelain and R.A. 2011) Let $P(M)$ be the base matroid polytope of a matroid M having as 1-skeleton the hypercube. Then, $P(M)$ is indecomposable.

Corollary Let M be a binary matroid. If $G(M)$ contains a vertex X having exactly d neighbours where $d = \dim(P(M))$ then $P(M)$ is indecomposable.

Remark The d -dimensional hypercube is the base graph of a binary matroid.

Corollary (Chatelain and R.A. 2011) Let $P(M)$ be the base matroid polytope of a matroid M having as 1-skeleton the hypercube. Then, $P(M)$ is indecomposable.